

2017

RAPPORT

EN SYSTEMATISK OVERSIKT

Samsvar mellom
kronologisk alder og
skjelettalder basert på
Greulich og Pyle-atlaset
for aldersestimering

Utgitt av	Folkehelseinstituttet Avdeling for kunnskapsoppsummering i Kunnskapssenteret
Titel	Samsvar mellom kronologisk alder og skjelettalder basert på Greulich og Pyle-atlaset for alderestimering: en systematisk oversikt
English title	Agreement between chronological age and bone age based on the Greulich and Pyle atlas for age estimation: a systematic review
Ansvarlig	Camilla Stoltenberg, direktør
Forfattere	Pål Skage Dahlberg, prosjektleader, Oslo universitetssykehus Annhild Mosdøl, Folkehelseinstituttet Kristoffer Yunpeng Ding, Folkehelseinstituttet Øyvind Bleka, Oslo universitetssykehus Veslemøy Rolseth, Oslo universitetssykehus Gyri Hval Straumann, Folkehelseinstituttet Marianne Skjerven-Martinsen, Oslo universitetssykehus Gerd Jorunn Møller Delaveris, Oslo universitetssykehus Gunn Elisabeth Vist, Folkehelseinstituttet
ISBN	978-82-8082-822-4
Publikasjonstype	Systematisk oversikt
Antall sider	46 (120 inklusiv vedlegg)
Oppdragsgiver	Folkehelseinstituttet
Emneord(MeSH)	Age Determination by Skeleton
Sitering	Dahlberg PS, Mosdøl A, Ding KY, Bleka Ø, Rolseth V, Straumann GH, Skjerven-Martinsen M, Delaveris GJM, Vist GE. Samsvar mellom kronologisk alder og skjelettalder basert på Greulich og Pyle-atlaset for alderestimering: en systematisk oversikt. [Agreement between chronological age and bone age based on the Greulich and Pyle-atlas for age estimation: a systematic review] Rapport 2017. Oslo: Folkehelseinstituttet, 2017.

Innhold

INNHOLD	3
HOVEDBUDSKAP	5
SAMMENDRAG	6
KEY MESSAGES (ENGLISH)	10
EXECUTIVE SUMMARY (ENGLISH)	11
FORORD	15
INNLEDNING	16
Beskrivelse av Greulich og Pyle-atlaset	17
Bruken av Greulich og Pyle-atlaset	18
Valideringsstudier av Greulich og Pyle-atlaset	19
METODE	21
Inklusjonskriterier	21
Litteratursøking	21
Artikkellutvelgning	22
Kvalitetsvurdering og dataekstraksjon	22
Analyser	24
Etikk	25
RESULTATER	26
Resultater av litteratursøket	26
Beskrivelse av de inkluderte studiene	27
Andre studier vurdert fra fulltekst	28
Kvalitetsvurdering av inkluderte studier basert på QUADAS-2	28
Differansen mellom kronologisk alder og GP skjelettalder	30
Gjennomsnittlig kronologisk alder i utviklingsstadier fra Greulich og Pyle-atlaset	33
DISKUSJON	37
Hovedfunn	37
Kvaliteten på forskningsresultatene	37
Styrker og svakheter	40
Overensstemmelse med andre oversikter	41
Kunnskapshull	42
KONKLUSJON	43

REFERANSER	44
VEDLEGG (APPENDIXES)	47
Appendix 1: Literature search	47
Appendix 2: Description of included studies with quality assessment	51
Appendix 3: Relevant datasets using the Greulich and Pyle atlas	65
Appendix 4: Relevant datasets using other index tests than Greulich and Pyle atlas to classify bone age on hand x-rays	74
Appendix 5: Studies excluded after full text assessment	84
Appendix 6: All analysed results	106

Hovedbudskap

Greulich og Pyle-atlaset benyttes til å anslå alder på barn og unge. Denne systematiske oversikten sammenfatter den forskningsbaserte dokumentasjonen på samsvaret mellom skjelettalder fra Greulich og Pyle-atlaset (heretter kalt GP skjelettalder) og kronologisk alder.

Totalt 19 artikler oppfylte kriteriene våre. Femten studier fremstilte resultater for GP skjelettalder fra kronologisk alder og fire fremstilte resultater som beskriver kronologisk alder fra GP skjelettalder. Disse fremstillingene er ikke sammenlignbare, og vi har derfor utført to separate analyser.

Forskjellen mellom GP skjelettalder og kronologisk alder var sjeldent over ett år for gjennomsnittet i hver aldersgruppe.

Studiene som har beskrevet fordelingen av kronologisk alder fra GP skjelettalder er de som er mest relevante hvis Greulich og Pyle-atlaset skal anvendes til alderestimering. Her gjorde vi ingen samlet analyse, ettersom tre av fire studier var så preget av fenomenet «aldersmimikering» at resultatene ikke var pålitelige. Vi vurderte at kun en av studiene var utført på en slik måte at den gir en god nok beskrivelse av metodens evne til å anslå alder.

Tittel:

Samsvar mellom kronologisk alder og skjelettalder basert på Greulich og Pyle-atlaset for alderestimering: en systematisk oversikt

Publikasjonstype:

Systematisk oversikt

En systematisk oversikt er resultatet av å

- innhente
- kritisk vurdere og
- sammenfatte

relevante forskningsresultater ved hjelp av forhåndsdefinerte og eksplisitte metoder.

Svarer ikke på alt:

- Ingen studier utenfor de eksplisitte inklusjonskriteriene
- Ingen helseøkonomisk evaluering
- Ingen anbefalinger

Hvem står bak denne publikasjonen?

Folkehelseinstituttet

Når ble litteratursøket utført?

Søk etter studier ble avsluttet
Januar 2017.

Eksterne fagfeller:

- Bjørn Anton Graff, Klinikk for medisinsk diagnostikk, Vestre Viken HF
- Lil-Sofie Ording Müller, Oslo universitetssykehus

Sammendrag

Innledning

Hvert år kommer det unge asylsøkere til Norge som ikke vet hvor gamle de er eller ikke kan dokumentere dette. For å sikre at barn får de rettigheter de har krav på og at voksne ikke blir behandlet som barn, er det nødvendig å fastsette en kronologisk alder. I Norge har det i flere år vært benyttet evaluering av modning av skjelett i hånd og av tannutvikling for å estimere alder på asylsøkere hvor det foreligger tvil. Disse metodene har i stor grad blitt kritisert for ikke å være presise, men per dags dato er det ikke funnet bedre metoder. I 2016 fikk Folkehelseinstituttet et nasjonalt fagansvar for å evaluere og forbedre metodene (fra 1.1. 2017 overført med Avdeling for Rettsmedisinske fag til Oslo universitetssykehus). Den rettsmedisinske faggruppen har i samarbeid med Kunnskapssenteret for helsetjenesten i Folkehelseinstituttet gjennomført en systematisk kartlegging av det vitenskapelige grunnlaget for flere metoder som benyttes til medisinske aldersvurderinger.

Formålet med denne systematiske oversikten er å sammenfatte den forskningsbaserte dokumentasjonen om samsvaret mellom kronologisk alder og skjeletalder basert på modningsstadier fra Greulich og Pyle-atlaset (GP skjeletalder), og eventuelt å belyse variasjoner mellom ulike populasjoner. Siden Greulich og Pyle-atlaset er den mest anvendte metoden for estimering av kronologisk alder basert på håndrøntgen, valgte vi å fokusere analysen på dette systemet. Parallelt har vi også gjennomført en systematisk oversikt på aldersestimering med Demirjians utviklingsstadier av visdomstener.

Metode

Vi søkte etter studier i Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, Embase og Google Scholar. Søket ble avsluttet mai 2016. Det ble utført ett felles søk for studier som benyttet røntgen av tenner eller hånd, samt CT eller MR av kragben, kne og ankel til aldersvurdering for barn og unge mellom 10 og 25 år. Søket for studier som benyttet Greulich og Pyle-atlaset på håndrøntgen ble oppdatert i januar 2017. To personer leste uavhengig av hverandre tittel og sammendrag for 10640 referanser. Av disse fant vi 658 publikasjoner som kunne være relevante for aldersestimering basert på håndrøntgen. To personer vurderte disse uavhengig av hverandre i fulltekst. Vi inkluderte 19 studier som har estimert alder basert på Greulich og Pyle-atlaset og samtidig har oppgitt data for hvert kronologiske år. Vi har inkludert to ulike måter å fremstille dataene på (i denne rapporten kalt fremstilling A og B). Begge metodene forutsetter en kjent kronologisk alder og en observert skjeletalder:

A) tar utgangspunkt i kronologisk alder og fremstiller gjennomsnittlig forskjell mellom kronologisk alder og GP skjeletalder basert på modningsstadier i Greulich og Pyle atlaset for aldersgrupper separat (ikke slått sammen aldre i mer enn årlige intervaller).

B) tar utgangspunkt i skjeletalder og fremstiller gjennomsnittlig kronologisk alder fra GP skjeletalder i Greulich og Pyle-atlaset.

To personer vurderte uavhengig av hverandre risiko for systematiske skjevheter i de inkluderte studiene ved QUADAS-2 sjekklisten for å vurdere kvaliteten av studier av diagnostisk nøyaktighet.

Resultat

Vi fant 19 studier som hadde sammenlignet kronologisk alder med GP skjeletalder basert på håndrøntgen og GP-atlaset. Studiene inkluderte fra 68 til 2614 personer med kjent kronologisk alder. Alle unntatt én studie inkluderte både gutter og jenter. Det var fem studier fra Tyrkia, fire studier fra India, og én studie hver fra henholdsvis Canada, Frankrike, Italia, Iran, Kina, Nederland, Spania, Taiwan, Pakistan og USA. Vi vurderte at flertallet av studiene hadde enten lav eller uklar risiko for systematiske skjevheter basert på QUADAS sjekklisten. Ett unntak var studiene med fremstilling B, hvor tre av fire studier hadde resultater med høy risiko for en spesiell form for seleksjonsskjehet kalt aldersmimikering.

Det klare flertall av studiene har oppgitt resultatene som gjennomsnittlig forskjell mellom kronologisk alder og GP skjeletalder for aldersgrupper basert på modningsstadier i Greulich og Pyle-atlaset (fremstilling A). Analysen viser at variasjonen mellom studiene var større enn man kan forvente ved tilfeldighet, men samtidig var det relativt moderate variasjoner mellom studier fra ulike deler av verden. Det var sjeldent at differansen mellom GP skjeletalder og kronologisk alder oversteg ett år i gjennomsnitt for en gruppe.

Fire studier, inkludert to vi analyserte basert på forfatternes rådata, fremstilte resultatene som gjennomsnittlig kronologisk alder i modningsstadier fra Greulich og Pyle-atlaset (fremstilling B). Vi fant at resultatene fra tre av disse studiene i høy grad var påvirket av alderssammensetningen av testpersonene. Denne skjeheten er tidligere beskrevet som aldersmimikering (*engelsk: age mimicry*). Aldersmimikering fører til at gjennomsnittsalder for hvert utviklingsstadium preges av alderssammensetningen i den inkluderte studiepopulasjonen. Kun én av de fire studiene med denne fremstillingsmåten hadde en stor inkludert populasjon som var relativt jevnt fordelt på alder:

Chaumoitre 2016. Resultatene fra denne, som er en forholdsvis stor studie på en multietnisk populasjon i Marseille (Frankrike), fant at differensen mellom GP skjeletalderen på bildene i atlaset (modningsstadiene) aldri var større enn 0,5 år fra gjennomsnittet i kronologisk alder. Chaumoitre 2016 anga et pålitelig estimat for variasjonen i alder for denne populasjonen dersom man skal bruke atlaset til estimering av kronologisk alder. Basert på denne studien varierte bredden på 95 % prediksjonsintervaller for alrene 10-19 år for gutter fra 4,0 år til 5,9 år.

Diskusjon

Vi inkluderte studier med to ulike måter å fremstille resultatene på (fremstilling A og B). Flest studier presenterte funnene som gjennomsnittlig forskjell mellom kronologisk

alder og GP skjeletalder for separate aldersgrupper basert på årskull (fremstilling A). Denne analysemetoden gir samlede resultater for GP skjeletalderen til en gruppe individer av ett årskull (for eksempel alle gutter som er 14 år). Fremstillingen reflekterer at Greulich og Pyle-atlaset opprinnelig er utviklet for å beskrive barns normale skjelettmodning. Flere av de inkluderte studiene hadde som mål å utvikle nasjonale referansestandarder. Vi har oppsummert disse studiene, og funnet at samsvaret mellom GP skjeletalder og kronologisk alder i snitt er relativt godt, selv om visse aldersgrupper i enkeltstudier kan ha en gjennomsnittsforskjell på over ett år. Disse studiene viser at Greulich og Pyle-atlaset, som er utviklet basert på røntgenbilder av barn tatt på 1930-tallet, fremdeles beskriver normal skjelettutvikling relativt godt i ulike populasjoner som er undersøkt de siste 10-15 årene. Imidlertid kan ikke standardavvikene som viser hvordan GP skjeletalder fordeler seg fra kjent alder legges til grunn for prediksjonsintervall hvis Greulich og Pyle-atlaset skal brukes til å estimere kronologisk alder (det vil si et «omvendt» scenario).

Analyser av hvordan kronologisk alder fordeler seg fra GP skjeletalder (fremstilling B) er mest egnet for å anslå usikkerheten når Greulich og Pyle-atlaset benyttes til kronologisk aldersestimering. Disse studiene må imidlertid ha en studiepopulasjon som er jevnt fordelt på alder, og sørge for at nedre og øvre aldersgrense ikke settes henholdsvis for høyt eller for lavt. For at resultatene skal bli så riktige som mulig må den inkluderte populasjonen være tilstrekkelig stor, ha tilstrekkelig mange og omtrent like mange individer i hvert alderstrinn samt dekke hele den forventede aldersspredningen for de stadiene man ønsker å undersøke. Hvis ikke kan studien bli preget av fenomenet aldersmimikering, og resultatene blir upålidelige. Vi vurderte at kun én studie unngikk dette og hadde tilstrekkelig godt studiedesign til å gi en mer korrekt skildring av hvordan kronologisk alder fordeler seg fra skjeletalderen: Chaumoitre 2016.

Først når det finnes flere tilsvarende studier er det mulig å vurdere metodens pålitelighet på tvers av studier, regioner og etnisiteter. En alternativ løsning er et prosjekt der man samler inn grunndata der observasjonene av kronologisk alder og modningsstadium er angitt per individ. Tilgangen på slike rådata vil også gi mulighet til å statistisk modellere dataene slik at effekten av aldersmimikering minimeres også i datasett med en ujevn aldersfordeling.

Konklusjon

Vi har oppsummert studiene som fremstiller GP skjeletalder fra kjent kronologisk alder, og funnet at samsvaret mellom GP skjeletalder og kronologisk alder i snitt er relativt godt, selv om visse aldersgrupper i enkeltstudier kan ha en gjennomsnittsforskjell på over ett år. Disse studiene kan ikke legges til grunn når Greulich og Pyle skal brukes til å estimere kronologisk alder.

Studiene som viser hvordan kronologisk alder fordeler seg fra GP skjelettaldrene (modningsstadiene) illustrerer direkte med hvilken usikkerhet Greulich og Pyle-atlaset estimerer kronologisk alder for en populasjon. Vi fant kun én studie med fremstilling B som hadde et pålitelig studiedesign: Chaumoitre 2016. Denne studien viste at for den inkluderte populasjonen vil 95 % prediksjonsintervall være på det minste 4 år bredt og på det meste 5,9 år bredt for gutter mellom 10 og 19 år. Dette gir et inntrykk av hvor stor variasjonen kan være når metoden tas i bruk på en gitt populasjon.

For å utforske usikkerheten for Greulich og Pyle-atlaset til å estimere kronologisk alder for andre populasjoner trengs det flere studier med tilsvarende studiedesign som Chaumoitre 2016. En alternativ løsning dersom grunndata hadde vært tilgjengelig, er å anvende andre statistiske metoder for å unngå aldersmimikering av resultatene.

Key messages (English)

The Greulich and Pyle atlas is used to estimate the age of children and adolescents. This systematic review summarizes the scientific evidence on the agreement between skeletal age from the Greulich and Pyle atlas (hereafter called GP skeletal age) and chronological age.

A total of 19 studies met our criteria. Fifteen studies presented results for GP skeletal age from chronological age and four presented results for chronological age from GP skeletal age. These two approaches are not comparable and we have therefore conducted two separate analyzes.

The difference between GP skeletal age and chronological age was on average rarely more than one year for each age group.

The studies that presented the distribution of chronological age from GP skeletal age are the most relevant if the Greulich and Pyle atlas is used for age estimation. Here we did no meta-analysis, since results from three of the four studies were so affected by the phenomenon «age mimicry» that we do not have confidence in the results. We considered that only one study was conducted in such a way that it can adequately describe the method's ability to estimate age.

Title:

Agreement between chronological age and bone age based on the Greulich & Pyle-atlas for age estimation: a systematic review.

Type of publication:

Systematic review

A review of a clearly formulated question that uses systematic and explicit methods to identify, select, and critically appraise relevant research, and to collect and analyse data from the studies that are included in the review. Statistical methods (meta-analysis) may or may not be used to analyse and summarise the results of the included studies.

Doesn't answer everything:

- Excludes studies that fall outside of the inclusion criteria
- No health economic evaluation
- No recommendations

Publisher:

Norwegian Institute of Public Health

Updated:

Last search for studies:
January 2017.

Peer review:

- Bjørn Anton Graff, Research leader, Diagnostics, Vestre Viken HF
- Lil-Sofie Ording Müller, Consultant Paediatric Radiologist, Oslo University Hospital

Executive summary (English)

Background

Every year, young unaccompanied asylum seekers arrive in Norway, many without knowing how old they are or unable to document their age. To ensure that children receive their rights and that adults are not treated as children, it is necessary to assign a chronological age for these individuals. Evaluation of skeletal maturation of the hand and tooth development has been used for age assessment of asylum seekers in Norway in cases of doubt. These methods have been criticized for their lack of precision, but currently no better methods have been suggested.

In 2016, the Norwegian Institute of Public Health was assigned a national professional responsibility to evaluate and improve the methods used for medical age estimation (from 1.1. 2017 transferred with the Department of Forensic Sciences to Oslo University Hospital). Department of Forensic Sciences has, in cooperation with the Knowledge Centre for the Health Services, conducted a systematic examination of the scientific evidence about several of the medical age estimation methods.

The objective of this systematic review is to assess the scientific evidence on the agreement between chronological age and bone age based on the maturation stages of the Greulich and Pyle atlas (GP skeletal age), and, if possible, to describe any variations between different populations.

Since the Greulich and Pyle atlas is the most widely used method for age estimating based on x-ray of the hand, we chose to focus on this system. In parallel, we have also conducted a systematic review of age estimation using Demirjians development stages on wisdom teeth.

Method

We searched for studies in the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, Embase and Google Scholar. Our search date was May 2016. The literature search covered studies that use x-ray of the teeth or hand, and CT or MRI of the clavicle, knee and ankle for age estimation of children and adolescents between 10 and 25 years. The search for studies that used the Greulich and Pyle atlas was updated in January 2017. Two people independently read title and summary for 10640 references. Of these, we found 658 potentially relevant publications for age estimation using x-ray of the hand. Two people independently considered these studies in full text. We included 19 studies with results that present age estimation based on the Greulich & Pyle atlas. We included two different approaches to present the data (in this report referred

to as approach A and B). Both approaches require a known chronological age and an observed skeletal age:

A) is based on the chronological age and presents the average difference between chronological age and GP skeletal age based on the maturation stages of the Greulich and Pyle atlas for separate age groups (ages not combined in more than annual increments).

B) is based on the GP skeletal age and presents the average chronological age from GP skeletal age in the Greulich and Pyle atlas.

Two people independently assessed risk of bias in the included studies by the QUADAS-2 checklist to assess the quality of diagnostic accuracy studies.

Results

We found 19 studies that compared chronological age with skeletal age based on hand x-ray and the GP atlas. The studies included from 68 to 2614 persons with known chronological age. All but one study included both boys and girls. Five studies were from Turkey, four studies from India, and one study each from Canada, China, France, Iran, Italy, the Netherlands, Pakistan, Spain, Taiwan, and the United States respectively. We considered that the majority of studies had either low or unclear risk of bias based on the QUADAS-2 assessments. The exception was results from the studies with approach B, where three of four studies had a high risk of a particular form of selection bias called "age mimicry".

The majority of studies present results as the average difference between chronological age and GP skeletal age within age groups (approach A). The analysis shows that the variation between studies was greater than expected by chance, even though the variations between studies from different parts of the world were moderate. The difference between GP skeletal age and chronological age rarely exceeded one year for the average of a group in single studies.

Four studies, including two that we analyzed based on the authors' original data, present results as average chronological age within maturation stages from the Greulich and Pyle atlas (approach B). We found that results from three of these studies were influenced by the age composition of the included study population. This bias has previously been described as "age mimicry", which means that the average age observed in each development stage is affected by the included age range and number of participants in each age group. Only one of the four studies with results using approach B had a large study population with a relatively even distribution in age groups: Chaumoitre 2016. This study of a multi-ethnic population in Marseille (France), found that the difference between GP skeletal age from the atlas (maturation stages) was on average never greater than 0.5 years from chronological age. Chaumoitre 2016 gives an appropriate estimate of the variation in age for this population if the atlas is to be used for estimation of chronological age. The width of 95% prediction intervals for boys aged 10 to 19 years ranged from 4.0 years to 5.9 years.

Discussion

We included studies with two different analytical approaches (approach A and B). Most studies presented the average difference between chronological age and GP skeletal age for separate age groups based on age cohorts (approach A). This approach provides overall results for GP skeletal age for a group of individuals within an age cohort (for example, all boys who are 14 years), and reflects that the original use of the Greulich and Pyle atlas was to describe normal skeletal maturation. We have summarized these studies, and found that on average there is a relatively strong association between GP skeletal age and chronological age on a group level, although certain ages in individual studies can have an average difference of more than one year. These studies show that the GP atlas, that is based on radiographs taken of children in the 1930s, still describes normal skeletal development relatively well in different populations studied the last 10-15 years. However, standard deviations from studies presented as approach A cannot be used to estimate predict intervals if the GP atlas is used for estimation of chronological age (which is a "reverse" scenario).

Studies on how chronological age is distributed from GP skeletal age (approach B) are most appropriate to illustrate the uncertainty of the GP atlas when it is used for age estimation. However, these studies must have a study population with an even age distribution, and ensure that the lower and upper age limits are appropriate. For the results to be as correct as possible, the study population must be sufficiently large, have roughly the same number of individuals in each age group and cover the entire expected age range for the stages one aims to examine. Otherwise, the study will be affected by the phenomenon age mimicry and the results will be unreliable. We considered that only one study had a sufficiently good study design to describe appropriately of how chronological age is distributed from skeletal age: Chaumoitre 2016.

More studies similar to this example are needed to evaluate the method's ability to predict age in different populations. An alternative solution is to assemble primary data sets on chronological age and development stage for individuals. Such data will make it possible to use statistical models in order to minimize the effect of age mimicry even in datasets with an uneven age distribution.

Conclusion

We have summarized studies presenting GP skeletal age from known chronological age, and found that there is good agreement between the average values of GP skeletal age and chronological age, although certain age groups in individual studies may have an average difference of about one year. These studies cannot be used to estimate prediction intervals if the GP atlas is used for age estimation.

The studies showing how chronological age is distributed from GP skeletal ages (maturation stages) are most appropriate to illustrate the uncertainty of the GP atlas when it is used for age estimation. We found only one study with approach B that had a reliable study design: Chaumoitre 2016. This study showed that for the included population, 95% prediction intervals vary from 4 years to 5.9 years for boys between 10 and 19 years. This illustrates the uncertainty of the method when it is used on a given population to estimate age.

To explore the uncertainty of the Greulich and Pyle atlas for age estimation in other populations, more studies with the design of Chaumoitre 2016 is needed. An alternative solution is to assemble primary data sets of chronological age and development stage for individuals, and apply statistical models to minimize the effect of age mimicry.

Forord

Norske myndigheter overførte i 2016 det nasjonale ansvaret for aldersestimering av enslige unge asylsøkere til Område for rettsmedisinske fag ved Folkehelseinstituttet (nå Avdeling for rettsmedisinske fag ved Oslo universitetssykehus). I samarbeid med Kunnskapssenteret for helsetjenesten i Folkehelseinstituttet har vi kartlagt det vitenskapelige grunnlaget for flere metoder som benyttes til medisinske aldersvurderinger.

I denne systematiske oversikten oppsummerer vi vitenskapelig dokumentasjon om samsvaret mellom kronologisk alder og skjelettalder som definert i Greulich og Pyle-atlaset for aldersestimering av tenåringer. Parallelt har vi utarbeidet en systematisk oversikt om samsvaret mellom kronologisk alder og Demirjians utviklingsstadier for visdomstänner (1). Vi har utført et litteratursøk som også inkluderte aldersvurdering ved bruk av CT eller MR av krageben, kne eller ankel. Vi har valgt å skrive disse to systematiske oversiktene som separate dokument, men bruker den samme teksten på tvers av dokumentene der det er relevant. Dette gjelder spesielt i introduksjonen, i beskrivelse av metodene og i deler av diskusjonen.

Prosjektgruppen har bestått av personer fra Folkehelseinstituttet (FHI) og Oslo Universitetssykehus (OUS):

- Pål Skage Dahlberg, forsker, prosjektlededer, OUS
- Annhild Mosdøl, seniorforsker, FHI
- Veslemøy Rolseth, forsker, OUS
- Kristoffer Yunpeng Ding, forsker/statistiker, FHI
- Øyvind Bleka, forsker/statistiker, OUS
- Gyri Hval Straumann, forskningsbibliotekar, FHI
- Marianne Skjerven-Martinsen, forsker/lege, OUS
- Gerd Jorunn Møller Delaveris, overlege/sekjonsleder, OUS
- Gunn Elisabeth Vist, seniorforsker/sekjonleder, FHI

Vi takker Signe Flottorp, Brynjar Fure og Torbjørn Wisløff for intern fagfellevurdering og Lil-Sofie Ording Müller og Bjørn Anton Graff for ekstern fagfellevurdering av rapporten. Takk til Marit Johansen for fagfellevurdering av litteratursøket og Vigdis Underland, Jose F. Meneses-Echouez og Signe Flottorp for å ha lest artikler på ulike språk. Vi takker også van Rijn og medarbeidere og Zafar og medarbeidere som delte datasettene fra sine studier for reanalyser.

Signe Flottorp
Avdelingsdirektør

Gunn E Vist
Sekjonsleder

Pål Skage Dahlberg
prosjektleder

Innledning

Det finnes mange situasjoner der det kan råde tvil rundt en ung persons reelle kronologiske alder, blant annet i tilknytning til barnearbeid, prostitution, idrett, seksuell lavalder, kriminell lavalder og asylsaker. I Norge er det primært usikkerhet om alder i saker der utlendingsmyndighetene er i tvil om asylsøkeren er mindreårig eller ikke. En enslig mindreårig asylsøker er en asylsøker under 18 år som ikke har følge av foreldre eller andre med foreldreansvar. Ofte knytter det seg usikkerhet til unge asylsøkeres alder. Dette er fordi mange land ikke har rutiner eller systemer for registrering av fødsler og fordi mange unge asylsøkere ikke kjenner til egen alder eller ikke kan dokumentere dette. Alle personer som skal bli en del av det norske samfunnet må få en kronologisk alder, og denne er avgjørende for personens krav på blant annet beskyttelse, helsehjelp, omsorg og utdannelse.

De europeiske landene har ulik praksis for medisinske aldersvurderinger og aldersfastsettelse (2). I Norge er det Utlendingsdirektoratet som fastsetter alder. Dersom det mangler dokumenter som kan bekrefte søkernes alder eller identitet, er det aktuelt å tilby søkerne en medisinsk aldersundersøkelse. Når medisinske aldersvurderinger er gjennomført inngår det som et underlagsmateriale for aldersfastsettelsen.

Fra 1. januar 2016 fikk Folkehelseinstituttet (nå overført med Retsmedisinske fag til Oslo Universitetssykehus) i oppdrag fra Helse- og omsorgsdepartementet å ha et overordnet medisinskfaglig ansvar for de medisinske aldersvurderingene. Bakgrunnen for dette var at Utlendingsdirektoratet selv ikke har faglig kompetanse til å vurdere kvaliteten i de benyttede metodene eller til å drive forsknings- og utviklingsarbeid. Det var derfor et ønske om at ansvaret skulle legges hos et offentlig medisinskfaglig miljø med bred kompetanse innen sakkyndigvirksomhet og forskning.

Som et ledd i arbeidet for å bygge opp kompetanse innen medisinske aldersvurderinger, ble det innleddet et samarbeid mellom det rettsmedisinske miljøet og Kunnskapssenteret for helsetjenesten i Folkehelseinstituttet. Det utføres i den forbindelse systematiske forskningsoppsummeringer for flere av de mest sentrale metodene som ligger til grunn for dagens praksis for medisinske aldersvurderinger i Norge og Europa.

Det skjer en rekke biologiske endringer i takt med at et menneske vokser og utvikles. Vurderingen av spesifikke utviklingsstadier danner grunnlag for medisinske aldersvurderinger. De mest brukte metodene internasjonalt er basert på vurdering av skelett- (som oftest hånd) og tannutvikling.

Når det er tatt et røntgenbilde av hånd og håndrot, kan man vurdere bildet opp mot et graderingssystem som igjen vil gi et estimat av kronologisk alder. Det fins ingen konensus om hvilket graderingssystem som er best egnet til dette formålet, men systemet

det finnes flest vitenskapelige publikasjoner på er det såkalte Greulich og Pyle-atlaset (3).

Formålet med denne systematiske oversikten er å sammenfatte den forskningsbaserte dokumentasjonen om samsvaret mellom kronologisk alder og skjeletalder basert på modningsstadier fra Greulich og Pyle-atlaset, og eventuelt å belyse variasjoner mellom populasjoner. Denne systematiske oversikten vil sammen med de andre kunnskapsoppsummeringene inngå i et grunnlag for en videre drøfting og anbefaling av hvordan medisinske aldersvurderinger bør gjennomføres i Norge.

Beskrivelse av Greulich og Pyle-atlaset

Skjeletalder (*engelsk: bone age*) vurderes ofte hos barn og unge for å kontrollere vekst, vurdere vekstpotensial, estimere slutthøyde og følge skjelettmodning hos et enkelt individ fordi dette påvirkes både av biologisk variasjon og av en rekke endokrine tilstander og pediatriske sykdommer. En slik kontroll gir informasjon om pasientens relative skjelettmodning ved et gitt tidspunkt, og kan sammen med andre kliniske observasjoner gi grunnlag for å vurdere om pasienten har en utvikling som faller utenom normalområdet. Hos normale barn og unge bør skjeletalderen falle omtrent innen to standardavvik fra referansenormene. En kilde som beskriver referansenormene for skjelettmodning er Greulich og Pyle-atlaset (3).

Denne bruken, det vil si vurdering av et individ opp mot normal skjeletalder, var bakgrunnen for at Greulich og Pyle-atlaset ble utviklet. Valideringsstudien som er gjengitt i Greulich og Pyle-atlaset er derfor bygd opp slik at den inkluderte populasjonen er gruppet etter kronologisk alder, og at skjeletalderen (med gjennomsnitt og standardavvik) er beskrevet gruppevis for hver av disse kronologiske aldrene.

«The Radiographic Atlas of Skeletal Development of the Hand and Wrist» (3), også kjent som Greulich og Pyle-atlaset, ble første gang publisert i 1950 av Dr. William Walter Greulich og Dr. Sarah Idell Pyle. Det er fremdeles en av de mest brukte referansenormene for å bedømme skjeletalder. Atlaset består av referansebilder for jenter og gutter separat, og strekker seg fra fødsel og opp til 18 år for jenter og 19 år for gutter. Ved siden av hvert bilde står det en alder (i denne rapporten kalt GP skjeletalder). Fordelingen av GP skjeletalder i atlaset er ikke identisk for gutter og jenter. Figur 1 viser modningsstadier for henholdsvis gutter og jenter fra ti år og oppover.

Gutter	Jenter
10 år	10 år
11 år	11 år
11 år og 6 måneder	12 år
12 år og 6 måneder	
13 år	13 år
13 år og 6 måneder	13 år og 6 måneder
14 år	14 år
15 år	15 år
15 år og 6 måneder	
16 år	16 år
17 år	17 år
18 år	18 år
19 år	

Figur 1: En oversikt over modningsstadier (bilder) i Greulich og Pyle-atlaset for henholdsvis gutter og jenter fra 10 år og oppover.

Ved siden av hvert bilde i Greulich og Pyle-atlaset står en beskrivelse av viktige endringer man kan observere for dette stadiet. Metoden er komparativ, slik at den som vurderer skjelettalderen til en person, finner bildet og beskrivelsen som ligner mest på røntgenbildet som er tatt. Stadiene er det man kaller diskrete, som betyr at analysen kun kan falle ned på akkurat de modningsstadiene som er gitt i atlaset. Det står i motsetning til kronologisk alder, som er en kontinuerlig skala.

Atlaset ble bygd opp på basis av ett datamateriale av hvite barn fra høyere sosioøkonomiske kår i Ohio på 1930-tallet. Bildene i atlaset er plukket ut ved at forfatterne skjønnsmessig fant det bildet de syntes best representerte et barn med alderen som står oppgitt ved bildet. Individet kunne ha et reelt aldersavvik på maksimalt +/- 2 % fra alderen som er angitt ved bildet. Forfatterne plukket ut 100 bilder av barn med samme alder og kjønn og sorterte bildene etter deres «relative skjelettstatus», fra den minst modne til den mest modne. Så plukket de ut et bilde som de syntes best representerte den «sentrale tendensen». Dette bildet lå ofte, men ikke alltid, omtrent i midten av de 100 sorterte bildene. Slik satte de sammen et atlas av omtrent 30 bilder av henholdsvis jenter og gutter, ettersom jenters håndskjelett modnes raskere enn gutter.

Bruken av Greulich og Pyle-atlaset

Den opprinnelige bruken av atlaset var altså som verktøy for å bedømme hvorvidt et barn hadde en skjelettutvikling innen normalområdet og slik brukes det fremdeles i klinisk praksis. Disse dataene er ikke direkte overførbare til estimering av kronologisk alder, som er en situasjon der man ønsker å vise gjennomsnitt og variasjon for kronolo-

gisk alder fordelt fra et modningsstadium (det vil si fra bildene i atlaset som angir skjelettalder). Det er en vesensforskjell mellom disse to måtene å strukturere dataene på, selv om de reflekterer det samme fenomenet: korrelasjonen mellom modning av skjelettet og økt kronologisk alder.

Det finnes en valideringsstudie i selve Greulich og Pyle-atlaset der det er angitt gjennomsnitt (mean) og standardavvik (SD) for ulike aldersgrupper av henholdsvis jenter og gutter. Resultatene av denne studien er oppgitt i atlases tabell V og VI, og formålet med disse tabellene er å kunne estimere hvorvidt et barn med kjent alder har en skjelettutvikling innen normalområdet. De inkluderte barna i tabellen er derfor sortert etter kronologisk alder før de statistiske dataene er oppgitt. Disse resultatene er derfor ikke overførbare til en situasjon der man ønsker å gjøre «det motsatte»: å estimere kronologisk alder basert på skjelettutviklingen. Statistikken for disse to situasjonene er ulik og man må legge ulike referansetabeller til grunn for beskrivelsen av disse.

I atlaset er det heller ikke oppgitt standardavvik for gutter 18 og 19 år og for jenter 16, 17 og 18 år, som er de avsluttende bildene for henholdsvis gutter og jenter i atlaset.

Særlig det helt avsluttende bildet (18 år for jenter og 19 år for gutter) er problematisk ettersom dette angir en ferdig utviklet hånd og dermed et endestadium. Hvis man skal bruke GP-atlaset til aldersestimering er det to hovedutfordringer knyttet til endestadiet og den statistiske beskrivelsen av dette:

1. Man vet ikke hvor lenge individene som er i endestadiet faktisk har vært der (altså kan en nittenåring med en helt ferdig utviklet hånd ha vært i dette stadiet siden vedkommende var for eksempel 16 år).
2. Når man skal beskrive fordelingen av kronologisk alder for endestadiet vil denne påvirkes av den øvre alderen for de inkluderte individer i studien som ligger til grunn (det vil påvirke gjennomsnitt og standardavvik for endestadiet om man inkluderer individer opp til for eksempel 20 år i stedet for til for eksempel 22 år).

Valideringsstudier av Greulich og Pyle-atlaset

Det finnes en stor mengde studier fra ulike deler av verden som har hatt som målsetting å validere Greulich og Pyle-atlaset. En del av disse studiene har ønsket å beskrive hvordan GP skjeletalder fordeler seg for ulike aldersgrupper i deres populasjon. Dette perspektivet samsvarer med Greulich og Pyle-atlasets opprinnelige intensjon og anvendelse. En annen del av studiene har ønsket å vise hvordan kronologisk alder fordeler seg fra GP skjeletalder. Disse studiene beskriver direkte hva et bilde i atlaset «betyr» i kronologisk alder, og dette perspektivet er mer relevant for atlases bruk til estimering av kronologisk alder. Disse to typene av studier fremstiller resultatene på ulikt vis, her angitt som fremstilling A og B:

- (A) Forskjell mellom kronologisk alder og estimert GP skjeletalder (Greulich og Pyle modningsstadier) for hele aldersgrupper (f. eks. gutter mellom 13,0 og 13,9 år).
- (B) Hvordan den kronologiske alderen fordeler seg ut fra bildene i atlaset (modningsstadiene eller GP skjeletalder)

Fremstilling A

Kronologisk alder	Antall		Skjelettalder
Alle gutter alder 11 år: 11.05 ± 0.24	20		10.80 ± 0.69
Alle gutter alder 12 år: 12.02 ± 0.23	30		11.63 ± 0.84
Alle gutter alder 13 år: 13.07 ± 0.25	42		12.75 ± 1.35
Alle gutter alder 14 år: 14.05 ± 0.24	30		14.18 ± 1.23

Fremstilling B

GP age category	n	mean	s
11 years	58	11.09	1.11
12 years	49	11.98	0.90
13 years	29	12.21	0.90
14 years	46	13.86	0.97

Figur 2: Fremstilling A (venstre figur) beskriver gjennomsnitt og standardavvik for GP skjelettalder for hver gruppe av kronologisk alder. Disse verdiene kan videre brukes til å beskrive differansen mellom kronologisk alder og GP skjelettalder. Dette krever at også korrelasjonskoeffisienten mellom GP skjelettalder og kronologisk alder er oppgitt. Fremstilling B (høyre figur) beskriver gjennomsnitt og standardavvik for kronologisk alder for hver GP skjelettalder (angitt som et bilde i atlaset).

Det finnes også flere andre måter å fremstille samsvaret mellom kronologisk alder og GP skjelettalder. Disse er imidlertid ofte vanskelige å sammenligne og sammenstille, og har derfor ikke blitt inkludert med kriteriene som vi har satt for denne systematiske oppsummeringen.

Metode

Vi har systematisk oppsummert studier som har vurdert samsvaret mellom kronologisk alder og GP skjeletalder for aldrene 14 år og oppover basert på røntgenbilder av hånd sammenlignet med Greulich og Pyle-atlaset (3), samt kartlagt studier av andre aldersestimeringssystemer basert på håndrøntgen. Siden vi arbeidet parallelt med flere systematiske oversikter om aldersestimering, har vi utført ett stort litteratursøk etter metoder basert på røntgen av tenner eller hånd, samt aldersvurdering ved bruk av CT eller MR av krageben, kne eller ankel som sammenholdes med data på kronologisk alder. Arbeidet baserte seg på Kunnskapssenterets metodebok «Slik oppsummerer vi forskning» (4), med følgende spesifikasjoner:

Inklusjonskriterier

Studiedesign: Studier som sammenlignet modningsstadier basert på røntgenbilder av hånd med kjent kronologisk alder

Populasjon: Personer med bekreftet alder mellom 10 og 25 år

Index test: Modningsstadier basert på røntgen av hånd, slik som Greulich og Pyle-atlaset.

Referansestest: Kjent kronologisk alder

Utfall: Gjennomsnittlig forskjell mellom kronologisk alder og GP skjelettalder: CA - SA.

Gjennomsnitt av kronologisk alder for hver GP skjeletalder.

Eksklusjonskriterier:

- Studier som ikke er presentert i fulltekst (konferanseabstrakt og postere)
- Studier som har inkludert døde mennesker (f.eks. levninger)
- Studier som har inkludert færre enn 50 personer mellom 10 og 25 års alder

Litteratursøking

Bibliotekar Gyri Hval Straumann gjennomførte litteratursøket og Marit Johansen fagfellevurderte dette. Det ble utført ett felles søk for studier som utførte aldersestimering basert på røntgen av tenner eller hånd, samt CT eller MR av krageben, kne eller ankel til aldersestimering i det angitte aldersspennet. Appendix 1 inneholder søkerestrategien.

Vi søkte etter primærstudier uten tids- og språkbegrensning i følgende databaser:

- Cochrane Central Register of Controlled Trials (CENTRAL)
- MEDLINE (Ovid) og Pubmed [sb]
- Embase (Ovid)
- Google Scholar

I tillegg søkte vi i registre over pågående studier som beskrevet i Appendix 1.

Artikkelutvelging

Alle identifiserte titler og sammendrag ble lest og vurdert av to prosjektmedarbeidere uavhengige av hverandre. Vi benyttet web-programmet Rayyan til å håndtere referansene i utvelgelsen (5) og fordele arbeidet mellom oss (PSD, KYD, AM, GHS, GEV). Vi markerte hvilke av de potensielt relevante artiklene som omhandlet aldersestimering basert på tenner, hånd eller CT og MR av kragebein, kne eller ankel. De utvalgte referansene for alderestimering basert på håndrøntgen ble deretter vurdert i fulltekst av to personer uavhengige av hverandre (PSD, KYD, AM, GEV, MSM) ut fra inklusjonskriteiene. Prosjektgruppen har lest og vurdert studier på engelsk, skandinaviske språk, kinesisk, japansk og polsk. Vi har fått hjelp av kolleger på Folkehelseinstituttet til å vurdere studier på fransk, tysk, spansk, portugisisk og italiensk. Vi har fått hjelp av andre bekjente til å vurdere studier på russisk og hebraisk.

I tillegg vurderte vi om de relevante artiklene presenterte data på en slik måte at de kunne inngå i analysene våre. Resultatene måtte enten være presentert som gjennomsnitt og standardavvik for kronologisk alder og GP skjeletalder, som differansen mellom disse to, for hvert år av kronologisk alder, eller som gjennomsnitt og standardavvik av kronologisk alder for hver av modningsstadiene fra Greulich og Pyle-atlaset (GP skjeletalder). I den første fremstillingen måtte også korrelasjonskoeffisienten mellom GP skjeletalder og kronologisk alder være oppgitt. Studier basert på andre alderestimeringsmetoder på håndrøntgen enn Greulich og Pyle ble kartlagt.

Kvalitetsvurdering og dataekstraksjon


Vi vurderte den metodologiske kvaliteten av de inkluderte studiene ved hjelp av QUADAS-2 (6). AM vurderte alle de inkluderte studiene og MSM, GEV og PSD ca 1/3 hver, før vurderingene ble sammenlignet og eventuell uenighet løst ved diskusjon. QUADAS-2 er en sjekkliste for å vurdere anvendbarhet og risiko for systematiske skjevheter i valideringsstudier av diagnostiske tester. Vi anser at QUADAS-2 er den sjekklisten for kvalitetsvurdering som passer best for denne systematiske oversikten. Vi tilpasset vurderingen av risikoen for systematiske skjevheter som beskrevet under.

QUADAS-2 sjekklisten har fire domener (6): Utvelgelse av deltakere; Indekstest; Referansetest; og Flyt og tidsaspekt. I metoden er det angitt støttespørsmål som benyttes i vurderingen. Det første domenet vurderer om det er risiko for at deltakere er valgt slik


at resultatene ikke blir representative for populasjonen de kom fra (seleksjonsskjevhets). Det andre domenet vurderer om det kan være risiko for systematisk skjevheter ved gjennomføring eller tolkning av indekstesten - i dette tilfellet spesielt om de som vurderte røntgenbildet hadde kjennskap til barnets kronologiske alder. Siden kjent alder var et inklusjonskriterium vurderte vi at alle studiene hadde lav risiko for systematisk skjevheter for domenet «Referansetest». I domenet «Flyt og tidsaspekt» vurderes det om gjennomføring av studien og analysene er utført på en slik måte at systematiske skjevheter ikke introduseres underveis.

Ujevn fordeling av alder og aldersspennet til studiepopulasjonen kan gi opphav til fenomenet aldersmimikering (7) (illustrert i figur 3).

Aldersmimikering oppstår når individene som er inkludert i en studie ikke har jevn aldersfordeling, enten ved at visse aldersgrupper er overrepresentert eller at nedre og/eller øvre inkluderte alder er slik at det kan påvirke resultatene. Problemet kan oppstå når man forsøker å skildre en variabel (her kronologisk alder) ved å fordele individene på diskrete stadier (her modningsstadium) og deretter beskrive variabelen per stadi.



La oss si man gjør en studie som inkluderer 30 individer, fordelt likt på tre aldersgrupper: 14 år (blå), 15 år (oransje) og 16 år (grønn). Disse kan fordele seg i åtte mulige utviklingsstadier angitt med bokstavene A til H. I den tenkte studien fordeler aldersgruppene seg i de åtte stadiene som til venstre.



Så bestemmer man seg for å doble antallet fjortenåringer i studien, men beholder antallet femten- og sekstenåringer. Utgangspunktet er illustrert på bildet til venstre. Illustrasjonen viser et eksempel på hvordan disse individene kan fordele seg på stadiene.

Hvis vi betrakter stadium D i de to eksemplene er det åpenbart at disse to oppsettene vil gi svært ulik gjennomsnittsalder i dette stadiet (fjortenåringene blir mer dominerende på det nederste bildet). I tillegg vil gjennomsnittsverdien og standardavvik påvirkes av at det ikke finnes individer som er 13 år eller yngre, samt 17 år og eldre. For å unngå effekten av aldersmimikering i denne typen studier må man undersøke tilstrekkelig mange individer som er jevn fordele i kronologisk alder og at aldersspennet minst dekker de modningsstadiene man er interessert i.

Figur 3: Illustrasjon av fenomenet aldersmimikering (7)(engelsk: age mimicry).

Aldersmimikering er en form for seleksjonsskjevhets, forstått som forvrengning av resultatene som følge av fremgangsmåten for innsamling av prøver. I QUADAS-2 sjekklisten faller fenomenet aldersmimikering inn under kategorien spektrumbias – som er et underpunkt under domenet «Utviegelse av deltakere» (6). Vi vurderte derfor studiene opp mot to ekstra spørsmål: 1) Om det var likt antall deltakere i årskull, og 2) om alderspennet var relevant for de analyserte modningsstadiene. Siden aldersmimikering kan ha stor innvirkning på de aktuelle statistiske analysene har vi vurdert dette forholdet separat for å tydeliggjøre den relevante informasjonen.

VR, AM og GEV hentet ut informasjon om populasjonen (alder, etnisitet/region og kjønn), land og årstall for når studien var utført, informasjon om studiedesign og hvilke alderstimeringsmetoder som var benyttet, og informasjonen ble sjekket av en annen person. KYD hentet ut relevante data fra studiene og PSD dobbeltsjekket uttrekket. Vi hentet ut informasjon om:

- Hvor studien var gjennomført, og eventuelt informasjon om etnisitet
- Datainnsamlingsperiode
- Deltakernes kjønn, alder og antall
- Studiedesign
- Metode for å velge ut deltakerne
- Informasjon om alderstimeringsmetoden
- Formålet med studien

Vi hentet ut følgende data (avhengig av hvordan resultatene var oppgitt):

- Gjennomsnitt og standardavvik (SD) av kronologisk alder (CA)
- Gjennomsnitt og SD av GP skjeletalder (SA)
- Gjennomsnittet av differansen mellom CA og SA for gitte kronologiske alderskategorier (hele eller halve år)
- Korrelasjonskoeffisienten (Pearson's) mellom CA og SA.

Analyser

For studier som kun oppga quartilene av kronologisk alder, og ikke standardavvikene (SD), beregnet vi SD ved bruk av normalantagelse.

For den første analysen er vi interessert i å beskrive hvordan differansen mellom kronologisk alder og GP skjeletalder, CA-SA, fordeler seg (basert på fremstilling A). Noen av studiene beregnet SD av denne differansen direkte (basert på differansene på individnivå), mens andre studier oppga SD av kronologisk alder og GP skjeletalder, og korrelasjonen mellom disse to variablene. For de sistnevnte studiene estimerte vi SD for differansen mellom kronologisk alder og GP skjeletalder (for gitte kronologiske alderskategorier) med følgende formel:

$$SD_{diff} = \sqrt{SD_{CA}^2 + SD_{SA}^2 - (2 * Corr * SD_{CA} * SD_{SA})}$$

der SD_{CA} er SD av kronologisk alder, SD_{SA} er SD av GP skjeletalder og $Corr$ er korrelasjonskoeffisienten mellom CA og SA.

De studiene som oppga gruppert gjennomsnitt og SD av kronologisk alder og GP skjelettalder, men ikke korrelasjonskoeffisienten, ble satt til separat liste.

Databehandling av originale datasett

For mange studier manglet det informasjon slik at vi ikke kunne inkludere dem i metaanalysen. For alle disse studiene etterspurte vi den nødvendige informasjonen for å kunne inkludere dem. Vi mottok det originale datasettet fra to av disse studiene. Fra disse ekstraherte vi nødvendig informasjon slik som beskrevet ovenfor, og inkluderte dem i metaanalysen.

Utfallsmål i analysene

Det er to primære utfall i dette prosjektet:

- Gjennomsnittlig forskjell mellom kronologisk alder og GP skjelettalder: CA - SA.
- Gjennomsnitt av kronologisk alder for hver gitte GP skjelettalder.

Estimatene av disse to målene presenteres med 95% konfidensintervall (95 % KI).

Ved å beregne gjennomsnittlig forskjell mellom kronologisk alder og GP skjelettalder får vi målt hvor presist Greulich og Pyle-atlaset angir kronologisk alder på populasjonsnivå/gruppenivå. Gjennomsnitt og varians av kronologisk alder for hver gitte GP skjelettalder vil være knyttet til å kunne anslå hvilken kronologisk alder som bør oppgis (prediksjon) for en person som har observert en gitt skjelettalder (praksis).

Statistisk utførelse

Vi utførte metaanalysen ved bruk av statistikkprogrammet R (versjon 3.3.2) med R-pakken "metafor". Denne utfører inferens om heterogeniteten mellom studiene ved å anta en «random effects» modell.

Vurdering av heterogenitet

For å vurdere statistisk heterogenitet mellom studiene, benyttet vi p-verdien fra heterogenitetesten (basert på «Cochran's Q» testobservator), og andel av den totale variasjonen som kan forklares av heterogenitet ved I^2 . Dersom p-verdien fra testen er lav (f.eks. $<0,1$), indikerer dette at den observerte forskjellen mellom studiene er større enn forskjellen man forventer ved tilfeldig variasjon mellom resultatene i studiene. I^2 statistikken anvendes for å kvantifisere nivået av statistisk heterogenitet. En viktig konsekvens av vurderingen om heterogenitet er at der det er stor heterogenitet mellom studiene, så må det samlede resultatet fra en metaanalyse tolkes med forsiktighet.


Etikk

Vi har valgt å ikke ha en drøfting av etikk i denne rapporten. De etiske aspektene ved aldersestimering basert på Greulich og Pyle-atlasets modningsstadier er spesielt knyttet til hvordan metoden anvendes. Dette drøftes i flere andre publikasjoner (8-10).

Resultater

Resultater av litteratursøket

Søket i elektroniske databaser og registre fram til mai 2016 ga 10059 antatt unike referanser. I tillegg utførte vi et oppdateringssøk kun etter studier som benyttet Greulich og Pyle-atlaset i januar 2017. Dette ga ytterligere 581 treff. Vi vurderte at 9982 referanser enten ikke var relevante eller de ble vurdert i fulltekst for ett av de andre delprosjektene. Vi hentet inn 658 publikasjoner i fulltekst som vi antok kunne være relevante for aldersestimering basert på håndrøntgen. Vi inkluderte 19 studier. Håndtering av de resterende 639 referansene beskrives under.



Figur 4: Flytskjema for søkeresultater og håndtering av referanser

I tillegg kontaktet vi forfattere som hadde relativt store, relevante datasett, men hvor resultatene ikke var presentert slik i publikasjonen at de kunne benyttes i de planlagte analysene. Vi spurte om forfatterne var villige til å gi ytterligere opplysninger eller presentere resultatene slik at de kunne inngå i analysene, alternativt om de var villige til å dele det originale datasettet slik at vi kunne reanalyse disse.

Beskrivelse av de inkluderte studiene

Vi inkluderte 19 studier som analyserte samsvaret mellom kronologisk alder og skjelettalder basert på modningsstadier fra Greulich og Pyle-atlaset. Tabell 1 gir en enkel beskrivelse av de inkluderte studiene. Ytterligere beskrivelser finnes i Appendix 2.

Tabell 1: Beskrivelse av de inkluderte studiene

Forfatter id (referanse)	Land	Etnisitet* eller region	Antall**	Gutter/ jenter	Aldersspenn
Bala 2010 (11)	India	Nord-vest	160	80/80	8-14 år
Buken 2007 (12)	Tyrkia	Nord-vest, Kaukasiere	492	251/292	11-20 år
Cantekin 2012 (13)	Tyrkia	Øst	767	342/425	7-17 år
Chaumoitre 2016 (14)	Frankrike	Etnisk variert	2614	1423/1191	1-20 år
Chiang 2005 (15)	Taiwan	-	370	230/140	0-18 år
Griffith 2016 (16)	Kina	Hong Kong	535	276/259	0-18 år
Gungor 2015 (17)	Tyrkia	Sør	535	259/276	10-18 år
Jimenez-Castellanos 1996 (18)	Spania	Sør	239	139/100	0-14 år
Johnston 1963 (19)	USA	«White»	120	58/62	7-17 år
Koc 2001 (20)	Tyrkia	Sør-Øst	255	255/0	7-17 år
Mohammed 2015 (21)	India	Andhra Pradesh	660	330/330	9-20 år
Nahid 2010 (22)	Iran	Kurdistan	228	102/126	2-22 år
Patel 2015 (23)	India	Gujarat	180	90/90	6-16 år
Patil 2012 (24)	India	Maharashtra	375	194/181	0-19 år
Suri 2013 (25)	Canada	«White»	68	35/33	9-18 år
Tise 2011 (26)	Italia	Kaukasiere	484	349/125	11-19 år
Van Rijn 2001 (27)	Nederland	Kaukasiere	572	278/294	5-20 år
Zafar 2010 (28)	Pakistan	Etnisk variert	889	535/354	0-18 år
Öztürk 2016 (29)	Tyrkia	To grupper	849	375/514	9-19 år

* Basert på eventuell betegnelse benyttet av artikkelforfatterne.

** Totalt antall deltakere og alderspenn inkludert i studien. I analysene av differansen mellom kronologisk alder og estimert skjelettalder er kun årskullene fra 14 år og oppover inkludert.

For ytterligere detaljer om gjennomføring av studiene, se Appendix 2.

De fleste artiklene var publisert etter 2000, med unntak av to studier publisert i 1963 og 1996. Det var fem studier fra Tyrkia, fire studier fra India, og én studie hver fra henholdsvis Canada, Frankrike, Italia, Iran, Kina, Nederland, Spania, Taiwan, Pakistan og USA. Alle studiene inkluderte begge kjønn, bortsett fra én studie som bare undersøkte

gutter (20). I tabell 1 er totalt antall deltakere og aldersspenn i studien presentert. Studiene varierte i størrelse fra kun 68 deltakere (25) til totalt 2614 deltakerne (14). Det smaleste aldersspennet i studien var fra 8-14 år med totalt 7 årskull (11), og det bredeste var 2-22 år med 21 årskull (22). I våre analyser av differansen mellom kronologisk alder og GP skjelettalder (fremstilling A) er kun årskullene fra 14 år og oppover inkludert. For analyser av gjennomsnittlig kronologisk alder fra GP skjelettalder (fremstilling B) er hele studiepopulasjonen benyttet. Vi mottok og reanalyserede datasett fra to studier av van Rijn og medarbeidere (27) og Zafar og medarbeidere (28).

Andre studier vurdert fra fulltekst

Studier basert på Greulich og Pyle-atlaset med mulig relevante datasett

Vi identifiserte ytterligere 107 studier som hadde samlet inn data både av håndskjelettmodning basert på Greulich og Pyle-atlaset og kronologisk alder. Disse studiene ble lagt til side fordi resultatene var presentert på en slik måte at de ikke kunne inkluderes i våre analyser. Noen av disse studiene var analysert med tanke på en annen problemsstilling enn vår. Vi har samlet informasjon om disse studiene fordi de representerer mulige relevante datasett for videre analyser. Vi mener at videre arbeid i størst mulig grad bør basere seg på datasett som allerede finnes snarere enn å utføre flere nye studier. Appendix 3 inneholder en oversikt over disse studiene.

Studier som benytter andre indeks-tester enn Greulich og Pyle

Ytterligere 124 studier hadde validert alderestimeringsmetoder med utgangspunkt i røntgenbilder av hånd som benyttet andre indeks-tester enn Greulich og Pyle-atlaset. Appendix 4 gir en oversikt over disse studiene.

Ekskluderte studier

Av de 658 referansene som ble innhentet i fulltekst, ekskluderte vi 408 etter vurdering mot inklusjons- og eksklusjonskriteriene. Se Appendix 5 for liste over ekskluderte studier med begrunnelse for eksklusjon. Blant disse var det også 19 artikler som ikke ble hentet inn fordi de var vanskelig å skaffe til veie eller svært dyre i innkjøp.

Kvalitetsvurdering av inkluderte studier basert på QUADAS-2

Basert på spørsmålene i QUADAS-2 sjekklisten (6) vurderte vi at ni studier hadde lav risiko og ti studier hadde uklar risiko for at deltakere ikke var representative for populasjonen de kom fra. Seksten studier hadde spesifisert at de som vurderte røntgenbildene var blindet for deltakernes kronologiske alder, mens de øvrige ikke omtalte blinding (uklar risiko for systematisk skjevhetsrisiko). Siden kjent kronologisk alder var et inklusjonskriterium vurderte vi at alle studiene hadde lav risiko for systematisk skjevhetsrisiko for referansestesten. Vi vurderte at én av studiene hadde ekskludert deltakere underveis i analysene på en slik måte at det introduserte høy risiko for systematiske skjevheter.

(15). For ytterligere fire studier vurderte vi at det var uklart om flyt og tidsaspekt i studien kan ha introdusert systematiske skjevheter i resultatene.

Tabell 2: Kvalitetsvurdering av de inkluderte studiene basert på QUADAS-2 sjekklistens fire domener, samt ekstra vurdering av risiko for aldersmimikering.

Forfatter, år	Domener for kvalitetsvurdering basert på QUADAS-2				
	Seleksjonsskjevhet Utvelgelse av deltakere	Alders- Mimikering ¹	Tolkning av indekstesten	Referanse- test	Flyt og tidsaspekt
Bala 2010	?	-	?	+	+
Buken 2007	+	²	-	+	+
Cantekin 2012	?	-	+	+	+
Chaumoitre 2016	+	+	+	+	+
Chiang 2005	?	-	+	+	-
Griffith 2016	?	-	+	+	?
Gungor 2015	+	-	+	+	+
Jimenez-Castellanos 1996	?	-	+	+	?
Johnston 1963	?	-	?	+	?
Koc 2001	?	-	+	+	+
Mohammed 2015	+	-	+	+	+
Nahid 2010	?	-	+	+	?
Patel 2015	+	-	?	+	+
Patil 2012	?	-	+	+	+
Suri 2013	?	-	+	+	+
Tise 2011	+	-	+	+	+
Van Rijn 2001	+	³	+	+	+
Zafar 2010	+	³	+	+	+
Öztürk 2016	+	-	+	+	+

Lav risiko, uklar risiko og høy risiko for systematiske skjevheter i studien.

¹ Vurderingene gjelder for studier med analyser i fremstillingsmåte B.

² Høy risiko for systematisk skjevhet i aldergruppene 18 og 19 år.

³ Forfatterne har gitt oss tilgang til originale data som er re-analyseret for denne systematiske oversikten med fremstillingsmåte B. Symbolet viser at antall individer og fordeling mellom aldersgrupper gir høy risiko for aldersmimikering i disse analysene.

For ytterligere detaljer om gjennomføring av studiene, se beskrivelser i Appendix 2.

I analyser av gjennomsnittlig kronologisk alder i modningsstadier fra Greulich og Pyle-atlaset er det risiko for systematiske skjevheter i form av aldersmimikering (7). Fire

studier inngikk i disse analysene. Vi vurderte at kun én av studiene hadde en inkludert populasjon med et relativt likt antall deltagere fra hvert årskull og at aldersspennet var relevant for de analyserte modningsstadiene. Vi vurderte at risikoen for systematiske skjevheter i form av aldersmimikering i denne ene studien var lav. I de tre andre studiene var det stor variasjon i antall deltagere per årskull. En av disse hadde også et relativt smalt aldersspenn med deltagere fra 11 til 19 år (8 årskull). Vi vurderte at risikoen for systematiske skjevheter i form av aldersmimikering i disse tre studiene var høy.

Differansen mellom kronologisk alder og GP skjelettalder


Syten studier presenterte resultater som differansen mellom kronologisk alder og GP skjelettalder for hvert aldersgruppe separat (fremstillingsmåte A). For tre av disse studiene (16;27;28) var hvert år definert med leveårets start som midtpunkt (f.eks. barn i et alderskull fra 13,5 år til 14,49 år) mens de resterende studiene definerte halvåret som midtpunkt (f.eks. barn i et alderskull 13,0 år til 13,99 år). Siden resultatene omfatter separate analyser av to kjønn og mange årskull har vi valgt å legge alle analyser i Appendix 6 og kun presentere eksempler og oppsummeringer i teksten.

Vi vurderte først om det var hensiktsmessig å sammenfatte resultatene fra disse studiene i metaanalyser. Alle studiene hadde tilsvarende design og gjennomføring, og var godt utført for denne problemstillingen. Studiedeltagerne er gruppert for kronologisk alder før resultatene fremstilles, og aldersfordeling av deltagere i studiene som helhet vil derfor ikke påvirke resultatene. Vi vurderte at studiepopulasjonene kunne betraktes som én populasjon – det vil si barn og unge i en gitt aldersgruppe. Vi vurderte at det ikke var tilstrekkelig informasjon til å gruppere studiene i kategorier etter region eller etnisitet av deltakerne.

Appendix 6 inneholder alle metaanalysene av differansen mellom kronologisk alder og estimert skjelettalder for alle aldersgrupper fra 14 år til 19 år for gutter (Figurene A1-A6) og fra 14 år til 18 år for jenter (Figurene A7-A11). Figurene 5 og 6 viser disse analysene for gutter og jenter med kronologisk alder 17 år. Årskullet 17 år er valgt som eksempel fordi det ligger mellom to viktige grenseverdier i mange juridiske sammenhenger. Figur 5 viser resultatene fra 12 studier for gutter og figur 6 resultatene fra 11 studier for jenter. For hver studie angis antall observasjoner, den gjennomsnittlige differansen mellom kronologisk alder og GP skjelettalder for alle deltakerne, samt usikkerheten i estimatene som 95% KI¹ for gjennomsnittet i hver studiepopulasjon. Dette presenteres både som et punktestimat med en vannrett linje for 95% KI og som tall til

¹ Klammene på hver side av firkantene angir 95% konfidensintervall for populasjonsgjennomsnittet som studien baserer seg på. Gjennomsnittsverdien vil variere hvis man hadde gjort studien flere ganger for samme populasjon. Konfidensintervallet angir usikkerheten på estimatet av populasjonsgjennomsnittet. Det vil si at dersom samme studie hadde vært utført med samme antall på samme populasjon, ville populasjonsgjennomsnittet havnet innen klammene med 95% sannsynlighet (eller 95 av 100 ganger). Jo smalere konfidensintervall jo mer presist vil estimatet for populasjonsgjennomsnittet være. Konfidensintervallet blir generelt bredere dersom estimatet er basert på færre observasjoner.

høyre i figurene. Flere studier bidro med relativt få observasjoner, fra kun 4 deltagere fra én studie til maksimalt 31 deltagere i dette alderskullet.



Figurene 5 og 6: Differansen (i år) mellom kronologisk alder og GP skjeletalder (modningsstadier fra Greulich og Pyle-atlastet) i de inkluderte studiene for henholdsvis gutter og jenter med kronologisk alder 17 år.


CA: kronologisk alder; SA: GP skjeletalder; 95% CI: 95 % konfidensintervall for gjennomsnittet

For alderskullet 17 år varierer gjennomsnittsdifferansen mellom kronologisk alder og GP skjeletalder i de ulike studiene fra en underestimering på 0,6 år til en overestimering på 0,95 år for gutter og en underestimering på 0,70 år til en overestimering på 0,44 år for jenter. Metaanalysen (angitt med «diamanten» og verdiene under hver av figurene) viser en samlet gjennomsnittsforskjell over alle studiene mellom kronologisk alder og GP skjeletalder på 0,06 år (95% KI: -0,25 til 0,38) for gutter og -0,13 år (95% KI: -0,35 til 0,09) for jenter. For begge kjønn er det samlede gjennomsnittlige avviket mellom kronologisk alder og GP skjeletalder derfor mindre enn 2 måneder. Sammen

med relativt snevre konfidensintervall bekrefter disse analysene at det er et godt samsvar mellom kronologisk alder og GP skjelettalder på gruppenivå.

Resultatene fra heterogenitetstestene i figur 5 og 6 viser imidlertid at det er større variasjon/forskjell mellom studiene enn det man vil forvente bare på grunn av tilfeldigheter. Andelen av variansen i metaanalysene som kan tilskrives heterogenitet angis ved I^2 -verdien og er på 91 % for guttene og 87 % for jentene. I diskusjonen drøfter vi hvilke faktorer som kan ha bidratt til denne heterogeniteten. Vi valgte å ikke gå videre med analyser for å forsøke å forklare hvilke faktorer som kan ha bidratt til heterogeniteten (se Diskusjons-kapittelet for begrunnelser).

Hvis man ser på gjennomsnittsdifferansene mellom kronologisk alder og GP skjelettalder i individuelle studier slik de er presentert for alle årskull av gutter fra 14 år til 19 år (Figurene A1-A6) og fra 14 år til 18 år for jenter (Figurene A7-A11) kan man trekke ut følgende funn: vi ser at denne gjennomsnittlige differansen mellom de to variablene i hver studie sjeldent overstiger ett år for gutter (kun i fire av totalt 68 verdier). For gutter i alderen fra 17 år til 19 år er disse differansene aldri over ett år. For jenter er variasjonen noe større (10 av 62 har gjennomsnittsdifferanse på mer enn ett år), men også for de eldste jentene (17 og 18 år) er gjennomsnittsdifferansene fra individuelle studier aldri over ett år. Dette til tross for at det er relativt få observasjoner for hver aldersgruppe i flere av studiene. Figur 7 presenterer det samlede estimatet fra hver av metaanalysene for gutter og jenter i alle årskull (hentet fra Figurene A1-A11).



Figur 7: Samlet gjennomsnittsdifferanse (95 % KI) mellom kronologisk alder og GP skjelettalder fra metaanalysene for alle analyserte årskull og begge kjønn. Figuren angir også p-verdi fra heterogenitettesten og andelen av den totale variasjonen som kan forklares av studieheterogenitet gjennom I^2 -verdien. Hver diamant illustrerer estimert gjennomsnitt med 95% konfidensintervallene.

Figuren viser at gjennomsnittsforskjellen mellom estimert alder og skjeletalder ligger tett på verdien 0, med unntak av gutter 19 år. Totalt var antallet observasjoner for gutter 19 år kun fra 32 personer. Det var gjennomgående større variasjon mellom hver studie i metaanalysene enn det man kan forvente bare på grunn av tilfeldigheter, angitt som p-verdier og I^2 -verdiene som varierer fra 65 % til 94% for alle aldersgruppene. Mulige årsaker til heterogeniteten og vår vurdering om å ikke utføre videre analyser av denne er lik for alle aldersgruppene (se diskusjonskapittelet).


Fra figur 7 kan man se en viss tendensen til at samlet sett (på tvers av alle studiene) har yngre og eldre gutter (14 og 18 år) noe høyere gjennomsnittlig kronologisk alder enn GP skjeletalder. For de andre aldersgruppene ser vi ingen systematisk forskjell mellom GP skjeletalder og den kronologiske alderen. For jentene observerer vi at GP skjeletalder i gjennomsnitt ligger foran kronologisk alder frem til og med 17 år. Fra de er 18 år ser kronologisk alder ut til å ligge foran GP skjeletalder for jenter. Disse trendene er basert på en sammenstilling av flere studier, men kan likevel skyldes tilfeldig variasjon.

Gjennomsnittlig kronologisk alder i utviklingsstadier fra Greulich og Pyle-atlaset

Fire studier (14;26-28), inkludert to studier der vi har reanalysert dataene selv (27;28), presenterte resultater for gjennomsnittlig kronologisk alder i et utviklingsstadium fra Greulich og Pyle-atlaset (fremstilling B). Vi vurderte først om det var hensiktsmessig å sammenfatte resultatene fra disse studiene i metaanalyser. Gjennom QUADAS-2 vurderingene hadde vi vurdert at tre av disse studiene har høy risiko for at resultatene påvirkes av aldersmimikering. Vi vurderer at effekten av aldersmimikering for tre av disse studiene potensielt er så stor at det er høy risiko for at estimatene av gjennomsnittlig kronologisk alder i hvert utviklingsstadium er systematisk forskjøvet. Derfor vurderer vi at det ikke er hensiktsmessig å slå resultatene sammen i metaanalyser, men presenterer funnen grafisk i plottene.

Figurene 8 og 9 viser gjennomsnittlig kronologisk alder i utviklingsstadiet 17 år i Greulich og Pyle-atlaset for henholdsvis gutter og jenter. Appendix 6 inneholder tilsvarende framstilling for alle utviklingsstadiene som er oppgitt fra 10 år til 19 år for gutter (Figurene A12-A23) og 10 til 18 år for jenter (Figurene A7-A11). Siden det er ulikt aldersspenn i studiene er det ulikt antall studier for hvert årskull.

Figur 8 viser resultatene fra 4 studier for gutter og figur 9 resultatene fra de samme 4 studiene for jenter. For hver studie angis antall observasjoner, gjennomsnittlig kronologisk alder blant deltakere som fikk vurdert skjelettutviklingen til stadiet 17 år, samt usikkerheten i estimatene som 95% KI for gjennomsnittet, grafisk og med tall. For eksempel viser figur 9 at laveste gjennomsnittsalder for jenter i utviklingsstadium 17 år er 15,7 år i studien fra Pakistan og høyeste gjennomsnittsalder er 16,8 år i studien fra Italia. Det er større heterogenitet mellom studiene enn det en ville forvente bare på grunn av tilfeldigheter. Det er grunn til å tro at en vesentlig andel av den observerte heterogeniteten for denne fremstillingen av resultater er knyttet til fenomenet aldersmimikering.




Figurer 8 og 9: Gjennomsnittlig kronologisk alder for individer i utviklingsstadiet 17 år fra Greulich og Pyle-atlaset for henholdsvis gutter og for jenter.

CA: kronologisk alder; 95% CI: 95 % konfidensintervall for populasjonsjennomsnittet


Tre av de fire studiene som er presentert her, Tise 2001 (26), van Rijn 2001 (27) og Zafar 2010 (28), har studiepopulasjoner med relativ ujevn fordeling av antall deltakere i de ulike aldergruppene. Det er forventet at dette kan påvirke resultatene i stor grad, noe som også kan observeres.

To eksempler vises her, mens Appendix 6 viser tilsvarende plott for alle studiene for gutter og jenter separat. Eksempel på en studie med ujevn aldersfordeling i studiepopulasjonen er Tise 2011 (26). Funnene fra studien for alle guttene i studien vises grafisk i figur 10.



Figur 10. Grafisk fremstilling av resultater fra Tise 2011 (26) for gutter med GP skjelettalder fra 10 til 18 år. Den øverste delen av figuren viser hver GP skjelettalder for gutter fra 10 til 18 år, med antall, gjennomsnitt (punktet i midten) med ett standardavvik ut til hver side (avslutning på linjene) og 95% konfidensintervall for populasjonsgjennomsnitt (parantes). Den nedre delen av figuren viser den inkluderte populasjonens fordeling på kronologisk alder.

Den eneste inkluderte studien med fremstilling B som har en relativ jevn aldersfordeling er Chaumoitre 2016 (14). I tillegg er antall observasjoner i hver aldersgruppe relativt høy. Funnene fra studien for gutter vises grafisk i figur 11. Vi betrakter dette som et eksempel på en godt designet studie som vil være i stand til å beskrive hvordan den inkluderte populasjonens kronologiske alder fordeler seg fra modningsstadiene (GP skjelettalder) på en mer korrekt måte. Ideelt sett skulle flere studier vært designet på denne måten.



Figur 11. Grafisk fremstilling av resultater fra Chaumoitre 2016 (14) for gutter med GP skjelettalder fra 10 til 19 år. Den øverste delen av figuren viser hver GP skjelettalder for gutter fra 10 til 19 år, med antall, gjennomsnitt (punktet i midten) med ett standardavvik ut til hver side og 95% konfidensintervall for populasjonsjennomsnitt (parentes). Den nedre delen av figuren viser den inkluderte populasjonens fordeling på kronologisk alder.

Tilsvarende plot for alle de fire studiene og begge kjønn ligger i Appendix 6. Det kan vi suelt observeres at det er store forskjeller mellom studiene fremstilt i figur 10 og 11. Aldersmimikering er ikke den eneste forklaringen på forskjellene mellom studiene, men det er en åpenbar kilde til noe av den observerte heterogeniteten, og effekten er mulig å forklare teoretisk i tillegg til at den kan fremvises empirisk.

Diskusjon

Hovedfunn

I denne rapporten har vi systematisk oppsummert primærforskning om samsvaret mellom kronologisk alder og skjelettalder som definert i Greulich og Pyle-atlaset. Vi inkluderte 19 studier som oppfylte kriteriene og som oppga tilstrekkelig data til at vi kunne inkludere dem.

GP skjelettalder øker når kronologisk alder øker. I oppsummeringen fant vi at gjennomsnittsdifferansen mellom disse to variablene sjeldent er større enn et år for hver kronologisk aldersgruppe i hver enkelt studie. For eksempel var denne differansen aldri større enn 0,5 år i noen aldersgruppe i en stor studie av Chaumoitre 2016 (14).

Hvis man ønsker å bruke atlaset til estimering av kronologisk alder er det essensielt å beregne et prediksjonsintervall som sier noe om sikkerhetsmarginen for hvor gammelt et individ kan være (i kronologisk alder) gitt den observerte GP skjelettalderen (fremstillingsmåte B). Chaumoitre 2016 var den eneste av studiene som vi vurderte til å være godt nok utført til å kunne beregne et slikt prediksjonsintervall direkte. Her fant vi at bredden på 95% prediksjonsintervallene varierer fra 4,0 år til 5,9 år for gutter i alderen 10-19 år (se figur 12).

Resultatene fra Chamouitre 2016 er en beskrivelse av den inkluderte populasjonen, som beskrives som multi-etnisk fra Marseille (Frankrike). Det er ikke uproblematisk å anvende resultatene fra Chaumoitre 2016 på enhver populasjon før den aktuelle populasjonen er testet og kartlagt på samme måte.

Kvaliteten på forskningsresultatene

Det har vært rettet mye kritikk mot bruken av Greulich og Pyle-atlaset til estimering av kronologisk alder. Blant annet har ett av ankepunktene vært at atlaset er basert på kaukasiske barn fra høyere sosioøkonomiske kår i Ohio (USA) på 1930-tallet. Selv om flere nye systemer har vært lansert til estimering av alder basert på røntgen av hånd (30-33) forblir Greulich og Pyle-atlaset det mest anvendte. Resultatene i denne oppsummen tyder på at de oppgitte skjelettaldrene fra metoden, i gjennomsnitt, samsvarer relativt godt med kronologisk alder også i populasjoner som er undersøkt de siste 15 årene. Estimering av kronologisk alder på individnivå ut fra et gitt skjelettmødningsstadium er imidlertid usikkert ved anvendelse i praksis.

Studier med fremstilling B er mer egnet til å skildre atlatsets prediksjonsevne for enkeltindivider enn fremstilling A. Av disse fant vi at kun én studie (14) hadde tilstrekkelig godt studiedesign til å unngå fenomenet aldersmimikering.

GP skjelettalder fordelt på grupperte kronologiske aldersintervaller (fremstilling A)

I denne fremstillingen er individene gruppert etter kronologisk alder, og resultatene presenterer GP skjelettalder for hver gruppe. Svært mange studier innen dette kognitivfeltet er organisert på denne måten. Disse studiene har ønsket å beskrive hvordan skjelettalderen fordeler seg for ulike aldersintervaller av deres populasjon. Dette perspektivet samsvarer med Greulich og Pyle-atlatsets opprinnelige intensjon og anvendelse. Denne fremstillingen er mest egnet til å fremstille skjelettutviklingen for en gruppe av en populasjon, la oss si gutter med kronologisk alder 14 år, representert som gjennomsnitt og standardavvik av GP skjelettalder. Fremstilling A påvirkes ikke av aldersmimikering ettersom man ikke fordeler individene i studien på stadier (fenomenet er beskrevet i detalj i figur 3).

Vi kan se fra analysene at kronologisk alder og GP skjelettalder følger hverandre. Det betyr, som forventet, at når kronologisk alder øker, så øker også GP skjelettalder for alle aldersgrupper og begge kjønn i vår analyse. Analysen viser at det i gjennomsnitt (på gruppenivå) er relativt liten forskjell mellom kronologisk alder og GP skjelettalder. Det gir en indikasjon på metodens validitet. Vi observerer imidlertid større variasjon mellom studiene i metaanalysen enn det man ville forventet med statistisk tilfeldighet alene. Forskjeller mellom populasjoner og regioner kan spille inn, men mange andre faktorer kan også ha bidratt til heterogeniteten. Det finnes flere kilder til målefeil i analysene som er utført, blant annet den subjektive tolkningen av bildene opp mot atlatset og røntgenbilder som ikke viser detaljene ved håndskjelettet godt nok (bildene er ofte tatt ved mistenkede skader). Vi valgte å ikke gå videre med analyser for å forsøke å forklare hvilke faktorer som kan ha bidratt til heterogeniteten. På grunn av den store variasjonen mellom studiene, må resultatene basert på sammenslåingen av studiene («diamantene» nederst i hver analyse) tolkes med varsomhet.

Fremstilling A angir ikke direkte hvordan kronologisk alder fordeler seg for gitt GP skjelettalder. Det finnes statistiske metoder som baserer seg på å bygge en modell som beskriver skjelettutviklingen for gitt kronologisk alder (34-36). Disse metodene krever selvsagt at dataene er angitt på individnivå. Et problem med studiene fra litteraturen er at de kun oppgir dataene gruppevis i form av gjennomsnitt og standardavvik, noe som gjør det ekstra krevende å bygge en fornuftig statistisk modell.

Kronologisk alder fordelt fra GP skjelettalder (fremstilling B)

Fremstilling B beskriver hvordan kronologisk alder fordeler seg direkte fra modningsstadiene (bildene i atlatset), angitt som GP skjelettalder. Det er denne fremstillingen som er mest relevant for å beskrive hvor god en metode er til estimere den kronologiske alderen til et individ med ukjent alder. Ulemplen med denne fremstillingen er at den kan være sårbar for fenomenet aldersmimikering. Fenomenet aldersmimikering

drøftes mer inngående i vår parallelle systematiske oversikt om samsvaret mellom kronologisk alder og Demirjians utviklingsstadier for visdomstennere (1).

Vi fant totalt fire studier som ble inkludert i metaanalysen. Tre av studiene (26-28) hadde et studieutvalg som ikke var jevnt fordelt på kronologisk alder (se vedlegg 6). Et slikt ujevnt utvalg vil kunne påvirke resultatene (gjennomsnitt og standardavvik av kronologisk alder) i stor grad. Vi har derfor ikke slått sammen resultatene til et samlet estimat, ettersom vi anser aldersmimikering som et så betydelig problem at det ikke er forsvarlig å gjøre en metaanalyse.

Den eneste studien som har et studieutvalg som er relativt jevnt fordelt på kronologisk alder er Chaumoitre 2016 (14). Dette er et eksempel på en godt designet studie som kan beskrive hvordan den inkluderte populasjonens kronologiske alder fordeler seg på de gitte modningsstadiene (GP skjeletalder) på en mer pålitelig måte. Chaumoitre har brukt resultatene til å beregne 95% prediksjonsintervall, som er gjengitt i figur 12.

SA	Antall	Gj.snitt	St.av.	95% PI-grenser	
				Nedre	Øvre
10	75	10.42	1.18	8.05	12.78
11	58	11.09	1.11	8.86	13.33
11.5	54	11.71	1.16	9.37	14.06
12.5	50	12.28	1.12	10.02	14.55
13	92	13.15	1.00	11.16	15.14
13.5	62	13.66	0.98	11.68	15.64
14	64	14.44	1.05	12.33	16.55
15*	25	15.01	1.01	12.89	17.12
15.5*	14	15.19	0.99	12.98	17.40
16	25	16.28	1.39	13.36	19.21
17	76	16.52	1.13	14.25	18.78
18	53	17.79	1.07	15.62	19.95
19	238	19.35	1.20	16.99	21.72

Figur 12. Tabellen er en gjengivelse av tabell 6 fra Chaumoitre 2016 (14) som viser fordelingen av kronologisk alder fra GP skjeletalder for gutter. Tabellen viser antall individer for hver skjeletalder (Antall), gjennomsnittlig kronologisk alder (Gj. Snitt), Standardavvik (St.av.) og 95% prediksjonsintervall er til høyre i tabellen gjengitt med nedre og øvre grenseverdier. Markeringen "*" betyr at prediksjonsintervallene må brukes med forsiktighet siden de individbaserte dataene ikke passer med normalantagelsen.

Det siste stadiet, som kalles 19 år, er et endestadium som angir en ferdig utviklet hånd. Dette stadiet skiller seg fra alle andre ved at det ikke tar slutt. I en tverrsnittsstudie, der man observerer ett individ én gang, er det derfor umulig å vite hvor lenge individet har vært i et endestadium. Dataene for dette stadiet blir også påvirket av den øvre alderen til den inkluderte populasjonen (øvre «cut-off»). Chaumoitre 2016 har oppgitt at det ikke er noen individer over 21 år i den inkluderte populasjonen (14). Hvis denne alderen hadde vært satt til for eksempel 23 år, ville dette ha påvirket gjennomsnittsverdi og

fordeling av kronologisk alder for endestadiet. I en studie som Chaumoitre 2016 vil endestadiet således skille seg fra de andre stadiene og må tolkes med forsiktighet.

Fra Chaumoitre 2016 ser vi at bredden på 95 % prediksjonsintervaller for aldersintervaller 10-19 år er fra 4,0 til 5,9 år. Dette innebærer at 95 % av de aldersestimerte individene (fra samme populasjon som studien) statistisk sett er estimert til å ligge innenfor de angitte prediksjonsintervallene dersom man hadde brukt denne metoden systematisk på denne populasjonen.

Ideelt sett hadde vi ønsket å finne flere studier med like store og jevnere fordelt aldersgrupper som Chaumoitre 2016 for å kunne belyse Greulich og Pyle-atlasets presisjon og varians til estimering av kronologisk alder for populasjoner i ulike deler av verden. De inkluderte studiene hadde i utgangspunktet lignende design, der det ble utført en blindet vurdering av hvilket skjelettstadium røntgenbildene lignet mest på. Fremstillingen av resultatene er det som begrenser hvorvidt vi kan benytte dataene eller ikke. Det betyr at dersom vi hadde fått tilgang til de individbaserte dataene fra disse studiene, kunne vi bruke statistisk metoder for å fremstille resultatene på en mer egnert måte, og bidra til å belyse bruken av Greulich og Pyle-atlaset for estimering av kronologisk alder.

Regionale forskjeller

Et sentralt spørsmål innen feltet medisinsk aldersestimering er hvor store forskjellene i skjelettmodning er mellom ulike populasjoner fra ulike deler av verden, enten de er forårsaket av genetisk heterogenitet eller ytre årsaker som levekår og levevaner. Fra et medisinsk perspektiv er det spesielt relevant at ulike populasjoner og etniske grupper kan ha genetisk likhet gjennom slektskap, liknende sosioøkonomisk status og levekår, og liknende levevaner (37). Disse faktorene kan hver for seg påvirke vekstmønster, men det er svært krevende å måle den relative betydningen av hver enkelt faktor. I denne systematiske oversikten har vi ikke definert kriterier for inndeling av studier etter geografiske regioner eller etnisiteter. Heterogeniteten vi ser mellom de ulike studiene kan ha en rekke årsaker. Det finnes mange kilder til målefeil i analysene som er utført, blant annet den subjektive tolkningen av bildene opp mot atlaset og røntgenbilder som ikke viser detaljene ved håndskjelettet godt nok (bildene er ofte tatt ved mistenkede skader). De aller fleste av gruppene i analysene våre er små, noe som øker variasjon og kan være en feilkilde. Analysene våre viser imidlertid at en gjennomsnittsforskjell på mer enn ett år mellom GP skjeletalder og kronologisk alder for en gruppe er sjeldent i aldersintervallet vi har sett på, til tross for de kjente kildene til skjevheter.

Styrker og svakheter

Styrken ved denne systematiske oversikten er den systematiske og transparente tilnærmingen vi har benyttet for å svare på spørsmålet. Vi har gjennomført systematiske litteratursøk i mange elektroniske databaser. Vi har benyttet klare inklusjonskriterier og eksklusjonskriterier. To personer har uavhengig av hverandre vurdert hver refe-

ranse etter disse kriteriene, samt uavhengig av hverandre vurdert kvaliteten på de inkluderte studiene. Uavhengige vurderinger er med på å redusere feil og redusere sjansen for feilkilder. Det skal være mulig å etterprøve alle ledd i arbeidet som vi har utført og vurderingene som vi har lagt til grunn for våre konklusjoner.

Selv om vi har utført et grundig litteratursøk med høy grad av sensitivitet, er det alltid mulig at det finnes studier som vi ikke har identifisert. En svakhet ved systematiske oversikter er at de kan bli utdatert når det publiseres flere nye studier, noe som skjer kontinuerlig. Datagrunnlaget i denne systematiske oversikten er antatt å være oppdagert fram til januar 2017.

Overensstemmelse med andre oversikter

En tidligere metodevurdering fra Kunnskapssenteret (nå Kunnskapssenteret i Folkehelseinstituttet), Graff og medarbeidere 2006 (8), omhandlet flere metoder for aldersvurdering av personer mellom 16 og 20 år, deriblant Greulich og Pyle-atlaset. Deres søk ble utført i 2006 og de inkluderte 11 studier som omhandler bruk av Greulich og Pyle-atlaset. Graff og medarbeidere presenterte analysene for begge kjønn, alle aldersgruppene og etniske grupper samlet, mens vi presenterer gutter og jenter separat og for hver kronologiske aldersgruppe. Vi hadde også noe ulike krav til dataformatet på inkluderte studier. Disse forskjellene fører til av det kun er to studier som er overlappende blant de inkluderte i vår rapport og rapporten til Graff og medarbeidere 2006. De andre ni studiene i Graff og medarbeidere er på vår liste over relevante datasett (Appendix 4), og 13 av våre inkluderte studier ble publisert etter deres søkedato.

En systematisk oversikt av Serinelli og medarbeidere (38) har også undersøkt metoder for å vurdere alder etter røntgen av hender. Oversikten til Serinelli og medarbeidere baserte seg på et søk fra desember 2009 og omhandlet både Greulich og Pyle, Tanner-Whitehouse og Fels metodene. For hver aldersvurderingsmetode ble aldersgruppene presentert samlet, mens gutter og jenter ble behandlet separat, og de ble også delt inn i etniske grupper for hvert kjønn. Tre av de fem studiene som er i metaanalysen til Serinelli og medarbeidere er også med i vår oversikt. Tolv av studiene som vi har inkludert er publisert etter søkedatoen i denne oversikten.

Både Graff 2006 (8) og Serinelli 2011 (38) oppsummerte studier med fremstilling A, altså studier som beskriver fordeling av skjeletalder for individer gruppert etter kronologisk alder. Vi har tidligere redegjort for hvorfor denne fremstillingen ikke direkte beskriver usikkerheten ved bruk av Greulich og Pyle-atlaset til estimering av kronologisk alder.

En forskjell mellom vår oversikt og de to tidligere oversiktene er at de begge har delt opp etter etnisitet, mens vi ikke har gjort det. Vi har konkludert med at variasjonen mellom studiene muligens helt kan skyldes tilfeldigheter og metodiske skjevheter. Vi har på grunnlag av dette begrensede datagrunnlaget ment at det ikke er forsvarlig å gå videre til å undersøke andre forklaringsvariabler (subgruppeanalyser), slik som etnisitet eller regionale forskjeller. Vi vurderte også at informasjonen fra studiene ikke var tilstrekkelig til å gruppere studiene i kategorier. For eksempel hadde studien til

Chaumoitre (14) en multi-etnisk populasjon som trolig inneholdt immigranter fra tidligere franske kolonier og Nord-Afrika. Videre vurderte vi at deltakerne i studiene fra Tyrkia (12;13;17;20;29) kunne representere ulike etnisiteter, siden ulike regioner i Tyrkia har ulike populasjonssammensetninger.

Kunnskapshull

Dersom Greulich og Pyle-atlaset skal anvendes til estimering av kronologisk alder i tilfeller der denne er ukjent, er det studier som Chaumoitre 2016 (14) (med fremstilling B) som direkte viser metodens usikkerhet. Denne studien viser hvordan kronologisk alder fordeler seg fra modningsstadiene/skjelettaldrene (bildene i atlaset) direkte. Studien inkluderte mange individer som var relativt jevnt fordelt på kronologisk alder, noe som er en forutsetning for å få representative resultater for å beskrive hvordan kronologisk alder på individnivå fordeler seg på ulike modningsstadier. Flere studier av denne typen kan gi et mer komplett bilde av metodens pålitelighet på tvers av ulike populasjoner og regioner.

En annen mulig løsning er å samle inn grunndata fra mange studier og sammenstille disse. Alle studiene er i utgangspunktet basert på de samme dataoppsettene; registrert kronologisk alder og observert GP skjeletalder for hvert individ. Med en stor og variert mengde av slike datasett kunne man sammenstille og presentere data på en optimal måte for å løse problemet knyttet til fenomenet aldersmimikering.

Konklusjon

I denne systematiske oversikten har vi oppsummert vitenskapelig litteratur om samsvaret mellom skjeletalder gitt i Greulich og Pyle-atlaset og kronologisk alder.

Flertallet av studiene undersøkte hvordan GP skjeletalder fordeler seg for gitte aldersgrupper (kalt fremstilling A i denne rapporten). Dette er den tradisjonelle bruken av atlaset og grunnen til at det ble etablert. Vi har oppsummert disse studiene, og funnet at samsvaret mellom GP skjeletalder og kronologisk alder i snitt er relativt godt, selv om visse aldersgrupper i enkeltstudier kan ha en gjennomsnittsforskjell på over ett år. Disse studiene kan ikke legges til grunn når Greulich og Pyle skal brukes til å estimere kronologisk alder (som blir et «omvendt» scenario).

Studiene som undersøkte hvordan kronologisk alder fordeler seg fra skjelettaldrene (kalt fremstilling B i denne rapporten) illustrerer direkte med hvilken usikkerhet Greulich og Pyle-atlaset estimerer kronologisk alder for en populasjon. Vi fant kun én studie med fremstilling B med en studiedesign og gjennomføring som gir en god beskrivelse av metodens evne til å predikere alder: Chaumoitre 2016. Denne studien viser at for den inkluderte populasjonen vil bredden på 95 % prediksjonsintervall være på det minste 4 år og på det meste 5,9 år for gutter i aldersintervallet 10-19 år. Dette gir et inntrykk av hvor stor variasjonen kan være når metoden tas i bruk på en gitt populasjon.

For å utforske usikkerheten for Greulich og Pyle-atlaset til å estimere kronologisk alder for andre populasjoner, trengs det relevante resultater som er representert på fremstilling B med et tilstrekkelig godt studiedesign. Dersom grunndata hadde vært tilgjengelig, kunne vi anvendt andre statistiske metoder der vi først beskriver hvordan GP skjeletalderen utvikler seg når kronologisk alder øker, og deretter undersøker hvilke kronologiske aldre som mest sannsynlig beskriver en gitt GP skjeletalder.

Referanser

1. Rolseth V, Mosdøl A, Dahlberg PS, Ding KY, Bleka Ø, Skjerven-Martinsen M, et al. Demirjians utviklingsstadier på visdomstrenner for estimering av kronologisk alder: en systematisk oversikt. Rapport 2017. Oslo: Folkehelseinstituttet; 2007.
2. European Asylum Support Office. EASO Age assessment practice in Europe. Luxemburg: Office EAS; 2013.
3. Press SU, red. Greulich W, Pyle S. Radiograph Atlas of Skeletal Development of the Hand and Wrist. 2nd utg. Stanford, California, USA1959.
4. Nasjonalt kunnskapssenter for helsetjenesten. Slik oppsummerer vi forskning. Håndbok for Nasjonalt kunnskapssenter for helsetjenesten. 4. reviderte utg. Oslo: helsetjenesten. Nkf; 2015.
5. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5(1):210.
6. Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155(8):529-36.
7. Bocquet-Appel J-P, Masset C. Farewell to paleodemography. *Journal of Human Evolution* 1982;11(4):321-33.
8. Graff BA, ten Have M, Hofmann B, Norderhaug I. Aldersvurdering av mindreårige asylsøkere. Oslo: Nasjonalt kunnskapssenter for helsetjenesten; 2006.
9. NOAS, Redd Barna. Over eller under 18? : Aldersvurdering av enslige mindreårige asylsøkere. NOAS; 2016.
10. Thevissen PW, Kvaal SI, Willems G. Ethics in age estimation of unaccompanied minors. *J Forensic Odontostomatol* 2012;30 Suppl 1:84-102.
11. Bala M, Pathak A, Jain RL. Assessment of skeletal age using MP3 and hand-wrist radiographs and its correlation with dental and chronological ages in children. *Journal of the Indian Society of Pedodontics and Preventive Dentistry* 2010;28(2):95-9.
12. Buken B, Safak AA, Yazici B, Buken E, Mayda AS. Is the assessment of bone age by the Greulich-Pyle method reliable at forensic age estimation for Turkish children? *Forensic Science International* 2007;173(2):146-53.
13. Cantekin K, Celikoglu M, Miloglu O, Dane A, Erdem A. Bone age assessment: the applicability of the Greulich-Pyle method in eastern Turkish children. *Journal of Forensic Sciences* 2012;57(3):679-82.
14. Chaumoitre K, Saliba-Serre B, Adalian P, Signoli M, Leonetti G, Panuel M. Forensic use of the Greulich and Pyle atlas: prediction intervals and relevance. *Eur Radiol* 2017;27(3):1032-43.
15. Chiang KH, Chou ASB, Yen PS, Ling CM, Lin CC, Lee CC, et al. The reliability of using Greulich-Pyle method to determine children's bone age in Taiwan. *Tzu Chi Medical Journal* 2005;17(6):417-20+53.
16. Griffith JF, Cheng JCY, Wong E. Are western skeletal age standards applicable to the Hong Kong Chinese population? A comparison of the Greulich and Pyle

- method and the tanner and whitehouse method. Hong Kong Medical Journal 2007;13(3 Supplement 3):28-32.
- 17. Gungor OE, Celikoglu M, Kale B, Gungor AY, Sari Z. The reliability of the Greulich and Pyle atlas when applied to a Southern Turkish population. European journal of dentistry 2015;9(2):251-4.
 - 18. Jimenez-Castellanos J, Carmona A, Catalina-Herrera CJ, Vinuales M. Skeletal maturation of wrist and hand ossification centers in normal Spanish boys and girls: a study using the Greulich-Pyle method. Acta Anatomica 1996;155(3):206-11.
 - 19. Johnston FE. Skeletal age and its prediction in Philadelphia children. Hum Biol 1963;35:192-202.
 - 20. Koc A, Karaoglanoglu M, Erdogan M, Kosecik M, Cesur Y. Assessment of bone ages: is the Greulich-Pyle method sufficient for Turkish boys? Pediatrics International 2001;43(6):662-5.
 - 21. Mohammed RB, Rao DS, Goud AS, Sailaja S, Thetay AA, Gopalakrishnan M. Is Greulich and Pyle standards of skeletal maturation applicable for age estimation in South Indian Andhra children? Journal of pharmacy and bioallied sciences 2015;7(3):218-25.
 - 22. Nahid G, Abdorrahim A, Gharib SM, Anvar E. Assessment of bone age in Kurdish children in IRAN. Pakistan Journal of Medical Sciences 2010;26(1):36-9.
 - 23. Patel PS, Chaudhary AR, Dudhia BB, Bhatia PV, Soni NC, Jani YV. Accuracy of two dental and one skeletal age estimation methods in 6-16 year old Gujarati children. Journal of forensic dental sciences : JFDS 2015;7(1):18-27.
 - 24. Patil ST, Parchand MP, Meshram MM, Kamdi NY. Applicability of Greulich and Pyle skeletal age standards to Indian children. Forensic Science International 2012;216(1):200.e1-4.
 - 25. Suri S, Prasad C, Tompson B, Lou W. Longitudinal comparison of skeletal age determined by the Greulich and Pyle method and chronologic age in normally growing children, and clinical interpretations for orthodontics. American Journal of Orthodontics & Dentofacial Orthopedics 2013;143(1):50-60.
 - 26. Tise M, Mazzarini L, Fabrizzi G, Ferrante L, Giorgetti R, Tagliabracchi A. Applicability of Greulich and Pyle method for age assessment in forensic practice on an Italian sample. International Journal of Legal Medicine 2011;125(3):411-6.
 - 27. van Rijn RR, Lequin MH, Robben SG, Hop WC, van Kuijk C. Is the Greulich and Pyle atlas still valid for Dutch Caucasian children today? Pediatric Radiology 2001;31(10):748-52.
 - 28. Zafar AM, Nadeem N, Husen Y, Ahmad MN. An appraisal of Greulich-Pyle Atlas for skeletal age assessment in Pakistan. JPMA - Journal of the Pakistan Medical Association 2010;60(7):552-5.
 - 29. Öztürk F, Karataş O, Mutaf H, Babacan H. Bone age assessment: comparison of children from two different regions with the Greulich-Pyle method In Turkey. Austr J of Forensic Sciences 2016;48(6):694-703.
 - 30. Gilsanz V, Ratib O. Hand bone age. A digital atlas of skeletal maturity. Berlin, Heidelberg: Springer; 2005.
 - 31. Roche A, Chumlea W, Thissen D. Assessing the skeletal maturity of the hand-wrist: FELS method. Springfield, IL: Charles C. Thomas; 1988.
 - 32. Tanner J, Healy M, Goldstein H, Cameron N. Assessment of Skeletal Maturity and Prediction of Adult Height (TW3) Method. London, UK: WB Saunders, Harcourt Publishers Ltd; 2001.
 - 33. Tanner J, Whitehouse R, Cameron N, Marshall W, Healy M, Goldstein H. Assessment of Skeletal Maturity and Prediction of Adult Height (TW2 Method). 2nd utg. London, UK: Academic Press; 1983.

34. Boldsen JL, Milner GR, Konigsberg LW, Wood JW. Transition analysis: a new method for estimating age from skeletons. CAMBRIDGE STUDIES IN BIOLOGICAL AND EVOLUTIONARY ANTHROPOLOGY 2002;73-106.
35. Konigsberg LW, Herrmann NP, Wescott DJ, Kimmerle EH. Estimation and evidence in forensic anthropology: age-at-death. *Journal of Forensic Sciences* 2008;53(3):541-57.
36. Thevissen PW, Fieuws S, Willems G. Human dental age estimation using third molar developmental stages: does a Bayesian approach outperform regression models to discriminate between juveniles and adults? *Int J Legal Med* 2010;124(1):35-42.
37. Bhopal RS. Ethnicity, Race, and Health in Multicultural Societies: Foundations for better epidemiology, public health, and health care. Oxford: Oxford University Press; 2007.
38. Serinelli S, Panetta V, Pasqualetti P, Marchetti D. Accuracy of three age determination X-ray methods on the left hand-wrist: a systematic review and meta-analysis. *Leg Med (Tokyo)* 2011;13(3):120-33.

Vedlegg (Appendixes)

Appendix 1: Literature search

Database: Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to Present

Search date: 2016-05-19

- 1 Age Determination by Teeth/ (1422)
- 2 Age Determination by Skeleton/ (3937)
- 3 (age adj3 (determinat* or estimat* or assess*)).ti. (2851)
- 4 ((forensic or radiological) adj age).ti,ab. (158)
- 5 ((age or maturation or mature or ossification) adj5 (determinat* or estimat* or assess* or examinat*)).ti,ab. (41703)
- 6 (hand\$1 or wrist\$ or carpal or metacarpal or metacarpus or dental or teeth or tooth or third molar* or clavicle* clavicula* or collar bone* or femur or tibia* or fibula* or knee or knees or foot or feet or ankle or ankles).ti,ab. (904235)
- 7 (MRI or MR imag* or magnetic resonance imag* or ct scan* or cat scan* or (comput* adj2 tomograp*) or roentgen or x-ray* or xray* or radiolog* or radiograp*).ti,ab. (1032026)
- 8 5 and 6 and 7 (1297)
- 9 1 or 2 or 3 or 4 or 8 (7491)
- 10 exp Animals/ (20185560)
- 11 Humans/ (15941900)
- 12 10 not (10 and 11) (4243660)
- 13 9 not 12 (7007)
- 14 (greulich adj2 pyle).ti,ab. (238)
- 15 (tanner adj2 whitehouse).ti,ab. (246)
- 16 demirjian.ti,ab. (218)
- 17 haavikko.ti,ab. (20)
- 18 kullman.ti,ab. (6)
- 19 nortje.ti,ab. (5)
- 20 liversidge.ti,ab. (10)
- 21 kvaal.ti,ab. (13)
- 22 or/14-21 (674)
- 23 13 or 22 (7178)

Database: Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to Present

Search date: 2017-01-03

- 1 Age Determination by Skeleton/ (4617)
- 2 (age adj3 (determinat* or estimat* or assess*)).ti. (3296)
- 3 ((forensic or radiological) adj age).ti,ab. (201)
- 4 ((age or maturation or mature or ossification) adj5 (determinat* or estimat* or assess* or examinat*)).ti,ab. (50880)
- 5 (hand\$1 or wrist\$ or carpal or metacarpal or metacarpus).ti,ab. (423891)

- 6 (roentgen or x-ray* or xray* or radiolog* or radiograp*).ti,ab. (738776)
- 7 4 and 5 and 6 (534)
- 8 1 or 2 or 3 or 7 (7429)
- 9 exp Animals/ (23024664)
- 10 Humans/ (18186804)
- 11 9 not (9 and 10) (4837860)
- 12 8 not 11 (7013)
- 13 (greulich adj2 pyle).ti,ab. (286)
- 14 12 or 13 (7059)

Database: Embase 1974 to 2016 May 18

Search date: 2016-05-19

- 1 age determination/ (5176)
- 2 (age adj3 (determinat* or estimat* or assess*)).ti. (3291)
- 3 ((forensic or radiological) adj age).ti,ab. (199)
- 4 ((age or maturation or mature or ossification) adj5 (determinat* or estimat* or assess* or examinat*)).ti,ab. (57474)
- 5 (hand\$1 or wrist\$ or carpal or metacarpal or metacarpus or dental or teeth or tooth or third molar* or clavicle* clavicula* or collar bone* or femur or tibia* or fibula* or knee or knees or foot or feet or ankle or ankles).ti,ab. (1087091)
- 6 (MRI or MR imag* or magnetic resonance imag* or ct scan* or cat scan* or (comput* adj2 tomograp* or roentgen or x-ray* or xray* or radiolog* or radiograp*).ti,ab. (1334461)
- 7 4 and 5 and 6 (1656)
- 8 1 or 2 or 3 or 7 (8121)
- 9 exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or animal cell/ or nonhuman/ (23089391)
- 10 human/ or normal human/ or human cell/ (17222575)
- 11 9 not (9 and 10) (5913580)
- 12 8 not 11 (7315)
- 13 (greulich adj2 pyle).ti,ab. (338)
- 14 (tanner adj2 whitehouse).ti,ab. (279)
- 15 demirjian.ti,ab. (208)
- 16 haavikko.ti,ab. (19)
- 17 kullman.ti,ab. (7)
- 18 nortje.ti,ab. (4)
- 19 liversidge.ti,ab. (18)
- 20 kvaal.ti,ab. (11)
- 21 or/13-20 (794)
- 22 12 or 21 (7692)

Database: Embase 1974 to 2016 December 30

Search date: 2017-01-03

- 1 age determination/ (5542)
- 2 (age adj3 (determinat* or estimat* or assess*)).ti. (3437)
- 3 ((forensic or radiological) adj age).ti,ab. (225)
- 4 ((age or maturation or mature or ossification) adj5 (determinat* or estimat* or assess* or examinat*)).ti,ab. (62094)
- 5 (hand\$1 or wrist\$ or carpal or metacarpal or metacarpus).ti,ab. (502099)
- 6 (roentgen or x-ray* or xray* or radiolog* or radiograp*).ti,ab. (776099)
- 7 4 and 5 and 6 (641)
- 8 1 or 2 or 3 or 7 (7816)
- 9 exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or animal cell/ or nonhuman/ (24312061)
- 10 human/ or normal human/ or human cell/ (18443969)

- 11 9 not (9 and 10) (5914908)
 12 8 not 11 (7052)
 13 (greulich adj2 pyle).ti,ab. (374)
 14 12 or 13 (7225)

Database: Central

Search date: 2016-05-19

#1	MeSH descriptor: [Age Determination by Skeleton] explode all trees	99
#2	MeSH descriptor: [Age Determination by Teeth] explode all trees	5
#3	(age near/3 (determinat* or estimat* or assess*)):ti	30
#4	((forensic or radiological) next age)	0
#5	((age or maturation or mature or ossification) near/5 (determinat* or estimat* or assess* or examinat*))	3474
#6	(hand or hands or wrist or wrists or carpal or metacarpal or metacarpus or dental or teeth or tooth or third molar* or clavicle* clavicula* or collar bone* or femur or tibia* or fibula* or knee or knees or foot or feet or ankle or ankles)	78361
#7	(MRI or (MR next imag*) or (magnetic next resonance next imag*) or ct-scan* or cat-scan* or (comput* near/2 tomograp*) or roentgen or x-ray* or xray* or radiolog* or radiograp*)	52159
#8	#5 and #6 and #7	236
#9	(greulich near/2 pyle)	6
#10	(tanner near/2 whitehouse)	12
#11	demirjian	11
#12	haavikko	1
#13	kullman	17
#14	nortje	9
#15	liversidge	9
#16	kvaal	5
#17	#1 or #2 or #3 or #4 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 in Trials	197

Database: Central

Search date: 2017-01-03

#1	MeSH descriptor: [Age Determination by Skeleton] explode all trees	99
#2	(age near/3 (determinat* or estimat* or assess*)):ti	34
#3	((forensic or radiological) next age)	1
#4	((age or maturation or mature) near/5 (determinat* or estimat* or assess* or examinat*))	3740
#5	(hand or hands or wrist or wrists or dental or teeth or tooth or (third next molar*) or carpal or metacarpal or metacarpus)	54096
#6	(roentgen* or x-ray* or xray* or radiolog* or radiograp*)	39127
#7	#4 and #5 and #6	167
#8	#1 or #2 or #3 or #7	288
#9	(greulich near/2 pyle)	7
#10	#8 or #9 in Trials	138

Database: PubMed

Search date: 2016-03-14

Search (((publisher [sb]) OR pubstatusaheadofprint)) AND (((age determinat*[Title/Abstract]) OR age estimat*[Title/Abstract]) OR age assess*[Title/Abstract])	46
---	----

Database: PubMed

Dato for søk: 2017-01-03

Search (((publisher [sb]) OR pubstatusaheadofprint)) AND (((age determinat*[Title/Abstract]) OR age estimat*[Title/Abstract]) OR age assess*[Title/Abstract])	76
---	----

Database: Google Scholar

Search date: 2016-03-23

"age estimation" OR "estimation of age" OR "estimating age" OR "age determination" OR "determination of age" OR "determining age" OR "age assessment" OR "assessing age" OR "assessment of age"	Leste første 100 treff.
---	-------------------------------

Database: Google Scholar

Dato for søk: 2017-01-03

"age estimation" OR "estimation of age" OR "estimating age" OR "age determination" OR "determination of age" OR "determining age" OR "age assessment" OR "assessing age" OR "assessing age" OR "assessment of age"	Leste første 100 treff.
--	-------------------------------

Database: Clinicaltrials.gov

Dato for søk: 2016-03-15

"age estimation" OR "estimation of age" OR "estimating age" OR "age determination" OR "determination of age" OR "determining age" OR "age assessment" OR "assessing age" OR "assessment of age"	16
greulich OR pyle OR demirjian OR haavikko OR kullman OR nortje OR liversidge OR kvaal	14

Database: Clinicaltrials.gov

Dato for søk: 2017-01-03

"age estimation" OR "estimation of age" OR "estimating age" OR "age determination" OR "determination of age" OR "determining age" OR "age assessment" OR "assessing age" OR "assessment of age"	16
greulich OR pyle	8

Database: WHO - International Clinical Trials Registry Platform (ICTRP)

Dato for søk: 2016-03-15

age estimation OR estimation of age OR estimating age OR age determination OR determination of age OR determining age OR age assessment OR assessing age OR assessment of age	20
greulich OR pyle OR demirjian OR haavikko OR kullman OR nortje OR liversidge OR kvaal	2

Database: WHO - International Clinical Trials Registry Platform (ICTRP)

Dato for søk: 2017-01-03

age estimation OR estimation of age OR estimating age OR age determination OR determination of age OR determining age OR age assessment OR assessing age OR assessment of age	22
greulich OR pyle	2

Appendix 2: Description of included studies with quality assessment

Abbreviations used: CA: Chronological age; SA: Skeletal age based on the index test.

Bala M, Pathak A, Jain RL. Assessment of skeletal age using MP3 and hand-wrist radiographs and its correlation with dental and chronological ages in children. Journal of the Indian Society of Pedodontics and Preventive Dentistry. 2010;28(2):95-9.		
Population: Country, ethnicity, place and year	India (North-West), from outpatient department of Pedodontics and Preventive Dentistry, Government Dental College and Hospital, Patiala, and various schools. Only children of normal weight (between 3rd and 97th percentile curves for a particular age). All radiographs taken at the same day (year not specified).	
Sample, sex and age	160 participants, 80 boys and 80 girls, aged 8-14 years.	
Study design	Cross-sectional.	
Index test	Radiographed for middle phalanx of third finger (MP_3) and hand-wrist of the right hand. Greulich & Pyle atlas.	
Other comments	Dental age was assessed from IOPA radiographs of right permanent maxillary canine based on Nolla's calcification stages	
Aim of the study	"...to assess skeletal age using MP 3 and hand-wrist radiographs and to find the correlation amongst the skeletal, dental and chronological ages"	
QUADAS- 2 assessment	Rating	Comment
Patient selection method:	"...children were selected from..." No further description of sampling method. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	
- Avoid inappropriate exclusions?	Yes	Weight criteria considered appropriate.
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Unclear	Not mentioned
DOMAIN 2: Index test interpretation	Unclear risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	All numbers included in results-tables
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Buken B, Safak AA, Yazici B, Buken E, Mayda AS. Is the assessment of bone age by the Greulich-Pyle method reliable at forensic age estimation for Turkish children? Forensic Science International. 2007;173(2):146-53.		
Population: Country, ethnicity, place and year	Turkey (North West). Schoolchildren in Central Düzce, and students at Abant Izzet Baysal University. Physically and mentally healthy, no history of chronic or severe illness, right handed with no history of trauma to the left hand and wrist. Caucasian with low-middle socioeconomic status. Data collection period not clear.	
Sample, sex and age	492 participants, 251 boys aged 11-17 years, 241 girls aged 11-19 years.	

Study design	Cross-sectional.	
Index test	Radiograph of right hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“...we investigated whether or not the Greulich–Pyle (G–P) method is sufficient at forensic age estimation for Turkish Children.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:		Random selection of schools in the area, random selection of children from these schools. Subjects 18 years and older were volunteers. No discussion of uniform distribution into age cohorts found.
- Consecutive or random sample of patients?	Unclear	
- Avoid inappropriate exclusions?	11-17 years: Yes 18-19 years: No	18-19 years: 20 female and 50 males were excluded because epiphyseal union were completed.
DOMAIN 1: Patient selection	High risk of bias for 18 and 19 years Low risk of bias for 11 to 17 years	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	The persons who interpreted the radiograph only knew the participants' gender.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Cantekin K, Celikoglu M, Miloglu O, Dane A, Erdem A. Bone age assessment: the applicability of the Greulich-Pyle method in eastern Turkish children. Journal of Forensic Sciences. 2012;57(3):679-82.		
Population: Country, ethnicity, place and year	Turkey (East). Children attending the Department of Orthodontics at the University of Ataturk. Physically and mentally healthy with no history of chronic or severe disease, no history of trauma on left hand-wrist. Data collection period not clear.	
Sample, sex and age	767 participants, 342 boys and 425 girls, aged 7-17 years	
Study design	Cross-sectional.	
Index test	Conventional roentgenograms of left hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“... to investigate whether or not the Greulich–Pyle (GP) method is adequate for Turkish children.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Random selection of radiographs at the Department. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	Random selection of radiographs at the Department
- Avoid inappropriate exclusions?	Unclear	
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	

- Index test interpreted without knowledge of CA?	Yes	Dental and skeletal maturation assessments performed without any knowledge of the children's CAs.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Unclear	Assumed
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Chaumoitre K, Saliba-Serre B, Adalian P, Signoli M, Leonetti G, Panuel M. Forensic use of the Greulich and Pyle atlas: prediction intervals and relevance Eur Radiol. 2017 Mar;27(3):1032-1043. doi: 10.1007/s00330-016-4466-4. Epub ahead of print 2016 Jun 29.		
Population: Country, ethnicity, place and year	France (South) with multi-ethnic sample. The authors write that France forbid ethic registration, but that Marseilles is a cosmopolitan melting pot with many immigrants. Patients consulting for minor injuries at University hospital of Marseilles from 2006 to 2011. Included if medical records contained no disease that could affect bone maturation.	
Sample, sex and age	2614 participants, 1423 boys aged 0-21 years and 1191 girls aged 0-20 years.	
Study design	Cross-sectional, from archives.	
Index test	Radiograph of right or left hand and wrist, Greulich & Pyle atlas. Both hands used indifferently.	
Aim of the study	"... to assess its [the Greulich & Pyle atlas] accuracy and to calculate the prediction intervals at 95% for forensic use."	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Individuals were consecutively recruited from the x-rays database in order to have sufficient number of patients by year of chronological age and equally distributed between boys and girls for prediction interval calculation. The article states that "they restrained the sample to boys aged less than 21 and girls less than 20 years, taking into account higher GP categories (19 for boys and 18 for girls) and not distorting the calculation of prediction interval."	
- Consecutive or random sample of patients?	Yes	Consecutive
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Uniform distribution of participants in age cohorts considered by authors. Number of participants in each age cohort varies from 58 to 102 observations among boys (highest numbers in age cohorts 12 and 13 years) and from 57 to 76 observations among girls (highest numbers in age cohort 14 years). Age range 0-21 years (boys)/0-20 years (girls). Our judgement is that the age cohorts are quite uniform and larger than in most studies, although not equal. Age range appropriate for the analyses. Low risk of bias	
- Index test interpreted without knowledge of CA?	Yes	The readers only knew the sex of the child and were blinded to the chronological age
DOMAIN 2: Index test interpretation	Low risk of bias	

- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	
DOMAIN 4: Patient flow and timing bias?	Low risk bias	

Chiang KH, Chou ASB, Yen PS, Ling CM, Lin CC, Lee CC, et al. The reliability of using Greulich-Pyle method to determine children's bone age in Taiwan. <i>Tzu Chi Medical Journal</i> . 2005;17(6):417-20+53.		
Population: Country, ethnicity, place and year	Taiwan (Hualien), children who came to an emergency outpatient department because of suspected trauma. Included radiographs that did not show bone or soft tissue abnormality and with no record of congenital disorder or developmental disturbances. Data collection period not clear.	
Sample, sex and age	370 participants, 230 boys and 140 girls, aged 0-18 years.	
Study design	Cross-sectional.	
Index test	Radiograph of the left hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“... we investigated whether or not the Greulich-Pyle method is sufficient for the determination of the skeletal age of children in Taiwan.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Consecutive children at an emergency outpatient department. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	Consecutive
- Avoid inappropriate exclusions?	Unclear	
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	Analysed by two radiologists who were unaware of the age of the children
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	No	Radiographs were excluded if the inter-observer variation was more than 12 months. 108 cases excluded because of poor film quality (58), inter-observer difference (40), and 10 of other causes.
DOMAIN 4: Patient flow and timing bias?	High risk of bias	

Griffith JF, Cheng JCY, Wong E. Are western skeletal age standards applicable to the Hong Kong Chinese population? A comparison of the Greulich and Pyle method and the Tanner and Whitehouse method. <i>Hong Kong Med J</i> 2007;13(Suppl 3):S28-32.		
Population: Country, ethnicity, place and year	China (Hong Kong). Children attending the accident and emergency departments of Prince of Wales, North District, and Tuen Mun hospitals with suspected fractures from October 2000 to December 2002.	
Sample, sex and age	535 participants, 276 females, 259 males, aged 10-18 years	

Study design	Cross-sectional.	
Index test	Radiographs of the hand and wrist, both the left and right sides were used. Greulich & Pyle atlas.	
Other comments	Radiographs also scored based on the Tanner and Whitehouse RUS and carpal scores.	
Aim of the study	“to (1) assess whether skeletal age standards currently used are applicable to modern day Hong Kong children/adolescents, and (2) to compare the GP and TW3 methods of skeletal age assessment with reference to modern day Hong Kong children/adolescents.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	All consecutive patients. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	Consecutive
- Avoid inappropriate exclusions?	Unclear	
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	“undertaken [...] without prior knowledge of the patient’s age”
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Unclear	
DOMAIN 4: Patient flow and timing bias	Unclear risk of bias	

Gungor OE, Celikoglu M, Kale B, Gungor AY, Sari Z. The reliability of the Greulich and Pyle atlas when applied to a Southern Turkish population. European journal of dentistry. 2015;9(2):251-4.		
Population: Country, ethnicity, place and year	Turkey (South). Radiographs were selected retrospectively from the archives of Akdeniz University. Children born and raised in Southern region of Turkey, no history of chronic or severe physical or mental illness, right-handed with no history of trauma or injury to the left hand wrist. Data collection period not clear.	
Sample, sex and age	535 participants, 259 boys and 276 girls, aged 10-18 years	
Study design	Cross-sectional, from archives.	
Index test	Radiograph of the left hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“to evaluate the applicability of Greulich and Pyle (GP) method for Southern Turkish population.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Retrospectively selected from an archive. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	Consecutive
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	Performed with the chronological age blinded.
DOMAIN 2: Index test interpretation	Low risk of bias	

- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Jiménez-Castellanos J, Carmona A, Catalina-Herrera CJ, Viñuales M. Skeletal maturation of wrist and hand ossification centers in normal Spanish boys and girls: a study using the Greulich-Pyle method. <i>Acta Anatomica</i> . 1996;155(3):206-11.		
Population: Country, ethnicity, place and year	Spain (South). Radiographs from the records of Seville University Hospital. Radiographs chosen to be made between ± 15 days from the child's birthday. Data collection period not clear.	
Sample, sex and age	239 participants, boys and girls, aged 1-14 years	
Study design	Cross-sectional, from archives.	
Index test	Radiograph of the hand and wrist, side not mentioned, Greulich & Pyle atlas.	
Aim of the study	"We aimed to establish both a chronological sequence of postnatal skeletal maturation in Spanish children and, at the same time intersexual differences."	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method not specified. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	Not specified.
- Avoid inappropriate exclusions?	Unclear	Not mentioned
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Unclear	In order to minimise subjectivity, the radiograph series was studied twice by the same researcher with an interval between observations, and also by two different workers, we judge that they was likely blinded to CA.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Unclear	
DOMAIN 4: Patient flow and timing bias?	Unclear of bias	

Johnston FE. Skeletal age and its prediction in Philadelphia children. <i>Human Biology</i> . 1963;35:192-202.		
Population: Country, ethnicity, place and year	USA (Philadelphia). From the files of the Philadelphia Center for Research in Child Growth, the Growth Center school series. Described as "White" schoolchildren. Selected based on "clinical normality" (unspecified). Examined serially throughout their school years. Data collection period not clear.	

Sample, sex and age	120 participants, 58 boys and 62 girls, age range for examinations 7-17 years.	
Study design	Mixed longitudinal sample	
Index test	Radiograph of the hand and wrist, side not mentioned, Greulich & Pyle atlas, standards from 1959.	
Aim of the study	“Information is presented here on the rate of skeletal maturation of Philadelphia white children as evaluated from standards based upon the Brush series. It, therefore, compares rates of development in normal Philadelphia children with their Cleveland age-peers.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method not specified. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	Selection method not specified.
- Avoid inappropriate exclusions?	Unclear	“Due to the inevitable dropouts, and to some later additions, the data are mixed longitudinal, and sample size varies for each age”.
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Unclear	Not described.
DOMAIN 2: Index test interpretation	Unclear risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Unclear	Not described.
DOMAIN 4: Patient flow and timing bias?	Unclear risk of bias	

Koc A, Karaoglanoglu M, Erdogan M, Kosecik M, Cesur Y. Assessment of bone ages: is the Greulich-Pyle method sufficient for Turkish boys? Pediatrics International. 2001;43(6):662-5.		
Population: Country, ethnicity, place and year	Turkey (South-East). Boys with weight and height outside 3 rd and 97 th growth percentile or with chronic diseases were excluded. Data collection period not clear.	
Sample, sex and age	255 participants, all boys, aged 7-17 years.	
Study design	Cross-sectional.	
Index test	Conventional roentgenograms of left hand and wrist, Greulich & Pyle atlas. Rated by two radiologists and one paediatrician separately, the average value used as bone age.	
Aim of the study	“...we investigated whether or not the GP method is sufficient for Turkish children for the determination of the skeletal age.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method not described. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	Selection method not described.
- Avoid inappropriate exclusions?	Unclear	Selection method not described.
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	

- Index test interpreted without knowledge of CA?	Yes	Rated without any knowledge of chronological age.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	Assumed ok.
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Mohammed RB, Rao DS, Goud AS, Sailaja S, Thetay AA, Gopalakrishnan M. Is Greulich and Pyle standards of skeletal maturation applicable for age estimation in South Indian Andhra children? Journal of pharmacy and bioallied sciences. 2015;7(3):218-25.

Population: Country, ethnicity, place and year	India (Andhra Pradesh). Randomly selected patients from the outpatient Department of Oral Medicine, GITAM Dental College, Andhra Pradesh. Parents belonged to middle socioeconomic status groups. Included children that were local residents, with left hand with neither bone (including fracture) or soft tissue abnormalities, and no congenital and growth/ nutritional disturbances or history of chronic illness. Data collection period not clear.	
Sample, sex and age	660 participants, 330 boys and 330 girls, aged 9-20 years	
Study design	Cross-sectional.	
Index test	Digital radiographs of left hand and wrist, Greulich & Pyle atlases.	
Aim of the study	"... to evaluate the applicability of the G-P standards to contemporary South Indian children of 9-20 years age and also the assess the possible relation between estimated SA and CA."	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Randomly selected from patients. No discussion of uniform distribution into age cohorts found in article, but each sex and age cohort has almost the same number of participants.	
	Rating	Comment
- Consecutive or random sample of patients?	Yes	Random.
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	Undertaken without prior knowledge of the CA.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	
DOMAIN 4: Patient flow and timing bias?	Low risk of bias	

Nahid G, Abdorrahim A, Gharib SM, Anvar E. Assessment of bone age in Kurdish children in Iran. Pakistan Journal of Medical Sciences. 2010;26(1):36-9.

Population: Country, ethnicity, place and year	Iran (Kurdistan state). Healthy children and adolescents recruited from schools. Only children of normal height and weight (between 3rd and 97th percentile curves for a particular age) without chronic illness, on medications, vitamin preparations, calcium supplements, or ill for longer than two weeks during previous six months, or hospitalized any time since birth. Data collection in 2007.	
Sample, sex and age	228 participants, 102 boys and 126 girls, aged 7-14 years.	
Study design	Cross-sectional.	
Index test	Radiograph of the left hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“To investigate whether or not the Greulich- Pyle method is sufficient for Kurdish children for determination of the skeletal age.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Recruited from schools. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	
- Avoid inappropriate exclusions?	Unclear risk	Many exclusion criteria.
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Low risk	Description of blinding.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	No	If the inter-observer variation was more than 12 months, the radiographs were excluded (22 case).
DOMAIN 4: Patient flow and timing bias?	Unclear risk of bias	

Patel PS, Chaudhary AR, Dudhia BB, Bhatia PV, Soni NC, Jani YV. Accuracy of two dental and one skeletal age estimation methods in 6-16 year old Gujarati children. Journal of forensic dental sciences : JFDS. 2015;7(1):18-27.		
Population: Country, ethnicity, place and year	India (Gujarat). Patients of the outpatient department of Oral Medicine and Radiology in Gandhinagar district. Subjects with all teeth of mandibular left quadrant at least partially erupted and right hand and wrist intact were included. Exclusion criteria were uncertain date of birth, missing, impacted, embedded or transposed left quadrant, trauma/injury to face or hand-wrist region, growth disorder/systemic illness. Data collection period not clear.	
Sample, sex and age	180 participants, 90 boys and 90 girls, aged 6-16 years.	
Study design	Cross-sectional.	
Index test	Radiograph of the right hand and wrist, Greulich & Pyle atlas.	
Comment	Subjects also scored according Demirjian's and Willem's method of dental age estimation.	
Aim of the study	“To test the applicability of Demirjian's and Willem's dental age assessment methods as well as Greulich and Pyle skeletal age assessment method in children residing in the Gandhinagar district.”	
QUADAS-2 assessment	Rating	Comment

Patient selection method:	Random selection from patients. No discussion of uniform distribution into age cohorts found in article, but each sex and age cohort has almost the same number of participants.	
- Consecutive or random sample of patients?	Yes	Random
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Unclear	No description of blinding
DOMAIN 2: Index test interpretation	Unclear risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	All x-rayed children were included in analysis- assumed
DOMAIN 4: Patient flow and timing bias?	Low risk of bias	

Patil ST, Parchand MP, Meshram MM, Kamdi NY. Applicability of Greulich and Pyle skeletal age standards to Indian children. Forensic Science International. 2012;216(1):200.e1-4.		
Population: Country, ethnicity, place and year	India (Maharashtra). Children attending at the radiology department, Government medical college, Nagpur for x-ray of inferior extremity for injury. Healthy children of middle socio-economic status, with no chronic diseases, endocrinopathies, metabolic disorders or any other disease affecting skeletal maturation. Data collection 2007- 2009.	
Sample, sex and age	375 participants, 194 boys and 181 girls, aged 0-19 years.	
Study design	Cross-sectional.	
Index test	Radiograph of the left hand and wrist if possible, otherwise the right, Greulich & Pyle atlas.	
Aim of the study	“In this study we have to find out whether Indian children have advanced skeletal age or lag behind skeletally or same as American children (GP atlas). So that applicability of Greulich and Pyle atlas to Indian children can be determined.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method not specified. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	Description of binding.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	All x-rayed children included in analysis.

DOMAIN 4: Patient flow and timing bias?	Low risk of bias
--	-------------------------

Suri S, Prasad C, Tompson B, Lou W. Longitudinal comparison of skeletal age determined by the Greulich and Pyle method and chronologic age in normally growing children, and clinical interpretations for orthodontics. American Journal of Orthodontics & Dentofacial Orthopedics. 2013;143(1):50-60..		
Population: Country, ethnicity, place and year	Canada (Toronto). Described as White Canadian children. Selected from Burlington Growth Centre archives the Faculty of Dentistry. Selected if serial annual hand-wrist radiographs available with 2 consecutive radiographs more than 2 years apart, in age range 9-18 years. Normal facial growth, no/minimal orthodontic treatment and no syndrome or specific disease or growth abnormalities. Data collection period not clear.	
Sample, sex and age	68 participants, 35 boys and 33 girls, aged 9-18 y	
Study design	Longitudinal study design, but analysed as cross-sectional study in this systematic review.	
Index test	Radiograph of the hand and wrist, side not mentioned, Greulich & Pyle atlas.	
Aim of the study	"...to comprehensively analyze the concordance between skeletal and chronologic ages determined by using the Greulich and Pyle method at different ages in the preadolescent and adolescent periods, and to determine any age- and sex-related differences in the concordance."	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method unclear. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Unclear	
- Avoid inappropriate exclusions?	Yes	Assumed ok.
DOMAIN 1: Patient selection	Unclear risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	Description of blinding
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	
DOMAIN 4: Patient flow and timing bias?	Low risk of bias	

Tisè M, Mazzarini L, Fabrizzi G, Ferrante L, Giorgetti R, Tagliabruni A. Applicability of Greulich and Pyle method for age assessment in forensic practice on an Italian sample. International Journal of Legal Medicine. 2011;125(3):411-6.		
Population: Country, ethnicity, place and year	Italy (Central Italia). Described as Italian-Caucasian children. Trauma patients hospitalized in the Azienda Ospedaliero Universitaria Ospedali Riuniti di Ancona. X-rays with evidence of bone fracture or soft tissue abnormalities were discarded. No information about patients' growth disorders or other illnesses were available. Data collection 2006-2007.	
Sample, sex and age	484 participants, 359 boys and 125 girls, aged 11-19 years	
Study design	Cross-sectional, retrospective from archives.	

Index test	Radiograph of the hand and wrist, side not mentioned, Greulich & Pyle atlas.	
Aim of the study	Since the Greulich and Pyle method is one of the most frequently used in age estimation, the aim of this study was to assess the reproducibility and accuracy of the method on a large Italian sample of teenagers, to ascertain the applicability of the Atlas at the critical age thresholds of 14 and 18 years.	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method unclear. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	
- Avoid inappropriate exclusions?	Yes	Assumed ok
DOMAIN 1: Patient selection	Low risk	
DOMAIN 1: Extra questions on age cohorts and age range	<p>Number of participants in each age cohort varies from 8 to 67 observations among boys (highest numbers in age cohort 15 years) and from 8 to 19 observations among girls (highest numbers in age cohort 12 years). Age range 11-19 years.</p> <p>Our judgement is that the size of the age cohorts vary considerably and some age cohorts have very few observations. Age mimicry is likely to influence analyses. Age range may limit analyses of the Greulich & Pyle development stages for lower and upper teenage age cohorts.</p> <p>High risk of bias</p>	
- Index test interpreted without knowledge of CA?	Yes	Description of blinding.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	Assumed ok.
DOMAIN 4: Patient flow and timing bias?	Low risk of bias	

van Rijn RR, Lequin MH, Robben SG, Hop WC, van Kuijk C. Is the Greulich and Pyle atlas still valid for Dutch Caucasian children today? <i>Pediatric Radiology</i> . 2001;31(10):748-52.		
Population: Country, ethnicity, place and year	Nederland (Amsterdam). Described as Dutch Caucasian. Data collection in 1997.	
Sample, sex and age	572 participants, 278 boys and 294 girls, aged 5-20 years.	
Study design	Cross-sectional.	
Index test	Radiograph of the left hand and wrist, Greulich & Pyle atlas.	
Aim of the study	“To compare skeletal age and calendar age in a healthy Dutch Caucasian population in order to test the applicability of this specific population”.	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Selection method unclear. No discussion of uniform distribution into age cohorts found in article.	
- Consecutive or random sample of patients?	Yes	
- Avoid inappropriate exclusions?	Yes	Assumed ok
DOMAIN 1: Patient selection	Low risk	

DOMAIN 1: Extra questions on age cohorts and age range	<i>The relevant analyses (average skeletal age in development stages from the Greulich & Pyle atlas) were not part of the results presented in the article. Re-analysed for this systematic review based on access to original data granted by authors.</i> Number of participants in each age cohort varies. Our judgement is that the size of the age cohorts vary considerably and some age cohorts have very few observations. Age mimicry is likely to influence analyses. High risk of bias	
- Index test interpreted without knowledge of CA?	Yes	Description of blinding.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok.
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	Assumed ok.
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Zafar AM, Nadeem N, Husen Y, Ahmad MN. An appraisal of Greulich-Pyle Atlas for skeletal age assessment in Pakistan. JPMA - Journal of the Pakistan Medical Association. 2010;60(7):552-5.		
Population: Country, ethnicity, place and year	Pakistan. Ethnically diverse. Children examined for indication of trauma. Consecutive patients admitted to one hospital in Karachi, Pakistan for indication of trauma. Less or equal to 216 months of age at time of exposure. Subjects ever investigated for metabolic, growth or nutritional disorders, or with a height or weight chart < 5 th and > 95 th centile were excluded. Chronological age from hospital records. 17 poor quality scans and 268 duplicate identities were excluded. Final Data collected 2005-2008.	
Sample, sex and age	889 participants, Children of both genders, aged up to 216 months	
Study design	Cross-sectional.	
Index test	Hand-wrist radiographs, left hand if possible, otherwise right hand. Assessed according to the Greulich-Pyle atlas. GP using hand-wrist of the right hand	
Aim of the study	“To assess accurate skeletal age (SA) in clinical and medico-legal decisions using the Greulich-Pyle (GP) atlas and to see its applicability across diverse populations in Karachi.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	“Consecutive patients admitted”	
- Consecutive or random sample of patients?	Yes	
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	<i>The relevant analyses (average skeletal age in development stages from the Greulich & Pyle atlas) were not part of the results presented in the article. Re-analysed for this systematic review based on access to original data granted by authors.</i> Number of participants in each age cohort varies. Our judgement is that the size of the age cohorts vary considerably and some age cohorts have very few observations. Age mimicry is likely to influence analyses. Age range may limit analyses of the Greulich & Pyle development stages upper teenage age cohorts.	

	High risk of bias	
- Index test interpreted without knowledge of CA?	Yes	Blinded assessor.
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in the analysis?	Yes	
DOMAIN 4: Patient flow and timing bias?	Low risk of bias	

Öztürk F, Karataş OH, Mutaf HI, Babacan H. Bone age assessment: comparison of children from two different regions with the Greulich–Pyle method In Turkey. Austr J of Forensic Sciences 2016;48(6):694-703.		
Population: Country, ethnicity, place and year	Turkey (two geographically different regions), Central Anatolian and Eastern Anatolian Turkish Caucasian children. Patients attending the Department of Orthodontics of Cumhuriyet University, Faculty of Dentistry in Central Anatolia and Inonu University in Eastern Anatolia.	
Sample, sex and age	849 subjects, 375 boys and 514 girls, aged 9-17 years	
Study design	Cross-sectional	
Index test	Digital radiography of the left hand and wrist. Greulich & Pyle atlas.	
Aim of the study	“...to identify and compare bone age assessments of Turkish children in regions of Central Anatolia and Eastern Anatolia using the Greulich-Pyle (GP) method.”	
QUADAS-2 assessment	Rating	Comment
Patient selection method:	Randomly selected from patients. No discussion of uniform distribution into age cohorts found in article.	
	Rating	Comment
- Consecutive or random sample of patients?	Yes	Random
- Avoid inappropriate exclusions?	Yes	
DOMAIN 1: Patient selection	Low risk of bias	
DOMAIN 1: Extra questions on age cohorts and age range	Not relevant for included analyses.	
- Index test interpreted without knowledge of CA?	Yes	“by the same investigator in a blind assessment.”
DOMAIN 2: Index test interpretation	Low risk of bias	
- CA interpreted without knowledge of SA?	Yes	Assumed ok
DOMAIN 3: Reference standard	Low risk of bias	
- All patients included in analysis?	Yes	
DOMAIN 4: Patient flow and timing bias	Low risk of bias	

Appendix 3: Relevant datasets using the Greulich and Pyle atlas

In addition to the 19 included studies, we identified 107 studies with data on skeletal maturation based on the Greulich & Pyle atlas and chronological age. The population criteria appear fulfilled, i.e. data for > 50 healthy children in age range 10-26 years. These studies were not included because the article did not provide the relevant figures or format on the data presented. Some of these studies were analysed to another problem than ours. We have gathered information about these studies because they may represent potential relevant datasets for further analyses. Some of these studies have used more than one scoring method on the same subjects as described.

Reference	Scoring method	Country	Participants	Sex	Age range
Alhadlaq AM, Al-Shayea EI. New method for evaluation of cervical vertebral maturation based on angular measurements. Saudi Medical Journal. 2013;34(4):388-94.	G&P and Fishman's	Saudi Arabia	197	Boys	10-15 years
Andersen E. Comparison of Tanner-Whitehouse and Greulich-Pyle methods in a large scale Danish Survey. American Journal of Physical Anthropology. 1971;35(3):373-6.	G&P, TW1 and TW2	Denmark	1009	Both	7 - 18 years
Anderson DL, Thompson GW, Popovich F. Interrelationships of dental maturity, skeletal maturity, height and weight from age 4 to 14 years. Growth. 1975;39(4):453-62.	G&P	Denmark	232	Both	4 -14 years
Arat M, Koklu A, Ozdiler E, Rubenduz M, Erdogan B. Craniofacial growth and skeletal maturation: a mixed longitudinal study. European Journal of Orthodontics. 2001;23(4):355-61.	G&P	Turkey	78	Both	10-17 years
Arat ZM, Rubenduz M. Changes in dentoalveolar and facial heights during early and late growth periods: a longitudinal study. Angle Orthodontist. 2005;75(1):69-74.	G&P	Turkey	62	Both	10-17 years
Awais M, Nadeem N, Husen Y, Rehman A, Beg M, Khattak YJ. Comparison between Greulich-Pyle and Girdany-Golden methods for estimating skeletal age of children in Pakistan. Journal of the College of Physicians and Surgeons--Pakistan : JCPSP. 2014;24(12):889-93.	G&P, Girdany-Golden	Pakistan	283	Both	0 -18 years
Beit P, Peltomaki T, Schatzle M, Signorelli L, Patcas R. Evaluating the agreement of skeletal age assessment based on hand-wrist and cervical vertebrae radiography. American Journal of Orthodontics and Dentofacial Orthopedics. 2013;144(6):838-47.	G&P	Switzerland	730	Both	6 -18 years
Berst MJ, Dolan L, Bogdanowicz MM, Stevens MA, Chow S, Brandser EA. Effect of knowledge of chronologic age on the variability of pediatric bone age determined using the Greulich and Pyle standards. American Journal of Roentgenology. 2001;176(2):507-10.	G&P	USA	107	Both	2-20 years

Bezerra IS, Topolski F, Franca SN, Brucker MR, Fernandes A. Assessment of skeletal and dental ages of children and adolescents with type 1 diabetes mellitus. <i>Pesquisa Odontologica Brasileira = Brazilian Oral Research.</i> 2015;29.	G&P and Nolla	Brazil	82	Both	5 -15 years
Blanksby BA, Brogan WF, McKnight HP, Sprague PL. A skeletal age survey of primary school children in Western Australia. <i>Australian Paediatric Journal.</i> 1975;11(3):169-71.	G&P	Australia	594	Both	6 -12 years
Bouchard C, Leblanc C, Malina RM, Hollmann W. Skeletal age and submaximal working capacity in boys. <i>Annals of Human Biology.</i> 1978;5(1):75-8.	G&P	Germany	237	Boys	8 -18 years
Bouchard C, Malina RM, Hollmann W, Leblanc C. Relationships between skeletal maturity and submaximal working capacity in boys 8 to 18 years. <i>Medicine & Science in Sports.</i> 1976;8(3):186-90.	G&P	Germany	237	Boys	8 -18 years
Brown T, Grave KC. Skeletal maturation in Australian Aborigines. <i>Australian Paediatric Journal.</i> 1976;12(1):24-30.	G&P	Australia	123	Both	5- 20 years
Buyukgebiz A, Eroglu Y, Karaman O, Kinik E. Height and weight measurements of male Turkish adolescents according to biological maturation. <i>Acta Paediatrica Japonica.</i> 1994;36(1):80-3.	G&P	Turkish, in USA	879	Boys	11-19 years
Calfee RP, Sutter M, Steffen JA, Goldfarb CA. Skeletal and chronological ages in American adolescents: current findings in skeletal maturation. <i>Journal of Childrens Orthopaedics.</i> 2010;4(5):467-70.	G&P	USA	138	Both	12-18 years
Canals M, Valenzuela C, Avendano A, Samith S. [Bone maturity in children older than 6 years of age. Wrist and hand. I. Quantitative analysis]. <i>Revista Chilena de Pediatria.</i> 1988;59(2):102-5.	G&P, TW2	Chile	1323	Both	6 -20 years
Carling C, Le Gall F, Malina RM. Body size, skeletal maturity, and functional characteristics of elite academy soccer players on entry between 1992 and 2003. <i>Journal of Sports Sciences.</i> 2012;30(15):1683-93.	G&P	France	158	Boys	13 years
Chaudhry, K., A. Agarwal, and U. Rehani, Interrelationship among Dental, Skeletal and Chronological Ages in Urban and Rural Female Children. <i>Jaypee's International Journal of Clinical Pediatric Dentistry,</i> 2010. 3(2): p. 79-86.	G&P	India	80	Both	8-14 years
Chaumoitre K, Lamtali S, Baali A, Saliba-Serre B, Lahmam A, Aboussad A, et al. Influence of socioeconomic status and body mass index on bone age. <i>Hormone Research in Paediatrics.</i> 2010;74(2):129-35.	G&P	Morocco	623	Both	6 -19 years
Chen ML, Chiang CH, Chen JS. Skeletal maturation and dietary intakes of Chinese children and young adults. <i>Chinese Medical Journal (Taipei).</i> 1977;24(2):132-41.	G&P	Chinese	4983	Both	2 -20 years

Cole AJ, Webb L, Cole TJ. Bone age estimation: a comparison of methods. <i>British Journal of Radiology</i> . 1988;61(728):683-6.	G&P, TW2	British	97	Both	mean 11 years
Coutinho S, Buschang PH, Miranda F. Relationships between mandibular canine calcification stages and skeletal maturity. <i>American Journal of Orthodontics & Dentofacial Orthopedics</i> . 1993;104(3):262-8.	G&P, TW & Demirjian	American	415	Both	7 -16 years
Cumming GR, Garand T, Borysyk L. Correlation of performance in track and field events with bone age. <i>Journal of Pediatrics</i> . 1972;80(6):970-3.	G&P	Canadian	259	Both	11 -18 years
De Donno A, Santoro V, Lubelli S, Marrone M, Lozito P, Intronà F. Age assessment using the Greulich and Pyle method on a heterogeneous sample of 300 Italian healthy and pathologic subjects. <i>Forensic Science International</i> . 2013;229(1):157.e1-6.	G&P	Italian	300	Both	10 -20 years
Ebri Torne B, Ebri Verde I. Comparative Study bone ages Greulich-Pyle Tanner-W2 and Ebri and between predicted adult height. <i>Pediatric Integral</i> . 2012;16(9):741.e1-e7.	G&P, TW2, Ebri	Spain	160	Both	0 -20 years
Edgren J, Fellman J, Lewin T. Skeletal maturation of the hand and wrist, a longitudinal study covering two years among children of Finnish Lapps. <i>Acta Morphologica Neerlando-Scandinavica</i> . 1974;12(1):1-7.	G&P	Finland	91	Both	3 -23 years
Fleshman K. Bone age determination in a paediatric population as an indicator of nutritional status. <i>Tropical Doctor</i> . 2000;30(1):16-8.	G&P	Nepalese	219		Unclear
Fry El. Tanner-Whitehouse and Greulich-Pyle skeletal age velocity comparisons. <i>American Journal of Physical Anthropology</i> . 1971;35(3):377-80.	G&P, TW	Hong Kong	233	Both	10 -16 years
Garamendi PM, Landa MI, Ballesteros J, Solano MA. Reliability of the methods applied to assess age minority in living subjects around 18 years old. A survey on a Moroccan origin population. <i>Forensic Science International</i> . 2005;154(1):3-12.	G&P & Demirjian	Morocco	114	Both	13 -25 years
Goldfarb CA, Strauss NL, Wall LB, Calfee RP. Defining ulnar variance in the adolescent wrist: Measurement technique and interobserver reliability. <i>Journal of Hand Surgery</i> . 2011;36(2):272-7.	G&P	USA	138	Both	12-18 years
Goldstein HS. Skeletal maturity and cognitive development of 12- to 17-year-old males. <i>Developmental Medicine & Child Neurology</i> . 1987;29(3):348-50.	G&P	USA	3516	Boys	12 -17 years
Gross GW, Boone JM, Bishop DM. Pediatric skeletal age: determination with neural networks. <i>Radiology</i> . 1995;195(3):689-95.	G&P	USA	521	Both	0 -18 years
Haavikko K, Kilpinen E. Skeletal development of Finnish children in the light of hand-wrist roentgenograms. <i>Proceedings of the Finnish Dental Society</i> . 1973;69(5):182-90.	G&P	Finland	1061	Both	2,5 -19 years

Hackman L, Black S. The reliability of the Greulich and Pyle atlas when applied to a modern Scottish population. <i>Journal of Forensic Sciences</i> . 2013;58(1):114-9.	G&P	UK	406	Both	0 -21 years
Haiter-Neto F, Kurita LM, Menezes AV, Casanova MS. Skeletal age assessment: A comparison of 3 methods. <i>American Journal of Orthodontics and Dentofacial Orthopedics</i> . 2006;130(4):435.e15-.e20.	G&P, TW3, Ek-løf & Ringertz	Brazil	360	Both	7 -15 years
Hala, L.A., et al., Comparison of accuracy between dental and skeletal age in the estimation of chronological age of Down syndrome individuals. <i>Forensic Science International</i> , 2016. 266: p. 578.e1-578.e10.	G&P	Brazil	216	Both	3-17 years
Hawley NL, Rousham EK, Norris SA, Pettifor JM, Cameron N. Secular trends in skeletal maturity in South Africa: 1962-2001. <i>Annals of Human Biology</i> . 2009;36(5):584-94.	G&P	South Africa	200	Both	9 -11 years
Heppe DH, Taal HR, Ernst GD, Van Den Akker EL, Lequin MM, Hokken-Koelega AC, et al. Bone age assessment by dual-energy X-ray absorptiometry in children: an alternative for X-ray? <i>British Journal of Radiology</i> . 2012;85(1010):114-20.	G&P and dual energy x-ray	Nether-land	95	Both	0-16 years
Kapalin V, Picko V. [the Skeletal Age of School Children and Its Dependences]. <i>Ceskoslovenska Hygiena</i> . 1964;9:342-56.	G&P	Czech	2500	Both	7 -15 years
Khan KM, Miller BS, Hoggard E, Soman A, Sarafoglou K. Application of ultrasound for bone age estimation in clinical practice. <i>Journal of Pediatrics</i> . 2009;154(2):243-7.	G&P, TW3	USA	100	Both	Mean age 10 years
Kim HJ, Yoon JR, Modi C, Modi H, Song HR, Song SY. Interrelationship of the Risser sign, knee epiphysis, and bone age in determining skeletal maturity: a case-control study. <i>Journal of Pediatric Orthopaedics, Part B</i> . 2011;20(3):173-7.	G&P, TW3	South Korea	293	Both	9 -18 years
Kim JR, Lee YS, Yu J. Assessment of bone age in pre-pubertal healthy Korean children: comparison among the Korean standard bone age chart, Greulich-Pyle method, and Tanner-Whitehouse method. <i>Korean Journal of Radiology</i> . 2015;16(1):201-5.	G&P, TW3 and KS (Korean standard)	South Korea	212	Both	7 -12 years
Kimura K. Age estimation from second metacarpals in children. <i>Okajimas Folia Anatomica Japonica</i> . 1992;69(4):177-82.	G&P	Japan	1241	Both	0 -19 years
Kimura K. Skeletal maturation in Japanese as assessed by the Oxford and Tanner-Whitehouse methods. <i>Kaibogaku zasshi</i> . 1972;0(5):358-72.	G&P	Japanese	598	Both	0-18 years
Kopczynska J. [Further studies on the skeletal age evaluated according to examination of bones of the hand and wrist in children in Warsaw. I]. <i>Pediatria Polska</i> . 1961;36:845-53.	G&P	Polish	5396	Both	7 -16 years
Kopczynska J. [Studies on developmental age of school children according to established skeletal age]. <i>Pediatria Polska</i> . 1959;34:859-70.	G&P	Polish	278	Both	6 -15 years

Krailassiri S, Anuwongnukroh N, Dechkunakorn S. Relationships between dental calcification stages and skeletal maturity indicators in Thai individuals. Angle Orthodontist. 2002;72(2):155-66.	G&P AND Demirjian	Thailand	361	Both	7 -19 years
Kullman L. Accuracy of two dental and one skeletal age estimation method in Swedish adolescents. Forensic Science International. 1995;75(2):225-36.	G&P	Sweden	72	Both	12 -19 years
Lee MM, Chan ST, Low WD, Chang KS. The relationship between dental and skeletal maturation in Chinese children. Archives of Oral Biology. 1965;10(6):883-91.	G&P	South China	5560	Both	6 -14 years
Lee MM. Maturation disparity between hand-wrist bones in Hong Kong Chinese children. American Journal of Physical Anthropology. 1971;34(3):385-95.	G&P	Hong Kong	571	Both	0 -17 years
Lenko HL. Prediction of adult height with various methods in Finnish children. Acta Paediatrica Scandinavica. 1979;68(1):85-92.	G&P, Tanner, Bayley&P, Walker, RWT	Finland	60	Both	7 -19 years
Lewis AB. Comparisons between dental and skeletal ages. Angle Orthodontist. 1991;61(2):87-92.	G&P	Unclear	694	Both	10 years old
Lewis CP, Lavy CBD, Harrison WJ. Delay in skeletal maturity in Malawian children. Journal of Bone and Joint Surgery - Series B. 2002;84(5):732-4.	G&P	Malawi	130	Both	2-28 years
Lin FQ, Zhang J, Zhu Z, Wu YM. Comparative study of Gilsanz-Ratib digital atlas and Greulich-Pyle atlas for bone age estimation in a Chinese sample. Annals of Human Biology. 2015;42(6):523-7.	G&P, Gilsanz-Ratib digital atlas	China	618	Both	0 -13 years
Lin NH, Ranjitkar S, Macdonald R, Hughes T, Taylor JA, Townsend GC. New growth references for assessment of stature and skeletal maturation in Australians. Australian Orthodontic Journal. 2006;22(1):1-10.	G&P	South Australia	2497	Both	9 -18 years
Linhares RV, Matta Mde O, Lima JR, Dantas PM, Costa MB, Fernandes Filho J. [Effects of sexual maturation on body composition, dermatoglyphics, somatotype and basic physical qualities of adolescents]. Arquivos Brasileiros de Endocrinologia e Metabologia. 2009;53(1):47-54.	G&P	Brazil	136	Boys	10 -14 years
Loder RT, Estle DT, Morrison K, Eggleston D, Fish DN, Greenfield ML, et al. Applicability of the Greulich and Pyle skeletal age standards to black and white children of today. American Journal of Diseases of Children. 1993;147(12):1329-33.	G&P	USA	841	Both	0 -18 years
Low WD. Relation between socio-economic status and skeletal maturation of Chinese children. Zeitschrift fur Morphologie und Anthropologie. 1972;64(1):1-11.	G&P	China	33990	Both	6 -20 years

Maggio, A., et al., Assessment of the accuracy of the Greulich and Pyle hand-wrist atlas for age estimation in a contemporary Australian population. Australian Journal of Forensic Sciences, 2016: p. 1-11.	G&P	Australia	360	Both	0-25 years
Makkad RS, Balani A, Chaturvedi SS, Tanwani T, Agrawal A, Hamdani S. Reliability of panoramic radiography in chronological age estimation. Journal of forensic dental sciences : JFDS. 2013;5(2):129-33.	G&P, Demirjian	India	270	Both	17 -25 years
Malik P, Rana V, Rehani U. To Evaluate the Relationship between Mandibular Canine Calcification Stages and Skeletal Age. Jaypees International Journal of Clinical Pediatric Dentistry. 2012;5(1):14-9.	G&P, Demirjian	India	147	girls	10 -13 years
Mansourvar M, Ismail MA, Raj RG, Kareem SA, Aik S, Gunalan R, et al. The applicability of Greulich and Pyle atlas to assess skeletal age for four ethnic groups. Journal of Forensic and Legal Medicine. 2014;22:26-9.	G&P	USA	184	Boys	1 -18 years
Maresh MM. Single versus serial assessment of skeletal age: either, both or neither? American Journal of Physical Anthropology. 1971;35(3):387-92.	G&P	USA	104	Both	10 -15 years
Mathiasen MS. Determination of bone age and recording of minor skeletal hand anomalies in normal children. Danish Medical Bulletin. 1973;20(3):80-5.	G&P	Denmark	1100	Both	1 -16 years
Mazess RB, Cameron JR. Skeletal growth in school children: maturation and bone mass. American Journal of Physical Anthropology. 1971;35(3):399-407.	G&P	USA	322	Both	6 -12 years
McCormack, S.E., et al., Relative Skeletal Maturation and Population Ancestry in Nonobese Children and Adolescents. Journal of Bone & Mineral Research, 2017. 32(1): p. 115-124.	G&P	USA	1592	Both	5-24 years
Michaut E, Niang I, Dan V. Bone maturation during puberty. Apropos of 227 adolescents from Dakar. [French]. Annales de radiologie. 1972;15(9):767-79.	G&P, TW	Senegal	227	Both	11 -15 years
Mora S, Boechat MI, Pietka E, Huang HK, Gilsanz V. Skeletal age determinations in children of European and African descent: applicability of the Greulich and Pyle standards. Pediatric Research. 2001;50(5):624-8.	G&P	USA	534	Both	0 -19 years
Moradi M, Sirous M, Morovatti P. The reliability of skeletal age determination in an Iranian sample using Greulich and Pyle method. Forensic Science International. 2012;223(1):372.e1-4.	G&P	Iran	425	Both	6 -18 years
Murthy KK, Srinivas CN, Lakshmi V, Kumar CV, Krishnaveni M. Assessment of skeletal and dental maturity levels for a given chronological age among Indian children. Journal of Contemporary Dental Practice [Electronic Resource]. 2012;13(3):310-5.	G&P, Nolla	India	260	Both	3 -15 years
Nahid G, Abdorrahim A, Gharib SM, Anvar E. Assessment of bone age in Kurdish children in IRAN. Pakistan Journal of Medical Sciences. 2010;26(1):36-9.	G&P	Kurdish	228	Both	7 -14 years
Napoli MM, Saraiva PA. Bone age. [Portuguese]. Revista do Hospital das Clinicas. 1985;40(5):210-5.	G&P	Brazil	217	Both	0 -20 years

Oh Y, Lee R, Kim HS. Evaluation of skeletal maturity score for Korean children and the standard for comparison of bone age and chronological age in normal children. <i>Journal of Pediatric Endocrinology and Metabolism</i> . 2012;25(3):279-84.	G&P and TW3	South Korea	378	Both	8-15 years
Onat T. Prediction of adult height of girls based on the percentage of adult height at onset of secondary sexual characteristics, at chronological age, and skeletal age. <i>Human Biology</i> . 1975;47(1):117-30.	G&P	Turkish	119	Girls	8.5-13.4 years
Ontell FK, Ivanovic M, Ablin DS, Barlow TW. Bone age in children of diverse ethnicity. <i>American Journal of Roentgenology</i> . 1996;167(6):1395-8.	G&P	USA	765	Both	3-18 years
Pawson IG. Radiographic determination of excessive bone loss in Alaskan Eskimos. <i>Human Biology</i> . 1974;46(3):369-80.	G&P	USA	154	Both	0->50 years
Paxton ML, Lamont AC, Stillwell AP. The reliability of the Greulich-Pyle method in bone age determination among Australian children. <i>Journal of Medical Imaging and Radiation Oncology</i> . 2013;57(1):21-4.	G&P	Australia	350	Both	0-18 years
Pinchi V, De Luca F, Focardi M, Pradella F, Vitale G, Ricciardi F, et al. Combining dental and skeletal evidence in age classification: Pilot study in a sample of Italian sub-adults. <i>Legal Medicine</i> . 2016;20:75-9.	G&P, TW3	Italy	274	Both	6-17 years
Pinchi V, De Luca F, Ricciardi F, Focardi M, Piredda V, Mazzeo E, et al. Skeletal age estimation for forensic purposes: A comparison of GP, TW2 and TW3 methods on an Italian sample. <i>Forensic Science International</i> . 2014;238:83-90.	G&P, TW3, TW2	Italy	307	Both	6-20 years
Rai V, Saha S, Yadav G, Tripathi AM, Grover K. Dental and skeletal maturity- A biological indicator of chronologic age. <i>Journal of Clinical and Diagnostic Research</i> . 2014;8(9):ZC60-ZC4.	G&P, TW2	India	150	Both	5-15 years
Rikhasor RM, Qureshi AM, Rathi SL, Channa NA. Skeletal maturity in Pakistani children. <i>Journal of Anatomy</i> . 1999;195:305-8.	G&P	India	750	Both	1-18 years
Roche AF, Roberts J, Hamill PV. Skeletal maturity of children 6-11 years, United States. <i>Vital & Health Statistics - Series 11: Data From the National Health Survey</i> . 1974;0(140):1-62.	G&P	USA	Ca 7000	Both	6-11 years
Roche AF, Roberts J, Hamill PV. Skeletal maturity of children 6-11 years: racial, geographic area, and socioeconomic differentials, United States. <i>Vital & Health Statistics - Series 11: Data From the National Health Survey</i> . 1975;0(149):1-81.	G&P	USA	Ca 7000	Both	6-11 years
Roche AF, Roberts J, Hamill PV. Skeletal maturity of youths 12--17 years racial, geographic area, and socioeconomic differentials. United States, 1966-1970. <i>Vital & Health Statistics - Series 11: Data From the National Health Survey</i> . 1978;0(167):1-98.	G&P	USA		Both	12-17 years
Roche AF, Roberts J, Hamill PV. Skeletal maturity of youths 12-17 years, United States. <i>Vital & Health Statistics - Series 11: Data From the National Health Survey</i> . 1976;0(160):1-90.	G&P	USA	Ca 7000	Both	12-17 years

Roche AF, Roberts J, Hamill PVV. Skeletal maturity of children 6-11 years. <i>Vital and Health Statistics Ser.</i> 1974;11(140).	G&P	USA	Ca 7000	Both	6-11 years
Russell DL, Keil MF, Bonat SH, Uwaifo GI, Nicholson JC, McDuffie JR, et al. The relation between skeletal maturation and adiposity in African American and Caucasian children. <i>Journal of Pediatrics.</i> 2001;139(6):844-8.	G&P	USA	252	Both	5-12 years
Santoro V, Roca R, De Donno A, Fiandaca C, Pinto G, Tafuri S, et al. Applicability of Greulich and Pyle and Demirjian aging methods to a sample of Italian population. <i>Forensic Science International.</i> 2012;221(1):153.e1-5.	G&P	Italian	532	Both	7-15 years
Santos C, Ferreira M, Alves FC, Cunha E. Comparative study of Greulich and Pyle Atlas and Maturos 4.0 program for age estimation in a Portuguese sample. <i>Forensic Science International.</i> 2011;212(1):276.e1-7.	G&P	Portugal	230	Both	12-20 years
Schedewie H, Braselman A, Willich E, Lachman R. The determination of bone age in the elbow as compared to the hand. A study in 390 children. <i>Revista Interamericana de Radiologia.</i> 1979;4(1):11-7.	G&P	Germany	408	Both	3-18 years
Schmidt S, Koch B, Schulz R, Reisinger W, Schmeling A. Comparative analysis of the applicability of the skeletal age determination methods of Greulich-Pyle and Thiemann-Nitz for forensic age estimation in living subjects. <i>International Journal of Legal Medicine.</i> 2007;121(4):293-6.	G&P, Thieman n & Nitz	Germany	349	Both	1-18 years
Schmidt S, Koch B, Schulz R, Reisinger W, Schmeling A. Studies in use of the Greulich-Pyle skeletal age method to assess criminal liability. <i>Legal Medicine.</i> 2008;10(4):190-5.	G&P	Germany	649	Both	1-18 years
Scoles PV, Salvagno R, Villalba K, Riew D. Relationship of iliac crest maturation to skeletal and chronologic age. <i>Journal of Pediatric Orthopedics.</i> 1988;8(6):639-44.	G&P	USA	388	Both	10-19 years
Shaikh AH, Rikhasor RM, Qureshi AM. Determination of skeletal age in children aged 8-18 years. <i>Jpmo.</i> 1998;0(4):104-6.	G&P	Pakistan	402	Both	8-18 years
Shilpa PH, Sunil RS, Sapna K, Kumar NC. Estimation and comparison of dental, skeletal and chronologic age in Bangalore south school going children. <i>Journal of the Indian Society of Pedodontics & Preventive Dentistry.</i> 2013;31(2):63-8 6p.	G&P	India	250	Both	6-15 years
So LL, Yen PK. Secular trend in skeletal maturation in southern Chinese girls in Hong Kong. <i>Zeitschrift fur Morphologie und Anthropologie.</i> 1990;78(2):145-53.	G&P	Hong Kong	117	Girls	11-12 years
So LL. Correlation of sexual maturation with skeletal age of southern Chinese girls. <i>Australian Orthodontic Journal.</i> 1997;14(4):215-7.	G&P	Hong Kong	117	Girls	11-12 years

So LL. Correlation of skeletal maturation with stature and body weight of southern Chinese girls in Hong Kong. Zeitschrift fur Morphologie und Anthropologie. 1991;78(3):307-12.	G&P	Hong Kong	117	Girls	11-12 years
So LL. Skeletal maturation of the hand and wrist and its correlation with dental development. Australian Orthodontic Journal. 1997;15(1):1-9.	G&P	China	117	Girls	12 years
Soudack M, Ben-Shlush A, Jacobson J, Raviv-Zilka L, Eshel I, Hamiel O. Bone age in the 21st century: is Greulich and Pyle's atlas accurate for Israeli children? Pediatric Radiology. 2012;42(3):343-8.	G&P	Israel	679	Both	0-18 years
Takatama H. Roentgenological studies on the skeletal development of the hand and wrist in Japanese and Ainu children. [Japanese]. Sapporo Medical Journal. 1979;48(3):260-84.	G&P	Japan	724	Both	6-15 years
Van Rijn RR, Lequin MH, Robben SGF, Hop WCJ, Van Kuijk C. The Greulich&Pyle atlas for determining the skeletal age can still be used in a contemporary Dutch white population. [Dutch]. Nederlands Tijdschrift voor Geneeskunde. 2003;147(15):701-4.	G&P	Nether-land	572	Both	5-20 years
Vignolo M, Milani S, DiBattista E, Naselli A, Mostert M, Aicardi G. Modified Greulich-Pyle, Tanner-Whitehouse, and Roche-Wainer-Thissen (knee) methods for skeletal age assessment in a group of Italian children and adolescents. European Journal of Pediatrics. 1990;149(5):314-7.	G&P, TW2	Italy	221	Both	4-17 years
Zabet D, Rerolle C, Pucheu J, Telmon N, Saint-Martin P. Can the Greulich and Pyle method be used on French contemporary individuals? International Journal of Legal Medicine. 2015;129(1):171-7.	G&P	France	190	Both	10-19 years
Zammit MP, Kalra V, Nelson S, Broadbent BH, Hans MG. Growth patterns of Labrador Inuit youth: II. Skeletal age. Arctic Medical Research. 1994;53(4):176-83.	G&P	USA	100	Both	5-18 years
Zhang A, Sayre JW, Vachon L, Liu BJ, Huang HK. Racial differences in growth patterns of children assessed on the basis of bone age. Radiology. 2009;250(1):228-35.	G&P	China	1390	Both	0-18 years
Zhang J, Lin F, Ding X. Maturation Disparity between Hand-Wrist Bones in a Chinese Sample of Normal Children: An Analysis Based on Automatic BoneXpert and Manual Greulich and Pyle Atlas Assessment. Korean Journal of Radiology. 2016;17(3):435-42.	G&P	China	397	Both	2-14 years

Appendix 4: Relevant datasets using other index tests than Greulich and Pyle atlas to classify bone age on hand x-rays

In addition to the studies based on the Greulich and Pyle atlas, we identified 124 studies that had collected data on hand skeletal maturation based on other classification systems and chronological age. The population criteria appear fulfilled, i.e. data for > 50 healthy children in age range 10-26 years. Some of these studies were analysed to another objective than ours. We have gathered information about these studies because they may represent potential relevant datasets for further analyses.

Reference	Scoring method	Country	Participants	Sex	Age range
Agathos D, Markostamos K, Toutountjakis N. Statural, bone and dental growth in Greek girls of the Athenian region. [French]. L' Orthodontie française. 1987;58:507-16.	Tanner	Greece	151	Girls	6-17 years
Agossou-Voyeme AK, Fachehou CR, Boco V, Hounnou GM, Biaou O. [Osseus age of the black children of Benin. A population study of 600 children aged from 9 to 18 years and living in Cotonou]. Morphologie. 2005;89(285):64-70.	TW2	Benin	600	Both	9-18 years
Aguiar LB, Caldas Mde P, Harter Neto F, Ambrosano GM. A methodology to measure cervical vertebral bone maturation in a sample from low-income children. Brazilian Dental Journal. 2013;24(1):30-4.	TW3	Brazil	ca 80	Both	7-15 years
Akrige M, Hilgers KK, Silveira AM, Scarfe W, Scheetz JP, Kinane DF. Childhood obesity and skeletal maturation assessed with Fishman's hand-wrist analysis. American Journal of Orthodontics and Dentofacial Orthopedics. 2007;132(2):185-90.	Fishman	USA	109	Both	9-16 years
Alkhal HA, Wong RW, Rabie AB. Correlation between chronological age, cervical vertebral maturation and Fishman's skeletal maturity indicators in southern Chinese. Angle Orthodontist. 2008;78(4):591-6.	Fishman	Chinese	400 (200x2)	Both	girls 10-15, boys 12 - 17
Al-Qtaitat, A., et al., Bone age determination of epiphyseal union around wrist joint and its correlation with chronological age: A radiological study in a Jordanian population. Biosciences Biotechnology Research Asia, 2016. 13(1): p. 67-73.	McKern & Stewart	Jordan	101	Both	12-22 years
Andersen E. Skeletal maturation of Danish school children in relation to height, sexual development, and social conditions. Acta Paediatrica Scandinavica. 1968;0:Suppl 185:11+.	Various methods	Denmark	ca 700		
Ashizawa K, Asami T, Anzo M, Matsuo N, Matsuoka H, Murata M, et al. Standard RUS skeletal maturation of Tokyo children. Annals of Human Biology. 1996;23(6):457-69.	TW2	Japan	1457	Both	3-18 years
Ashizawa K, Kumakura C, Zhou X, Jin F, Cao J. RUS skeletal maturity of children in Beijing. Annals of Human Biology. 2005;32(3):316-25.	TW3	China	1273	Both	6-8 years
Baughan B, Demirjian A, Levesque GY. Skeletal maturity standards for French-Canadian children of school-age with a discussion of the reliability and validity of such measures. Human Biology. 1979;51(3):353-70.	TW2	Canada	4084	Both	6-17 years

Benso L, Vannelli S, Pastorin L, Benso A, Milani S. Variation of bone age progression in healthy children. <i>Acta Paediatrica Supplement</i> . 1997;423:109-12.	TW2	Italy	407	Boys	7-12 years
Beunen G, Lefevre J, Ostyn M, Renson R, Simons J, Van Gerven D. Skeletal maturity in Belgian youths assessed by the Tanner-Whitehouse method (TW2). <i>Annals of Human Biology</i> . 1990;17(5):355-76.	TW2	Belgium	21174 boys, 9698 girls	Both	boys 12-20 years, girls 6-19 years
Beunen G, Malina RM, Ostyn M, Renson R, Simons J, Van Gerven D. Fatness and skeletal maturity of Belgian boys 12 through 17 years of age. <i>American Journal of Physical Anthropology</i> . 1982;59(4):387-92.	TW1+2	Belgium	14259	Boys	12-17 years
Beunen G, Ostyn M, Renson R, Simons J, Swalus P, Van Gerven D. Skeletal maturation and physical fitness of 12 to 15 year old boys. <i>Acta Paediatrica Belgica</i> . 1974;28:221-32.	TW and Healy	Belgium	7000	Boys	12 to 15
Beunen G, Ostyn M, Simons J, Renson R, Van Gerven D. Chronological and biological age as related to physical fitness in boys 12 to 19 years. <i>Annals of Human Biology</i> . 1981;8(4):321-31.	TW2	Belgium	21052	Boys	12 to 19
Beunen GP, Malina RM, Lefevre JA, Claessens AL, Renson R, Vanreusel B. Adiposity and biological maturity in girls 6-16 years of age. <i>International Journal of Obesity</i> . 1994;18(8):542-6.	TW	Belgium	6029	Girls	6 to 16
Bhat VJ, Kamath GP. Age estimation from root development of mandibular third molars in comparison with skeletal age of wrist joint. <i>American Journal of Forensic Medicine and Pathology</i> . 2007;28(3):238-41.	Kangne et al criteria AND Kullmann	India	735	Both	15 to 25
Briers PJ, Hoorweg J, Stanfield JP. The long-term effects of protein energy malnutrition in early childhood on bone age, bone cortical thickness and height. <i>Acta Paediatrica Scandinavica</i> . 1975;64(6):853-8.	TW	Uganda	72	Both	11 to 17 years
Buken B, Buken E, Safak AA, Yazici B, Erkol Z, Mayda AS. Is the "Gok Atlas" sufficiently reliable for forensic age determination of Turkish children? <i>Turkish Journal of Medical Sciences</i> . 2008;38(4):319-27.	Gok atlas	Turkey	772	Both	11 to 22
Buken B, Erzengin OU, Buken E, Safak AA, Yazici B, Erkol Z. Comparison of the three age estimation methods: which is more reliable for Turkish children? <i>Forensic Science International</i> . 2009;183(1):103.e1-7.	Various methods	Turkey	333	Both	11 to 16
Buken B, Safak AA, Buken E, Yazici B, Erkol Z, Erzengin OU. Is the tanner-whitehouse (TW3) method sufficiently reliable for forensic age determination of Turkish children?. [Turkish]. <i>Turkish Journal of Medical Sciences</i> . 2010;40(5):797-805.	TW3	Turkey	324	Both	girls 11 to 16, boys 11 to 16
Caldas Mde P, Ambrosano GM, Haite-Neto F. Use of cervical vertebral dimensions for assessment of children growth. <i>Journal of Applied Oral Science</i> . 2007;15(2):144-7.	TW3	Brazil	238	Both	7 to 16
Caltabiano M, Leonardi R, Zaborra G. [Evaluation of cervical vertebrae for determination of skeletal age]. <i>Rivista Italiana di Odontoiatria Infantile</i> . 1990;1(3):15-20.	Demirjian, unknown hand-method and cervical vertebrae	Italy	72	Both	10 to 15

Camacho-Basallo, P., et al., Five radiographic methods for assessing skeletal maturity in a Spanish population: is there a correlation? <i>Acta Odontologica Scandinavica</i> , 2016; p. 1-7.	Björk, Grave & Brown	Spain	200	Both	13 years
Cameriere R, De Luca S, Biagi R, Cingolani M, Farronato G, Ferrante L. Accuracy of three age estimation methods in children by measurements of developing teeth and carpal and epiphyses of the ulna and radius. <i>Journal of Forensic Sciences</i> . 2012;57(5):1263-70.	Cameriere Ferrante	Italy	288	Both	5 to 15
Cameriere R, Ferrante L, Ermenc B, Mirtella D, Strus K. Age estimation using carpal: study of a Slovenian sample to test Cameriere's method. <i>Forensic Science International</i> . 2008;174(2):178-81.	Cameriere & TW3	Slovenia	158	Both	6 to 16
Cameriere R, Ferrante L, Mirtella D, Cingolani M. Carpal and epiphyses of radius and ulna as age indicators. <i>International Journal of Legal Medicine</i> . 2006;120(3):143-6.	TW3	Italy	150	Both	5 to 17
Cameriere R, Ferrante L. Age estimation in children by measurement of carpal and epiphyses of radius and ulna and open apices in teeth: a pilot study. <i>Forensic Science International</i> . 2008;174(1):60-3.	Own method	Italy	150	Both	5 to 15
Chang HP, Liao CH, Yang YH, Chang HF, Chen KC. Correlation of cervical vertebra maturation with hand-wrist maturation in children. <i>Kaohsiung Journal of Medical Sciences</i> . 2001;17(1):29-35.	Fishman	Taiwan	503	Both	8 to 18
Chen LL, Xu TM, Jiang JH, Zhang XZ, Lin JX. Quantitative cervical vertebral maturation assessment in adolescents with normal occlusion: A mixed longitudinal study. <i>American Journal of Orthodontics and Dentofacial Orthopedics</i> . 2008;134(6):720.e1-e7.	Fishman	USA	87	Both	8 to 18
Coelho ESMJ, Figueiredo AJ, Simoes F, Seabra A, Natal A, Vaeyens R, et al. Discrimination of u-14 soccer players by level and position. <i>International Journal of Sports Medicine</i> . 2010;31(11):790-6.	Fels	Portugal	128	Boys	13 to 14
Cole TJ, Rousham EK, Hawley NL, Cameron N, Norris SA, Pettifor JM. Ethnic and sex differences in skeletal maturation among the Birth to Twenty cohort in South Africa. <i>Archives of Disease in Childhood</i> . 2015;100(2):138-43.	TW3	South Africa	607	Both	9 to 20
Cunha P, Moura DC, Guevara Lopez MA, Guerra C, Pinto D, Ramos I. Impact of ensemble learning in the assessment of skeletal maturity. <i>Journal of Medical Systems</i> . 2014;38(9):87.	Own method	Portugal	887	Both	0 to 20
De Luca S, Mangiulli T, Merelli V, Conforti F, Velandia Palacio LA, Agostini S, et al. A new formula for assessing skeletal age in growing infants and children by measuring carpal and epiphyses of radio and ulna. <i>Journal of Forensic and Legal Medicine</i> . 2016;39:109-16.	Own system	Italy	204	Both	1 to 16
Dhar S, Dangerfield PH. Skeletal maturity in normal children from Liverpool, United Kingdom. <i>Clinical Anatomy</i> . 1992;5(6):458-65.	TW2	UK	273	Both	2 to 17
El-Bakary AA, Attalla SM, Hammad SM, El-Ashry RA, De Luca S, Ferrante L, et al. Age estimation in Egyptian children by measurements of carpal and epiphyses of the ulna and radius. <i>Journal of Forensic Radiology and Imaging</i> . 2014;2(3):121-5.	Cameriere	Egypt	257	4 to 18	2 to 18
Engstrom C, Engstrom H, Sagne S. Lower third molar development in relation to skeletal maturity and chronological age. <i>Angle Orthodontist</i> . 1983;53(2):97-106.	Unclear, 5 stages	Sweden	221	Both	possibly 8 to 21

Figueiredo AJ, Goncalves CE, Coelho ESMJ, Malina RM. Youth soccer players, 11-14 years: maturity, size, function, skill and goal orientation. Annals of Human Biology. 2009;36(1):60-73.	Fels	Portugal	159	Boys	11- 14 years
Fishman LS. Maturational patterns and prediction during adolescence. Angle Orthodontist. 1987;57(3):178-93.	SMI	USA	2225 female, 1775 males	Both	9- 19 years
Fishman LS. Radiographic evaluation of skeletal maturation. A clinically oriented method based on hand-wrist films. Angle Orthodontist. 1982;52(2):88-112.	SMA method 4 stages	USA	170 fe-male, 164 males	Both	3 months to adult
Flores-Mir C, Burgess CA, Champney M, Jensen RJ, Pitcher MR, Major PW. Correlation of skeletal maturation stages determined by cervical vertebrae and hand-wrist evaluations. Angle Orthodontist. 2006;76(1):1-5.	Demirjian & Hagg and Taranger	Peru	140	Both	9,5 to 16,5
Flores-Mir C, Mauricio FR, Orellana MF, Major PW. Association between growth stunting with dental development and skeletal maturation stage. Angle Orthodontist. 2005;75(6):935-40.	Hägg & Taranger	Peru	140	Both	10-17 years
Freitas D, Maia J, Beunen G, Lefevre J, Claessens A, Marques A, et al. Skeletal maturity and socio-economic status in Portuguese children and youths: the Madeira growth study. Annals of Human Biology. 2004;31(4):408-20.	TW2, TW3	Portugal	507	Both	8- 6 years
Freitas D, Malina RM, Maia J, Lefevre J, Stasino-poulos M, Gouveia E, et al. Short-term secular change in height, body mass and Tanner-Whitehouse 3 skeletal maturity of Madeira youth, Portugal. Annals of Human Biology. 2012;39(3):195-205.	TW3	Portugal	2856	Both	4-17 years
Freitas DL, Lausen B, Maia JA, Gouveia ER, Thomis M, Lefevre J, et al. Skeletal Maturation, Body Size, and Motor Coordination in Youth 11-14 Years. Medicine & Science in Sports & Exercise. 2016;48(6):1129-35.	TW3	Portugal	613	Both	11-14 years
Gelbrich B, Frerking C, Weiss S, Schwerdt S, Stell-Izig-Eisenhauer A, Tausche E, et al. Combining wrist age and third molars in forensic age estimation: how to calculate the joint age estimate and its error rate in age diagnostics. Annals of Human Biology. 2015;42(4):389-96.	Thiemann-Nietz	Germany	383	Both	8 -19 years
Guimarey L, Moreno Morcillo A, Orazi V, Lemos-Marini SHV. Validity of the use of a few hand-wrist bones for assessing bone age. Journal of Pediatric Endocrinology and Metabolism. 2003;16(4):541-4.	TW2	Brazil, Argentina	205	Both	0,9- 17,4 years
Hauspie R, Bielicki T, Koniaruk J. Skeletal maturity at onset of the adolescent growth spurt and at peak velocity for growth in height: a threshold effect? Annals of Human Biology. 1991;18(1):23-9.	TW2	Poland	191	Boys	8- 19 years
Helm S. Relationship between dental and skeletal maturation in Danish schoolchildren. Scandinavian Journal of Dental Research. 1990;98(4):313-7.	TW2	Denmark	2744	Both	7- 14 years
Helm S. Skeletal maturity in Danish schoolchildren assessed by the TW2 method. American Journal of Physical Anthropology. 1979;51(3):345-52.	TW2	Denmark	3817	Both	7- 18 years
Himes JH, Huang Z, Haas JD, Rivera R, Pineda O. Serum alkaline phosphatase activity and skeletal maturation in Guatemalan adolescents. Annals of Human Biology. 1993;20(1):39-46.	TW2	Guate-mala	873	Both	11- 25 years
Houston WJ. Relationships between skeletal maturity estimated from hand-wrist radiographs and	TW2	UK	126	Both	8- 17 years

the timing of the adolescent growth spurt. European Journal of Orthodontics. 1980;2(2):81-93.					
Huo AH, Peng Y, Zeng JJ, Yu T, Li DH, Hu D. The CHN radiographic atlas method for assessing skeletal age of hand and wrist in 1397 children and result analysis. [Chinese]. Chinese Journal of Radiology (China). 2013;47(12):1074-6.	CHN atlas	China	1397	Both	1 - 18 years
Illich JZ, Hangartner TN, Skugor M, Roche AF, Goel PK, Matkovic V. Skeletal age as a determinant of bone mass in preadolescent females. Skeletal Radiology. 1996;25(5):431-9.	FELS	USA	456	Girls	8 -13 years
Jensen BL, Dahl E, Kreiborg S. Longitudinal study of body height, radius length and skeletal maturity in Danish boys with cleft lip and palate. Scandinavian Journal of Dental Research. 1983;91(6):473-81.	TW2	Denmark	85	Boys	6- 20 years
Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. BMJ. 2009;338:b490.	Fels	Uk	292	Boys	9- 16 years
Jones G, Ma D. Skeletal age deviation assessed by the Tanner-Whitehouse 2 method is associated with bone mass and fracture risk in children. Bone. 2005;36(2):352-7.	TW2	Australia	642	Both	9- 16 years
Joshi V, Yamaguchi T, Matsuda Y, Kaneko N, Maki K, Okano T. Skeletal maturity assessment with the use of cone-beam computerized tomography. Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. 2012;113(6):841-9.	GARCIA's table	Japan	100	Both	3- 35 years
Katzmarzyk PT, Malina RM, Beunen GP. The contribution of biological maturation to the strength and motor fitness of children. Annals of Human Biology. 1997;24(6):493-505.	TW2	USA	740	Both	7- 12 years
Kemper HC, Post GB, Twisk JW. Rate of maturation during the teenage years: nutrient intake and physical activity between ages 12 and 22. International Journal of Sport Nutrition. 1997;7(3):229-40.	TW2	Nether-land	200 longitudinal	Both	12- 22 years
Kimura K. Skeletal maturation in Japanese. A new analytical method (Japanese). Journal of the Anthropological Society of Nippon. 1972;80(4):319-36.	TW	Japan	598	Both	0 -18 years
Kimura K. Skeletal maturation of children in Okinawa. Annals of Human Biology. 1976;3(2):149-55.	TW & Ki-mura	Japan	264	Both	7- 15 years
Kimura K. Skeletal maturity and bone growth in twins. American Journal of Physical Anthropology. 1983;60(4):491-7.	TW2	Japan	179	Both	12- 18 years
Kimura K. Skeletal maturity of the hand and wrist in Japanese children by the TW2 method. Annals of Human Biology. 1977;4(4):353-6.	TW2	Japan	718	Both	0- 19 years
Kimura K. Skeletal maturity of the hand and wrist in Japanese children in Sapporo by the TW2 method. Annals of Human Biology. 1977;4(5):449-53.	TW2	Japan	258	Both	6- 18 years
Kumar V, Venkataraghavan K, Krishnan R, Patil K, Munoli K, Karthik S. The relationship between dental age, bone age and chronological age in underweight children. Journal of pharmacy and bio-allied sciences. 2013;5:S73-9.	Demirjian, Bjork, Grave & Brown	India	100	Both	8- 14 years
Lai EH, Chang JZ, Jane Yao CC, Tsai SJ, Liu JP, Chen YJ, et al. Relationship between age at menarche	NTUH-SMI (hand) +	Taiwan	304	Girls	8- 19 years

and skeletal maturation stages in Taiwanese female orthodontic patients. Journal of the Formosan Medical Association. 2008;107(7):527-32.	CVMS (cervical vertebra)				
Lai EH, Liu JP, Chang JZ, Tsai SJ, Yao CC, Chen MH, et al. Radiographic assessment of skeletal maturation stages for orthodontic patients: hand-wrist bones or cervical vertebrae? Journal of the Formosan Medical Association. 2008;107(4):316-25.	NTUH-SMI (hand) + CVMS (cervical vertebra)	Taiwan	709	Both	8 to 18 years
Lejarraga H, Guimarey L, Orazi V. Skeletal maturity of the hand and wrist of healthy Argentinian children aged 4-12 years, assessed by the TWII method. Annals of Human Biology. 1997;24(3):257-61.	TW2	Argentina	775	Both	4 to 12 years
Lequin MH, van Rijn RR, Robben SG, Hop WC, van Kuijk C. Normal values for tibial quantitative ultrasonometry in caucasian children and adolescents (aged 6 to 19 years). Calcified Tissue International. 2000;67(2):101-5.	Tanner,	Nether-land	596	Both	6 to 20 years
Li CS, Li CY, Xi HJ, Ren F, Huang KQ, Wen YF, et al. Assessment of development of wrist skeletal age of Tibetan adolescent in Lhasa. [Chinese]. Chinese Journal of Clinical Rehabilitation. 2005;9(23):36-8.	Fels	China	1496	Both	7 to 21 years
Li CY, Ren F, Li CS, Huang KQ, Wen YF, Guo ZY, et al. Assessment of skeletal age of hand-wrist in Tibetan adolescent of Lhasa with Fels method. [Chinese]. Chinese Journal of Clinical Rehabilitation. 2006;10(24):36-8.	Fels	China	1496	Both	7 to 21 years
Litsas G, Ari-Demirkaya A. Growth indicators in orthodontic patients. Part 1: comparison of cervical vertebral maturation and hand-wrist skeletal maturation. European Journal of Paediatric Dentistry. 2010;11(4):171-5.	Bjork ,Grave and Brown.	Turkey	393	Both	8 to 18 years
Litsas G, Ari-Demirkaya A. Growth indicators in orthodontic patients. Part 2: comparison of cervical bone age to hand-wrist skeletal age. Relationship with chronological age. European journal of paediatric dentistry : official journal of European Academy of Paediatric Dentistry. 2010;11(4):176-80.	Bjork ,Grave and Brown.	Turkey	393	Both	8 to 18 years
Little BB, Malina RM. Gene-environment interaction in skeletal maturity and body dimensions of urban Oaxaca Mestizo schoolchildren. Annals of Human Biology. 2007;34(2):216-25.	TW2	Mexico, USA	1398	Both	6 to 13 years
Liu B, Wang D, Wang Z. Comparison of skeletal development between rural and urban school-age children. [Chinese]. Zhongguo yi xue ke xue yuan xue bao. 1994;0(3):165-9.	Chinese method	China	1680	Both	7 to 18 years
Liu ZX, Cheng XG, Li XM, Yu W. Bone age of children and adolescent of X-ray film: Comparison of left- and right- hand. [Chinese]. Chinese Journal of Radiology (China). 2013;47(12):1070-3.	CHN left vs right hand	China	1380	Both	1 to 19 years
Maggio A, Flavel A, Hart R, Franklin D. Skeletal age estimation in a contemporary Western Australian population using the Tanner-Whitehouse method. Forensic Science International. 2016;263:e1-8.	TW	Australia	360	Both	0 to 25 years
Magnusson TE. Skeletal maturation of the hand in Iceland. Acta Odontologica Scandinavica. 1979;37(1):21-8.	Björk	Iceland	1426	Both	5 to >17 years
Mahajan S. Evaluation of skeletal maturation by comparing the hand wrist radiograph and cervical vertebrae as seen in lateral cephalogram. Indian Journal of Dental Research. 2011;22(2):309-16.	Fishman & cervical vertebrae	India	100	Both	8 to 18 years
Malina RM, Coelho ESMJ, Figueiredo AJ, Carling C, Beunen GP. Interrelationships among invasive and non-invasive indicators of biological maturation in	Fels	Portugal	180	Boys	11 to 14 years

adolescent male soccer players. <i>Journal of Sports Sciences</i> . 2012;30(15):1705-17.					
Malina RM, Dompier TP, Powell JW, Barron MJ, Moore MT. Validation of a noninvasive maturity estimate relative to skeletal age in youth football players. <i>Clinical Journal of Sport Medicine</i> . 2007;17(5):362-8.	Fels	USA	143	Boys	9 to 14 years
Malina RM, Himes JH, Stepick CD. Skeletal maturity of the hand and wrist in Oaxaca school children. <i>Annals of Human Biology</i> . 1976;3(3):211-9.	TW2	Mexico	394	Both	5 to 18 years
Malina RM, Little BB. Comparison of TW1 and TW2 skeletal age differences in American black and white and in Mexican children 6-13 years of age. <i>Annals of Human Biology</i> . 1981;8(6):543-8.	TW1 +2	USA, Mexico	2562	Both	6 to 13 years
Malina RM, Pena Reyes ME, Eisenmann JC, Horta L, Rodrigues J, Miller R. Height, mass and skeletal maturity of elite Portuguese soccer players aged 11-16 years. <i>Journal of Sports Sciences</i> . 2000;18(9):685-93.	Fels	Portugal	135	Boys	10 to 17 years
Malina RM, Pena Reyes ME, Figueiredo AJ, Coelho ESMJ, Horta L, Miller R, et al. Skeletal age in youth soccer players: implication for age verification. <i>Clinical Journal of Sport Medicine</i> . 2010;20(6):469-74.	Fels	Portugal, Spain	592	Boys	11 to 17 years
Malina RM. Skeletal maturation studied longitudinally over one year in American Whites and Negroes six through thirteen years of age. <i>Human biology; an international record of research</i> . 1970;42(3):377-90.	TW	USA	806	Both	4 to 14 years
Malo L, Lima S, Teixeira V, Canova F, Alves S. Skeletal maturation in a Portuguese population - comparison between the hand-wrist and cervical vertebral maturation. [Portuguese]. <i>Revista Portuguesa de Estomatologia, Medicina Dentaria e Ciurgia Maxilofacial</i> . 2014;55(2):102-9.	Grave & Brown	Portugal	285	Both	7 to 16 years
Mappes MS, Harris EF, Behrents RG. An example of regional variation in the tempos of tooth mineralization and hand-wrist ossification. <i>American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics</i> . 1992;101(2):145-51.	Fishman & Moorrees	USA	585 tann, 295 hand	Both	12 and 13 years
Marak FK, Sangma WBC, Singh MS, Kharrubon B. A roentgenographic study for age estimation in boys and girls of North-Eastern region of India. <i>International Journal of Medical Toxicology and Legal Medicine</i> . 2008;10(2):34-9.	Unclear for wrist, el- bow, knee, pelvis	India	200	Both	16 to 22 years
Marti Henneberg C, Vilardell Latorre E. Bone maturation in the infantile population (boys 8-17 years old and girls 6-16 years old) of Barcelona, Spain. Standard curves from a numeric method. [Spanish]. <i>Medicina Clinica</i> . 1975;64(2):73-8.	TW	Spain	1268	Both	6 to 17 years
Matsuoka H, Sato K, Sugihara S, Murata M. Bone maturation reflects the secular trend in growth. <i>Hormone Research</i> . 1999;52(3):125-30.	TW2	Japan, China	3123	Both	7 to 16 years
Mohammed RB, Kalyan VS, Tircouveluri S, Vigesna GC, Chirla A, Varma DM. The reliability of Fishman method of skeletal maturation for age estimation in children of South Indian population. <i>Journal of Natural Science Biology & Medicine</i> . 2014;5(2):297-302.	Fishman	India	330	Both	9 to 20 years

Mohammed RB, Reddy MA, Jain M, Singh JR, Sanghvi P, Thetay AA. Digital radiographic evaluation of hand-wrist bone maturation and prediction of age in South Indian adolescents. <i>Hand.</i> 2014;9(3):375-83.	Fishman	India	330	Both	8 to 18 years
Molinari L, Hermanussen M. The effect of variability in maturational tempo and midparent height on variability in linear body measurements. <i>Annals of Human Biology.</i> 2005;32(5):679-82.	TW3	Switzerland	232	Both	2 to 20 years
Murata M. Characteristics of pubertal growth in Japanese children from the standpoint of skeletal growth. <i>Acta Paediatrica Japonica.</i> 1992;34(2):236-40; discussion 40-2.	TW2	Japan	200	Both	7 to 18 years
Murata M. Population-specific reference values for bone age. <i>Acta Paediatrica Supplement.</i> 1997;423:113-4.	TW2	Japan	382	Both	1 to 20 years
Pasciuti E, Franchi L, Baccetti T, Milani S, Farronato G. Comparison of three methods to assess individual skeletal maturity. <i>Journal of Orofacial Orthopedics.</i> 2013;74(5):397-408.	Gianni's analysis	Italy	100	Both	6-18 years
Safer AN, Homel P, Chung DD. Lateral comparisons using Fishman's skeletal maturation assessment. <i>Angle Orthodontist.</i> 2015;85(3):408-12.	Fishman	USA	125	Both	8-20 years
Sahin Saglam AM, Gazilerli U. The relationship between dental and skeletal maturity. <i>Journal of Orofacial Orthopedics.</i> 2002;63(6):454-62.	Fishman	Turkey	422	Both	7-15 years
Schmeling A, Baumann U, Schmidt S, Wernecke KD, Reisinger W. Reference data for the Thiemann-Nitz method of assessing skeletal age for the purpose of forensic age estimation. <i>International Journal of Legal Medicine.</i> 2006;120(1):1-4.	Thiemann-Nitz	Germany	402	Both	10-18 years
Schmidt S, Nitz I, Schulz R, Schmeling A. Applicability of the skeletal age determination method of Tanner and Whitehouse for forensic age diagnostics. <i>International Journal of Legal Medicine.</i> 2008;122(4):309-14.	TW	Germany	92	Both	12-16 years
Schmidt S, Nitz I, Schulz R, Tsokos M, Schmeling A. The digital atlas of skeletal maturity by Gilsanz and Ratib: a suitable alternative for age estimation of living individuals in criminal proceedings? <i>International Journal of Legal Medicine.</i> 2009;123(6):489-94.	Gilsanz Ratib	Germany	180	Both	10-18 years
Shim J, Heo G, Lagravere MO. Correlation between three-dimensional morphological changes of the hyoid bone with other skeletal maturation methods in adolescents. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.</i> 2013;116(4):511-7.	Fishman	Canada	62	Both	11-17 years
Shim JJ, Bogowicz P, Heo G, Lagravere MO. Interrelationship and limitations of conventional radiographic assessments of skeletal maturation. <i>International orthodontics / College europeen d'orthodontie.</i> 2012;10(2):135-47.	Fishman	Canada	62	Both	11-17 years
Singer R, Kimura K. Body height, weight, and skeletal maturation in Hottentot (Khoikhoi) children. <i>American Journal of Physical Anthropology.</i> 1981;54(3):401-13.	TW2	South West Africa	120	Both	3-17 years
Soegiharto BM, Cunningham SJ, Moles DR. Skeletal maturation in Indonesian and white children assessed with hand-wrist and cervical vertebrae methods. <i>American Journal of Orthodontics and Dentofacial Orthopedics.</i> 2008;134(2):217-26.	Fishman	Indonesia, USA	2167	Both	8-15 years

Takai S, Akiyoshi T. Skeletal maturity of Japanese children in Western Kyushu. American Journal of Physical Anthropology. 1983;62(2):199-204.	TW2	Japan	985	Both	4 -15 years
Takai S. [Smoothed skeletal maturity curve of Japanese children by Tanner-Whitehouse 2 (TW2) method and its application]. Kaibogaku Zasshi - Journal of Anatomy. 1990;65(6):436-47.	TW2	Japan	6351	Both	6-18 years
Tanner J, Oshman D, Bahhage F, Healy M. Tanner-Whitehouse bone age reference values for North American children.[Erratum appears in J Pediatr. 2012 Dec;161(6):1180]. Journal of Pediatrics. 1997;131(1):34-40.	TW2	USA	450	Both	8-16 Years
Uysal T, Ramoglu SI, Basciftci FA, Sari Z. Chronologic age and skeletal maturation of the cervical vertebrae and hand-wrist: Is there a relationship? American Journal of Orthodontics and Dentofacial Orthopedics. 2006;130(5):622-8.	Björk and Grave and Brown	Turkey	2003	Both	5-24 years
Uysal T, Sari Z, Ramoglu SI, Basciftci FA. Relationships between dental and skeletal maturity in Turkish subjects. Angle Orthodontist. 2004;74(5):657-64.	Björk and Grave and Brown	Turkey	500	Both	7-20 years
Valente-dos-Santos J, Coelho-e-Silva MJ, Duarte J, Pereira J, Rebelo-Goncalves R, Figueiredo A, et al. Allometric multilevel modelling of agility and dribbling speed by skeletal age and playing position in youth soccer players. International Journal of Sports Medicine. 2014;35(9):762-71.	Fels	Portugal	83	Boys	11-15 years
van Lenthe FJ, Kemper HC, van Mechelen W. Skeletal maturation in adolescence: a comparison between the Tanner-Whitehouse II and the Fels method. European Journal of Pediatrics. 1998;157(10):798-801.	Fels, TW2	Nether-land	60	Both	12-16 years
Vignolo M, Naselli A, Magliano P, Di Battista E, Aiardi M, Aiardi G. Use of the new US90 standards for TW-RUS skeletal maturity scores in youths from the Italian population. Hormone Research. 1999;51(4):168-72.	TW-RUS	Italy	1831	Both	8-17 years
Xi HJ, Roche AF. Differences between the hand-wrist and the knee in assigned skeletal ages. American Journal of Physical Anthropology. 1990;83(1):95-102.	Fels	China	4902	Both	2-17 years
Xu YC, Ye LY, Li K, Wang ZY, Ke Y, Zhang QC. Accuracy of age estimation in 14-year-old females by two skeletal age standards. [Chinese]. Fa yi xue za zhi. 2006;22(5):328-9, 32.	CHN	China	51	Girls	14 years
Yang YM, Lee J, Kim YI, Cho BH, Park SB. Axial cervical vertebrae-based multivariate regression model for the estimation of skeletal-maturation status. Orthodontics & Craniofacial Research. 2014;17(3):187-96.	SMI, FMI	Finland	121	Both	6-18 years
Zhang S, Liu L, Hua J, Liu G. The observation of difference between skeletal age and chronological age with hand and wrist in Chinese Han adolescent. [Chinese]. Chinese Journal of Forensic Medicine. 2009;24(1):18-20.	RUS-CHN	China	12414	Both	3-18 years
Zhang S, Liu L, Zhang J, Liu G. The RUS-CHN radiographic atlas method of assessing skeletal age for adolescent. [Chinese]. Chinese Journal of Forensic Medicine. 2009;24(4):249-53.	RUS-CHN	China	5472	Both	13-18 years
Zhang S, Ma Z, Han Y, Shen X, Liu G, Xu R. The effects of the differences of skeletal development on age estimation in children and adolescent be-	RUS-CHN	China	11635	Both	3-18 years

tween large and middling cities in China. [Chinese]. Chinese Journal of Forensic Medicine. 2009;24(2):99-102+3.					
Zhang SY, Liu LJ, Han YS, Liu G, Ma ZG, Shen XZ, et al. [Reference values of differences between TW3-C RUS and TW3-C Carpal bone ages of children from five cities of China]. Zhonghua Erke Zazhi. 2008;46(11):851-5.	TW3	China	9408	Both	1-14 years
Zhang SY, Liu LJ, Wu ZL, Liu G, Ma ZG, Shen XZ, et al. Standards of TW3 skeletal maturity for Chinese children. Annals of Human Biology. 2008;35(3):349-54.	TW3	China	17 401	Both	2-20 years
Zhao XD, Hu Y, Pan SN, Li XF, Li XM, Cheng XG, et al. Evaluation of bone age of wrist-hand in adolescent of Shenyang and Beijing. [Chinese]. Chinese Journal of Radiology (China). 2013;47(12):1066-9.	CHN	China	1333	Both	4-18 years

Appendix 5: Studies excluded after full text assessment

Reference	Reason for exclusion
Avenarius DMF, Ording Muller LS, Eldevik P, Owens CM, Rosendahl K. The paediatric wrist revisited-findings of bony depressions in healthy children on radiographs compared to MRI. <i>Pediatric Radiology.</i> 2012;42(7):791-8.	Less than 50 subjects in relevant age range.
A. Erratum: Tanner-whitehouse bone age reference values for North American children (Journal of Pediatrics (1997) 131 (34-40)). <i>Journal of Pediatrics.</i> 2012;161(6):1180.	Not an empirical study published in full text format (abstracts, reviews, other)
A. EVALUATION of skeletal age in roentgenograms. <i>Nutrition Reviews.</i> 1958;16(4):101-2.	Not an empirical study published in full text format (abstracts, reviews, other)
Abdel-Kader HM. The potential of digital dental radiography in recording the adductor sesamoid and the MP3 stages. <i>British Journal of Orthodontics.</i> 1999;26(4):291-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Acheson RM, Fowler G, Fry EI, Janes M, Koski K, Urbano P, et al. Studies in the Reliability of Assessing Skeletal Maturity from X-Rays. I. Greulich-Pyle Atlas. Human biology; an international record of research. 1963;35:317-49.	Less than 50 subjects in relevant age range.
Acheson RM, Vicinus JH, Fowler GB. Studies in the reliability of assessing skeletal maturity from x-rays. 3. Greulich-Pyle Atlas and Tanner-Whitehouse method contrasted. <i>Human Biology.</i> 1966;38(3):204-18.	Less than 50 subjects in relevant age range.
Acheson RM, Vicinus JH, Fowler GB. Studies in the Reliability of Assessing Skeletal Maturity from X-Rays. II. The Bone-Specific Approach. <i>Human Biology.</i> 1964;36:211-28.	Less than 50 subjects in relevant age range.
Aggarwal ML, Pathak IC. Roentgenologic study of epiphyseal union in Punjabi girls for determination of age. <i>Indian Journal of Medical Research.</i> 1957;45(2):283-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Al Khal HA, Wong RW, Rabie AB. Elimination of hand-wrist radiographs for maturity assessment in children needing orthodontic therapy. <i>Skeletal Radiology.</i> 2008;37(3):195-200.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Andersen E. Skeletal development as a measure of biological development. <i>Acta Paediatrica Scandinavica.</i> 1967;0:Suppl 177:73-4.	Not an empirical study published in full text format (abstracts, reviews, other).
Anderson M. Use of the Greulich-Pyle "Atlas of Skeletal Development of the Hand and Wrist" in a clinical context. <i>American Journal of Physical Anthropology.</i> 1971;35(3):347-52.	Not an empirical study published in full text format (abstracts, reviews, other).
Armengaud G. Radiological examination of the ossification of the hand in Spanish children. <i>Electromedica.</i> 1973;41(4):188-91.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ashizawa K. Interpretation errors in the estimation of bone maturation stages by an observer using the Tanner Whitehouse method. [French]. <i>Bulletins et Memoires de la Societe d'Anthropologie de Paris.</i> 1974;13(3):335-50.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Bagherpour A, Pousti M, Adelianfar E. Hand skeletal maturity and its correlation with mandibular dental development. <i>Journal of Clinical & Experimental Dentistry.</i> 2014;6(3):e275-9.	Less than 50 subjects in relevant age range.
Bai WJ, Ning G, Qu HB, Sun XM, Xiang CF, Wu KM, et al. Comparison among three standards of TW2 to skeletal age determination in children with central precocious puberty. [Chinese]. <i>Journal of Forensic Medicine.</i> 2010;26(3):181-4.	A study population with chronic disease or developmental disorders.
Banik ND, Nayar S, Krishna R, Raj L, Gadekar NG. Skeletal maturation of Indian children. <i>Indian Journal of Pediatrics.</i> 1970;37(269):249-54.	The study population is not living persons between the ages 10-25 years
Bassett RB, Briggs C, Drummer OH. Age estimation using CT imaging of the third molar tooth, the medial clavicular epiphysis, and the spheno-occipital synchondrosis: a multifactorial approach. <i>Forensic Science International.</i> 2011;212(1):273.e1-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Bassed RB, Briggs C, Drummer OH. The incidence of asymmetrical left/right skeletal and dental development in an Australian population and the effect of this on forensic age estimations. International Journal of Legal Medicine. 2012;126(2):251-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Baumann U, Schulz R, Reisinger W, Heinecke A, Schmeling A, Schmidt S. Reference study on the time frame for ossification of the distal radius and ulnar epiphyses on the hand radiograph. Forensic Science International. 2009;191(1):15-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Beek FJA. Current validation of the Greulich and Pyle atlas for the determination of skeletal age. [Dutch]. Nederlands Tijdschrift voor Geneeskunde. 2003;147(15):689-90.	Not an empirical study published in full text format (abstracts, reviews, other).
Belkin V, Livshits G, Otremski I, Kobylansky E. Aging bone score and climatic factors. American Journal of Physical Anthropology. 1998;106(3):349-59.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Benson J, Williams J. Age determination in refugee children. Australian Family Physician. 2008;37(10):821-5.	Not an empirical study published in full text format (abstracts, reviews, other).
Berdikulov GB. [X-ray data on the times of synostosis of the bones in the hand and distal portion of the forearm in girls and young women of Uzbek nationality from 11 to 19 years of age]. Sudebno-Meditsinskaia Ekspertiza. 1980;23(2):23-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Beunen G, Cameron N. The reproducibility of TW2 skeletal age assessments by a self-taught assessor. Annals of Human Biology. 1980;7(2):155-62.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Beunen G, Malina RM, Claessens AL, Lefevre J, Thomis M. Ulnar variance and skeletal maturity of radius and ulna in female gymnasts. Medicine & Science in Sports & Exercise. 1999;31(5):653-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Beunen G, Ostyn M, Renson R, Simons J, Swalus P, van Gerven D, et al. Skeletal age and physical development of 12-year-old boys. [Dutch]. Archives belges de medecine sociale, hygiene, medecine du travail et medecine legale. 1972;0(2):102-19.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Beunen G, Ostyn M, Renson R, Simons J, VanGerven D. Patterns of TW-1 and TW-2 skeletal age differences in 12-19-year-old Belgian boys. Annals of Human Biology. 1983;10(5):479-82.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Biggerstaff RH. Forensic dentistry and the assessment of skeletal age using hand-wrist film standards. Dental Clinics of North America. 1977;21(1):159-66.	Not an empirical study published in full text format (abstracts, reviews, other).
Birch D, MacKenzie RG, Iverson EF. Age assessment of young people: Preliminary findings of a blinded study using a multifactorial developmental approach. Journal of Adolescent Health. 2013;1:S3-S4.	Not an empirical study published in full text format (abstracts, reviews, other).
Birkbeck JA, Lee M. Growth and skeletal maturation in British Columbia Indian Populations. American Journal of Physical Anthropology. 1973;38(3):727-38.	Less than 50 subjects in relevant age range.
Boechat MI, Lee DC. Graphic representation of skeletal maturity determinations. AJR American Journal of Roentgenology. 2007;189(4):873-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Bolboaca S, Jantschi L, Denes C, Cadariu AA. Skeletal maturity assessment client-server application. Rentgenologiya i Radiologiya. 2005;44(3):189-93.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Borms J, Broekhoff J, Beunen G. [Comparative studies of 3 methods of skeletal age determination]. Archives Belges de Medecine Sociale, Hygiene, Medecine du Travail et Medecine Legale - Belgisch Archief van Sociale Geneeskunde, Hygiene, Arbeidsgeneeskunde en Gerechtelijke Geneeskunde. 1973;31(5):343-53.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Borms J, Broekhoff J, Beunen G. Comparative study of three methods for the assessment of bone age. [Dutch]. Archives Belges de Medecine Sociale Hygiene Medecine du Travail et Medecine Legale. 1973;31(5):343-53.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Bowden BD. Epiphyseal changes in the hand/wrist area as indicators of adolescent stage. Australian Orthodontic Journal. 1976;4(3):87-104.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Bruns HA. [Roentgen diagnosis of skeletal age]. Monatsschrift fur Kinderheilkunde. 1969;117(9):582-3.	Not an empirical study published in full text format (abstracts, reviews, other).

Buckler JM. Comparison of systems of estimating skeletal age. <i>Archives of Disease in Childhood</i> . 1977;52(8):667-8.	Not an empirical study published in full text format (abstracts, reviews, other).
Buckler JM. Skeletal age changes in puberty. <i>Archives of Disease in Childhood</i> . 1984;59(2):115-9.	Less than 50 subjects in relevant age range.
Bugyi B, Kausz I. Radiographic determination of the skeletal age of the young swimmers. <i>The Journal of sports medicine and physical fitness</i> . 1970;10(4):269-70.	Not an empirical study published in full text format (abstracts, reviews, other).
Bull RK, Edwards PD, Kemp PM, Fry S, Hughes IA. Bone age assessment: a large scale comparison of the Greulich and Pyle, and Tanner and Whitehouse (TW2) methods. <i>Archives of Disease in Childhood</i> . 1999;81(2):172-3.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Buric M, Momcilovic B. Growth pattern and skeletal age in school girls with idiopathic scoliosis. <i>Clinical Orthopaedics & Related Research</i> . 1982;0(170):238-42.	A study population with chronic disease or developmental disorders.
Burov SA, Reznikov BD. [Ossification characteristics of the wrist and distal part of the forearm and their significance in the determination of age]. <i>Sudebno-Meditsinskaia Ekspertiza</i> . 1972;15(1):21-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Burov SA. [Ossification times for human extremities]. <i>Sudebno-Meditsinskaia Ekspertiza</i> . 1973;16(3):11-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Byun BR, Kim YI, Yamaguchi T, Maki K, Ko CC, Hwang DS, et al. Quantitative skeletal maturation estimation using cone-beam computed tomography-generated cervical vertebral images: a pilot study in 5- to 18-year-old Japanese children. <i>Clinical Oral Investigations</i> . 2015;19(8):2133-40.	Less than 50 subjects in relevant age range.
Cabello AS, Estrada RC. Automated system of bone maturity evaluation applied to the Tanner-Whitehouse 2 method. [Spanish]. <i>Anales Espanoles de Pediatría</i> . 1999;50(2):134-9.	Less than 50 subjects in relevant age range.
Cameriere R, De Luca S, Cingolani M, Ferrante L. Measurements of developing teeth, and carpal and epiphyses of the ulna and radius for assessing new cut-offs at the age thresholds of 10, 11, 12, 13 and 14 years. <i>Journal of Forensic & Legal Medicine</i> . 2015;34:50-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Cameron N, Mirwald RL, Bailey DA, Davies PS. The application of new height-prediction equations (Tanner-Whitehouse mark 2) to a sample of Canadian boys. <i>Annals of Human Biology</i> . 1985;12(3):233-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Canavese F, Charles YP, Dimeglio A, Schuller S, Rousset M, Samba A, et al. A comparison of the simplified olecranon and digital methods of assessment of skeletal maturity during the pubertal growth spurt. <i>Bone and Joint Journal</i> . 2014;96(11):1556-60.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Cao F, Huang HK, Pietka E, Gilsanz V. Digital hand atlas and web-based bone age assessment: system design and implementation. <i>Computerized Medical Imaging & Graphics</i> . 2000;24(5):297-307.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Capron JP, Sempe M, Giard MH, Pernier J, Rubel P. [Calculating system for the degree of skeletal maturation in the child and adolescent]. <i>Medical & Biological Engineering & Computing</i> . 1980;18(1):70-2.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Cardoso HF. Age estimation of adolescent and young adult male and female skeletons II, epiphyseal union at the upper limb and scapular girdle in a modern Portuguese skeletal sample. <i>American Journal of Physical Anthropology</i> . 2008;137(1):97-105.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Castriota-Scanderbeg A, Sacco MC, Emberti-Gialloreto L, Fraracci L. Skeletal age assessment in children and young adults: comparison between a newly developed sonographic method and conventional methods. <i>Skeletal Radiology</i> . 1998;27(5):271-7.	A study population with chronic disease or developmental disorders.
Chai HY, Swee TT, Seng GH, Wee LK. Multipurpose contrast enhancement on epiphyseal plates and ossification centers for bone age assessment. <i>Biomedical Engineering Online</i> . 2013;12:27.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Chai HY, Wee LK, Swee TT, Salleh Sh H, Chea LY. An artifacts removal post-processing for epiphyseal region-of-interest (EROI) localization in automated bone age assessment (BAA). <i>Biomedical Engineering Online</i> . 2011;10:87.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Chaitanya NCSK, Ali IM, Shashikanth MC. Age estimation using cervical vertebrae. Indian Journal of Forensic Medicine and Toxicology. 2009;3(1):11-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Charles YP, Canavese F, Schuller S, Steib JP, Rousset M, Dimeglio A. Comparison of simplified olecranon and digital methods of skeletal maturity assessment. European Spine Journal. 2014;23:S497-S8.	Not an empirical study published in full text format (abstracts, reviews, other).
Charlet G. [Dental age, skeletal age]. Revue d Odonto-Stomatologie. 1984;13(1):19-31.	Not an empirical study published in full text format (abstracts, reviews, other).
Chaumoitre K, Adalian P, Colavolpe N, Ramis O, Marciano S, Leonetti G, et al. [Value of the sesamoid bone of the thumb in the determination of bone age]. Journal de Radiologie. 2008;89(12):1921-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Cheng XG, Zeng JJ, Yu W. The status of wrist bone age assessment in China and research perspective. [Chinese]. Chinese Journal of Radiology (China). 2013;47(12):1061-2.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Cockill T, Hewitt A, Wright N, Elder C. Basic: Bone age study in children. Hormone Research in Paediatrics. 2015;84:270.	Not an empirical study published in full text format (abstracts, reviews, other).
Cole TJ. The evidential value of developmental age imaging for assessing age of majority. Annals of Human Biology. 2015;42(4):379-88.	Not an empirical study published in full text format (abstracts, reviews, other).
Conceicao EL, Cardoso HF. Environmental effects on skeletal versus dental development II: further testing of a basic assumption in human osteological research. American Journal of Physical Anthropology. 2011;144(3):463-70.	3, Less than 50 subjects in relevant age range.
Costacurta M, Sicuro L, Di Renzo L, Condo R, De Lorenzo A, Docimo R. Childhood obesity and skeletal-dental maturity. European journal of paediatric dentistry : official journal of European Academy of Paediatric Dentistry. 2012;13(2):128-32.	Less than 50 subjects in relevant age range.
Danaei SM, Karamifar A, Sardarian A, Shahidi S, Karamifar H, Ali-pour A, et al. Measuring agreement between cervical vertebrae and hand-wrist maturation in determining skeletal age: reassessing the theory in patients with short stature. American Journal of Orthodontics & Dentofacial Orthopedics. 2014;146(3):294-8.	A study population with chronic disease or developmental disorders.
Darmawan MF, Yusuf SM, Abdul Kadir MR, Haron H. Age estimation based on bone length using 12 regression models of left hand X-ray images for Asian children below 19 years old. Legal Medicine. 2015;17(2):71-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Das Gupta SM, Prasad V, Singh S. A roentgenologic study of epiphyseal union around elbow, wrist and knee joints and the pelvis in boys and girls of Uttar Pradesh. Journal of the Indian Medical Association. 1974;62(1):10-2.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Davis LM, Theobald BJ, Lines J, Toms A, Bagnall A. On the segmentation and classification of hand radiographs. International Journal of Neural Systems. 2012;22(5):1250020.	Not an empirical study published in full text format (abstracts, reviews, other).
Dembetembe KA. Greulich-Pyle method and determining skeletal maturation in African males. Clinical Anatomy. 2012;25:920.	Not an empirical study published in full text format (abstracts, reviews, other).
Demirjian A, Buschang PH, Tanguay R, Patterson DK. Interrelationships among measures of somatic, skeletal, dental, and sexual maturity. American Journal of Orthodontics. 1985;88(5):433-8.	Less than 50 subjects in relevant age range.
Deschamps JP, Benchemsi N. Bone maturation and socioeconomic and cultural conditions (Cross sectional study of 4,526 children and adolescents). [French]. Revue de Pediatrie. 1974;10(3):143-51.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Dhamija SC, Agarwal KN, Gupta SK, Katiyar GP. Evaluation of hand and foot ossification centres for assessment of bone age. Indian Pediatrics. 1976;13(3):201-8.	The study population is not living persons between the ages 10-25 years
Di Lorenzo P, Niola M, Pantaleo G, Buccelli C, Amato M. On the comparison of age determination methods based on dental development radiographic studies in a sample of Italian population. Dental Cadmos. 2015;83(1):38-45.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Dickhaus H, Wastl S. Computer assisted bone age assessment. Medinfo. 1995;8:709-13.	Not an empirical study published in full text format (abstracts, reviews, other).

Dimeglio A, Charles YP, Daures JP, De Rosa V, Kabore B. Accuracy of the Sauvegrain method in determining skeletal age during puberty. <i>Journal of Bone and Joint Surgery - Series A</i> . 2005;87(8):1689-96.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Dogaroiu C, Capatina CO, Gherghe EV, Avramou M. The importance of the ossification centre morphology in the left hand-wrist bones for age evaluation. <i>Romanian Journal of Legal Medicine</i> . 2014;22(2):105-8.	Less than 50 subjects in relevant age range.
Dopper T. [Development of the skeleton in child. Prerequisites for the assessment of bone age and skeletal maturity]. <i>Medizinische Welt</i> . 1976;27(4):142-5.	Not an empirical study published in full text format (abstracts, reviews, other).
Drayer NM, Cox LA. Assessment of bone ages by the Tanner-Whitehouse method using a computer-aided system. <i>Acta Paediatrica Supplement</i> . 1994;406:77-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Dreizen S, Spirakis CN, Stone RE. A comparison of skeletal growth and maturation in undernourished and well-nourished girls before and after menarche. <i>Journal of Pediatrics</i> . 1967;70(2):256-63.	Less than 50 subjects in relevant age range.
Dundaroz MR, Sarici SU, Denli M, Aydin HI, Kocaoglu M, Ozisik T. Bone age in children with nocturnal enuresis. <i>International Urology & Nephrology</i> . 2001;32(3):389-91.	Less than 50 subjects in relevant age range.
Duren D, Nahhas R, Sherwood R, Lee M, Choh A, Towne B, et al. Secular trend for earlier skeletal maturation in us children. <i>Journal of Bone and Mineral Research</i> . 2010;25:S54.	Not an empirical study published in full text format (abstracts, reviews, other).
Duren DL, Nahhas RW, Sherwood RJ. Do Secular Trends in Skeletal Maturity Occur Equally in Both Sexes? <i>Clinical Orthopaedics & Related Research</i> . 2015;473(8):2559-67.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Durka-Zajac M, Marcinkowska A, Mitus-Kenig M. Bone age assessment using cephalometric photographs. <i>Polish Journal of Radiology</i> . 2013;78(2):19-25.	Less than 50 subjects in relevant age range.
Dvorak J, George J, Junge A, Hodler J. Age determination by magnetic resonance imaging of the wrist in adolescent male football players. <i>British Journal of Sports Medicine</i> . 2007;41(1):45-52.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ebri Torne B, Ebri Verde I. [Metacarpophalangeal and carpal numeric indices to calculate bone age and predict adult size]. <i>Anales de Pediatría</i> . 2012;76(4):199-213.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ebri Torne B. [New method for evaluating ossification of the carpal bone. From a study with 5225 Spanish children]. <i>Pediatrie</i> . 1993;48(11):813-7.	Not an empirical study published in full text format (abstracts, reviews, other).
Eklof O, Ringertz H. A method for assessment of skeletal maturity. <i>Annales de Radiologie</i> . 1967;10(3):330-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Espina-Fereira A, Ortega Al, Barrios FA, Maldonado YJ, Fereira JL. [Metric and angular variables of the mandibular ramus on panoramic radiographs, as indicators for chronologic age]. <i>Investigacion Clinica</i> . 2007;48(4):403-18.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Faruch Bilfeld M, Dedouit F, Soumah M, Cartegnie S, Joffre F, Rousseau H, et al. [Value of radiographic evaluation of the second metacarpal in the determination of bone age]. <i>Journal de Radiologie</i> . 2008;89(12):1930-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Fazekas A, Farkas G, Kocsis SG. [Bone age of children in 2 communities in Hungary with different fluoride levels in the drinking water]. <i>Anthropologischer Anzeiger</i> . 1984;42(1):57-65.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ferrante L, Skrami E, Gesuita R, Cameriere R. Bayesian calibration for forensic age estimation. <i>Statistics in Medicine</i> . 2015;34(10):1779-90.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Fieuws S, Willems G, Larsen-Tangmose S, Lynnerup N, Boldsen J, Thevissen P. Obtaining appropriate interval estimates for age when multiple indicators are used: evaluation of an ad-hoc procedure. <i>International Journal of Legal Medicine</i> . 2016;130(2):489-99.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Figueiredo AJ, Coelho ESMJ, Cumming SP, Malina RM. Size and maturity mismatch in youth soccer players 11- to 14-years-old. <i>Pediatric Exercise Science</i> . 2010;22(4):596-612.	Less than 50 subjects in relevant age range.

Fischer B, Welter P, Gunther RW, Deserno TM. Web-based bone age assessment by content-based image retrieval for case-based reasoning. International Journal of Computer Assisted Radiology and Surgery. 2012;7(3):389-99.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Fishman LS. Can cephalometric x-rays of the cervical column be used instead of hand-wrist x-rays to determine patient's maturational age? American Journal of Orthodontics & Dentofacial Orthopedics. 2002;122(1):18A-9A.	Not an empirical study published in full text format (abstracts, reviews, other).
Fleischer-Peters A, Baykan C, Prestele H. Determination of the duration of the stages of maturation of the hand. [German]. Fortschritte der Kieferorthopadie. 1978;39(1):77-86.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Flor-Cisneros A, Roemmich JN, Rogol AD, Baron J. Bone age and onset of puberty in normal boys. Molecular & Cellular Endocrinology. 2006;254:202-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Forbes AP, Ronaghya HA, Majd M. Skeletal maturation of children in Shiraz, Iran. American Journal of Physical Anthropology. 1971;35(3):449-54.	The study population is not living persons between the ages 10-25 years
Forsyth DM. Estimation of young peoples' ages by a radiological method. Transactions of the Royal Society of Tropical Medicine & Hygiene. 1970;64(1):196-7.	Not an empirical study published in full text format (abstracts, reviews, other).
Friedrich RE, v Maydell LA, Ulbricht C, Scheuer HA. [Prosthetic restorations, dental root fillings and periodontal bone resorptions as a forensic-odontologic aid for determining the age above 18 years: A radiographic study of orthopantomograms from a group of teenagers and young adults]. Archiv fur Kriminologie. 2005;216(5):166-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Fry EI. Assessing skeletal maturity: comparison of the atlas and individual bone technics. Nature. 1968;220(5166):496-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Gaind BN, Ghosh S, Sarin GS, Satya Nath S. Skeletal growth in healthy children from a public school in Delhi. Indian Journal of Medical Research. 1980;72:527-36.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Gandini P, Mancini M, Andreani F. A comparison of hand-wrist bone and cervical vertebral analyses in measuring skeletal maturation. Angle Orthodontist. 2006;76(6):984-9.	Less than 50 subjects in relevant age range.
Gandini P, Rizzo S, Renzi P, Sfondrini G. Dental age and skeletal age: correlation study. [Italian]. Mondo ortodontico. 1989;14(2):207-10.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Gefferth K. [A method for the determination of biological bone age]. Acta Paediatrica Academiae Scientiarum Hungaricae. 1970;11(1):59-66.	3
George J, Nagendran J, Azmi K. Comparison study of growth plate fusion using MRI versus plain radiographs as used in age determination for exclusion of overaged football players. British Journal of Sports Medicine. 2012;46(4):273-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Gertych A, Zhang A, Sayre J, Pospiech-Kurkowska S, Huang HK. Bone age assessment of children using a digital hand atlas. Computerized Medical Imaging & Graphics. 2007;31(4):322-31.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ghosh S, Bhardawaj OP, Varma KP. A study of skeletal maturation of hand and wrist and its relationship to nutrition. Indian Pediatrics. 1966;3(4):145-52.	The study population is not living persons between the ages 10-25 years
Giua MR, Pasini M, Tecco S, Marchetti E, Giannotti L, Marzo G. Skeletal maturation in obese patients. American Journal of Orthodontics and Dentofacial Orthopedics. 2012;142(6):774-9.	Less than 50 subjects in relevant age range.
Graham CB. Assessment of bone maturation--methods and pitfalls. Radiologic Clinics of North America. 1972;10(2):185-202.	Not an empirical study published in full text format (abstracts, reviews, other).
Graham EA. Economic, racial, and cultural influences on the growth and maturation of children. Pediatrics in Review. 2005;26(8):290-4.	Not an empirical study published in full text format (abstracts, reviews, other).
Grau U, Klaus F. [Skeletal age and school achievements. Study on the prognostic value of the skeletal age as a measure for physical school readiness]. Praxis der Kinderpsychologie und Kinderpsychiatrie. 1975;24(4):146-55.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Grave K, Townsend G. Hand-wrist and cervical vertebral maturation indicators: how can these events be used to time Class II treatments? <i>Australian Orthodontic Journal</i> . 2003;19(2):33-45.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Grave KC, Brown T. Skeletal ossification and the adolescent growth spurt. <i>American Journal of Orthodontics</i> . 1976;69(6):611-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Green LJ, Sedita J. Genetic variability of skeletal maturity in human twins. <i>Journal of Dental Research</i> . 1971;50(6):1501.	Not an empirical study published in full text format (abstracts, reviews, other).
Green WT, Anderson M. Skeletal age and the control of bone growth. <i>Instructional Course Lectures</i> . 1960;17:199-217.	Not an empirical study published in full text format (abstracts, reviews, other).
Gresham H, Brown T, Barrett MJ. Skeletal and denture patterns in children from Yuendumu, Central Australia, and Melbourne. <i>Australian Dental Journal</i> . 1965;10(6):462-8.	Less than 50 subjects in relevant age range.
Grippaudo C, Garcovich D, Volpe G, Lajolo C. Comparative evaluation between cervical vertebral morphology and hand-wrist morphology for skeletal maturation assessment. <i>Minerva Stomatologica</i> . 2006;55(5):271-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Gron AM, Medicus H. Reproducibility of rating bone development (Tanner-Whitehouse system). <i>American Journal of Orthodontics</i> . 1971;60(2):201-2.	Not an empirical study published in full text format (abstracts, reviews, other).
Gupta M, Divyashree R, Abhilash P, MN AB, Murali K. Correlation between Chronological Age, Dental Age and Skeletal Age among Monozygotic and Dizygotic Twins. <i>Journal of International Oral Health</i> . 2013;5(1):16-22.	Less than 50 subjects in relevant age range.
Gupta S, Chada MK, Sharma A. Assessment of puberty growth spurt in boys and girls--a dental radiographic method. <i>Journal of the Indian Society of Pedodontics & Preventive Dentistry</i> . 1995;13(1):4-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Haavikko K. Skeletal age estimated in a few selected ossification centres of the hand wrist. A simple method for clinical use. <i>Proceedings of the Finnish Dental Society</i> . 1974;70(1):7-14.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Hackman L, Black S. Does mirror imaging a radiograph affect reliability of age assessment using the Greulich and Pyle atlas? <i>Journal of Forensic Sciences</i> . 2012;57(5):1276-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Harmsen M, Fischer B, Schramm H, Seidl T, Deserno TM. Support vector machine classification based on correlation prototypes applied to bone age assessment. <i>IEEE journal of biomedical and health informatics</i> . 2013;17(1):190-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Harris EF, Weinstein S, Weinstein L, Poole AE. Predicting adult stature: a comparison of methodologies. <i>Annals of Human Biology</i> . 1980;7(3):225-34.	Another objective than to compare age estimation using x-ray of the hand with known chronological age. & Less than 50 subjects in relevant age range.
Harrison MH, Turner MH, Jacobs P. Skeletal immaturity in Perthes' disease. <i>Journal of Bone & Joint Surgery - British Volume</i> . 1976;58(1):37-40.	Less than 50 subjects in relevant age range.
Hattner R, Frost HM. Mean skeletal age: its calculation, and theoretical effects on skeletal tracer physiology and on the physical characteristics of bone. <i>Henry Ford Hospital Medical Bulletin</i> . 1963;11:201-16.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Hawley N, Rousham E, Norris S, Pettifor J, Cameron N. Determinants of skeletal age deviation in South African children. <i>American Journal of Physical Anthropology</i> . 2010;141:125.	Not an empirical study published in full text format (abstracts, reviews, other).
Hawley NL, Rousham EK, Johnson W, Norris SA, Pettifor JM, Cameron N. Determinants of relative skeletal maturity in South African children. <i>Bone</i> . 2012;50(1):259-64.	The study population is not living persons between the ages 10-25 years
Healy MJ. Assessing physical maturity by scores for change in skeletal appearance. <i>Journal of Anatomy</i> . 1970;107:180.	Not an empirical study published in full text format (abstracts, reviews, other).
Hegde DY, Baliga S, Yeluri R, Munshi AK. Digital radiograph of the middle phalanx of the third finger (MP3) region as a tool for skeletal maturity assessment. <i>Indian Journal of Dental Research</i> . 2012;23(4):447-53.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Heikkinen E, Kiiskinen A, Kayhty B, Rimpela M, Vuori I. Assessment of biological age. Methodological study in two finnish populations. <i>Gerontologia</i> . 1974;20(1):33-43.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Heimendinger J. [Results of a Survey of Body Measurements of 5000 Basel Children 2-18 Years of Age]. <i>Helvetica Paediatrica Acta</i> . 1964;19:SUPPL 13:1-131.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Helm S, Siersbaek-Nielsen S, Skieller V, Bjork A. [Relationship between skeletal maturation of the hand and pubertal spurt in stature growth]. <i>Mondo Ortodontico</i> . 1977;19(6):64-74.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Henderson AS. The physiological maturity of adolescent psychiatric patients, juvenile delinquents and normal teenagers. <i>British Journal of Psychiatry</i> . 1969;115(525):895-905.	Another objective than to compare age estimation using x-ray of the hand with known chronological age. & Less than 50 subjects in relevant age range.
Heyworth BE, Osei DA, Fabricant PD, Schneider R, Doyle SM, Green DW, et al. The shorthand bone age assessment: A simpler alternative to current methods. <i>Journal of Pediatric Orthopaedics</i> . 2013;33(5):569-74.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Himes JH. An early hand-wrist atlas and its implications for secular change in bone age. <i>Annals of Human Biology</i> . 1984;11(1):71-5.	Less than 50 subjects in relevant age range.
Hinz R. [Radiography of the hand]. <i>Zahnärztliche Mitteilungen</i> . 1979;69(22):1387-9.	Not an empirical study published in full text format (abstracts, reviews, other).
Hirth L. [Estimate of age in a living person]. <i>Deutsche Zeitschrift für die Gesamte Gerichtliche Medizin</i> . 1959;48(2):188-94.	Not an empirical study published in full text format (abstracts, reviews, other).
Holderbaum RM, Veeck EB, Oliveira HW, Silva CL, Fernandes A. Comparison among dental, skeletal and chronological development in HIV-positive children: a radiographic study. <i>Pesquisa Odontologica Brasileira = Brazilian Oral Research</i> . 2005;19(3):209-15.	Less than 50 subjects in relevant age range.
Horak J, Kovarikova I, Vejvalka J, Kucerova K, Dvorakova M. Evaluating bone maturation of present Czech children population. [Czech]. <i>Ceska Radiologie</i> . 2006;60(6):424-33.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Horter MJ, Friesen S, Wacker S, Vogt B, Leidiger B, Roedl R, et al. [Determination of skeletal age : comparison of the methods of Greulich and Pyle and Tanner and Whitehouse]. [German]. <i>Der Orthopade</i> . 2012;41(12):966-76.	Less than 50 subjects in relevant age range.
Hsieh CW, Jong TL, Chou YH, Tiu CM. Computerized geometric features of carpal bone for bone age estimation. <i>Chinese Medical Journal</i> . 2007;120(9):767-70.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Hsieh CW, Liu TC, Jong TL, Tiu CM. Long-term secular trend of skeletal maturation of Taiwanese children between agricultural (1960s) and contemporary (after 2000s) generations using the Tanner-Whitehouse 3 (TW3) method. <i>Journal of Pediatric Endocrinology and Metabolism</i> . 2013;26(3):231-7.	The study population is not living persons between the ages 10-25 years
Hunt EE, Jr., Gleiser I. The estimation of age and sex of preadolescent children from bones and teeth. <i>American Journal of Physical Anthropology</i> . 1955;13(3):479-87.	Less than 50 subjects in relevant age range.
Hunter WS, Baumrind S, Popovich F, Jorgensen G. Forecasting the timing of peak mandibular growth in males by using skeletal age. <i>American Journal of Orthodontics and Dentofacial Orthopedics</i> . 2007;131(3):327-33.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Isiekwe M. Skeletal maturity of children in Lagos, Nigeria. <i>Odonto-stomatologie tropicale = Tropical dental journal</i> . 1987;10(2):111-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Jelliffe DB. Age assessment in field surveys of children of the tropics. <i>Journal of Pediatrics</i> . 1966;69(5):826-8.	Not an empirical study published in full text format (abstracts, reviews, other).
Johnson GF, Dorst JP, Kuhn JP, Roche AF, Davila GH. Reliability of skeletal age assessments. <i>American Journal of Roentgenology, Radium Therapy & Nuclear Medicine</i> . 1973;118(2):320-7.	Less than 50 subjects in relevant age range.
Johnston FE, Jahina SB. The contribution of the carpal bones to the assessment of skeletal age. <i>American Journal of Physical Anthropology</i> . 1965;23(4):349-54.	Less than 50 subjects in relevant age range.
Johnston FE. The Relationship of Certain Growth Variables to Chronological and Skeletal Age. <i>Human Biology</i> . 1964;36:16-27.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kaawach W, Ecklund K, Di Canzio J, Zurakowski D, Waters PM. Normal ranges of scapholunate distance in children 6 to 14 years old. <i>Journal of Pediatric Orthopedics</i> . 2001;21(4):464-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Kadam SS, Belagatti SL, Kulkarni VG. A study of epiphyseal union of base of first metacarpal bone radiologically for estimation of age. <i>Medico-Legal Update</i> . 2012;12(1):78-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kaipova ZA, Giyasov ZA, Adylov OA. Age determination by skeletal bones using X-ray method of investigation. [Russian]. <i>Sudebno-Meditsinskaya Ekspertisa</i> . 1991;34(3):21-3.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kalichman L, Cohen Z, Kobylansky E, Livshits G. Interrelationship between bone aging traits and basic anthropometric characteristics. <i>American Journal of Human Biology</i> . 2002;14(3):380-90.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kalichman L, Malkin I, Livshits G, Pavlovsky O, Batsevich V, Kobylansky E. Variation of skeletal biomarkers of biological aging in a Chuvashian population: a longitudinal study. <i>American Journal of Human Biology</i> . 2007;19(1):74-81.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kamal M, R, Goyal S. Comparative evaluation of hand wrist radiographs with cervical vertebrae for skeletal maturation in 10-12 years old children. <i>Journal of Indian Society of Pedodontics and Preventive Dentistry</i> . 2006;24(3):116-20.	Less than 50 subjects in relevant age range.
Kanbur NO, Kanli A, Derman O, Eifan A, Atac A. The relationships between dental age, chronological age and bone age in Turkish adolescents with constitutional delay of growth. <i>Journal of Pediatric Endocrinology and Metabolism</i> . 2006;19(8):979-85.	Less than 50 subjects in relevant age range.
Kaplowitz P, Srinivasan S, He J, McCarter R, Hayeri MR, Sze R. Comparison of bone age readings by pediatric endocrinologists and pediatric radiologists using two bone age atlases. <i>Pediatric Radiology</i> . 2011;41(6):690-3.	Less than 50 subjects in relevant age range.
Karami M, Moshirfatem A, Daneshvar P. Age determination using ultrasonography in young football players. <i>Advanced Biomedical Research</i> . 2014;3:174.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Karasik D, Pavlovsky O, Batsevich V, Livshits G, Kobylansky E. Use of the hand bones roentgenographs in the prediction of age in nine human populations. <i>Anthropologischer Anzeiger</i> . 2000;58(2):199-214.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kawashima M, Nanjo B, Sakura H, Ota S. Standard for bone age determination based on x-ray findings of the hand. [Japanese]. <i>Seikeigeka</i> . 1971;0(11):846-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kemp SF, Sy JP. Analysis of bone age data from national cooperative growth study substudy VII. <i>Pediatrics</i> . 1999;104(4):1031-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Khan KM, Gonzalez-Bolanos MT, Holm T, Miller BS, Sarafoglu K. Use of Automated Bone Age for Critical Growth Assessment. <i>Clinical Pediatrics</i> . 2015;54(11):1038-43.	Less than 50 subjects in relevant age range.
Khanna P, Bhargava S. Roentgen assessment of bone density in North Indian population. <i>Indian Journal of Medical Research</i> . 1971;59(10):1599-609.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
King DG, Steventon DM, O'Sullivan MP, Cook AM, Hornsby VP, Jefferson IG, et al. Reproducibility of bone ages when performed by radiology registrars: an audit of Tanner and Whitehouse II versus Greulich and Pyle methods. <i>British Journal of Radiology</i> . 1994;67(801):848-51.	Less than 50 subjects in relevant age range.
Kobylansky E, Livshits G, Pavlovsky O. Population biology of human aging: methods of assessment and sex variation. <i>Human Biology</i> . 1995;67(1):87-109.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kovi R, Korner A, Varkonyi I, Kis E. Bone age discrepancy on hand radiographs. <i>Pediatric Radiology</i> . 2000;30(7):500.	Not an empirical study published in full text format (abstracts, reviews, other).
Kozlowski K. Critical evaluation of radiographic methods of determination of bone age. [Polish]. <i>Pediatria Polska</i> . 1966;41(10):1233-6.	Not an empirical study published in full text format (abstracts, reviews, other).
Kristmundsdottir F, Burwell RG, Marshall WA, Small P. Cross-sectional study of skeletal maturation in normal children from Nottingham and London. <i>Annals of Human Biology</i> . 1984;11(2):133-9.	Less than 50 subjects in relevant age range.
Kucukkeles N, Acar A, Biren S, Arun T. Comparisons between cervical vertebrae and hand-wrist maturation for the assessment of	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

skeletal maturity. <i>Journal of Clinical Pediatric Dentistry</i> . 1999;24(1):47-52.	
Kullman L. Accuracy and precision in some dental radiographic methods. A methodological study with special considerations in age estimation in juveniles. <i>Swedish Dental Journal - Supplement</i> . 1995;104:1-38.	Not an empirical study published in full text format (abstracts, reviews, other).
Kumar MP, Anitha G. Computerized skeletal age assessment. <i>Journal of Clinical Orthodontics</i> . 2010;44(10):598-601; quiz 21.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kyrkanides S, Richter L. Mandibular asymmetry and anterograde notching in individuals with unilateral cleft lip and palate. <i>Cleft Palate-Craniofacial Journal</i> . 2002;39(1):30-5.	Less than 50 subjects in relevant age range.
Lacey KA, Parkin JM, Steel GH. Relationship between bone age and dental development. <i>Lancet</i> . 1973;2(7831):736-7.	Not an empirical study published in full text format (abstracts, reviews, other). and Less than 50 subjects in relevant age range.
Lacey KA, Parkin JM. The normal short child. Community study of children in Newcastle upon Tyne. <i>Archives of Disease in Childhood</i> . 1974;49(6):417-24.	Less than 50 subjects in relevant age range.
Lai XP, Yu XJ, Liu Y. Skeletal age determination in adolescence and the related issues. [Chinese]. <i>Chinese Journal of Forensic Medicine</i> . 2005;20(5):284-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Larsen ST, Demant S, Lynnerup N. GPA, TW2RUS, TW3RUS and FELS-a comparative study on bone-age of the left hand and wrist. <i>American Journal of Physical Anthropology</i> . 2012;147:189.	Presentation withdrawn
Lauterstein AM. A cross-sectional study in dental development and skeletal age. <i>Journal of the American Dental Association</i> . 1961;62:161-7.	Less than 50 subjects in relevant age range.
Lee J, Jung S, Hwang SH, Cho J, Ahn KM, Pyo JW, et al. Rus skeletal maturity of children in urban area in Korea. <i>Endocrine Reviews Conference: 95th Annual Meeting and Expo of the Endocrine Society, ENDO</i> . 2013;34(3).	Not an empirical study published in full text format (abstracts, reviews, other).
Lee J, Kim J, Cho J. Skeletal maturity of radius, ulna, and short bones in TW3 method for children in Korea. <i>Hormone Research in Paediatrics</i> . 2014;82:410.	Not an empirical study published in full text format (abstracts, reviews, other).
Leite HR, O'Reilly MT, Close JM. Skeletal age assessment using the first, second, and third fingers of the hand. <i>American Journal of Orthodontics and Dentofacial Orthopedics</i> . 1987;92(6):492-8.	Less than 50 subjects in relevant age range.
Leonard H, Thomson M, Glasson E, Fyfe S, Leonard S, Ellaway C, et al. Metacarpophalangeal pattern profile and bone age in Rett syndrome: further radiological clues to the diagnosis. <i>American Journal of Medical Genetics</i> . 1999;83(2):88-95.	Less than 50 subjects in relevant age range.
Li K, Ye K, Wang JW, Ye LY, Zhang QC. Study on development of hand-wrist bone in 14 year-old adolescents I: bone age standard for boys. [Chinese]. <i>Fa yi xue za zhi</i> . 2007;23(6):414-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Li K, Ye K, Zhang Z, Wang JW, Ye LY, Zhang QC. Development of hand-wrist bones of 14 year-old adolescents II. Standard of bony age for girls. [Chinese]. <i>Journal of Forensic Medicine</i> . 2008;24(1):15-7+24.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Li XM, Cheng XG, Yu W. The current status of bone age imaging assessment of wrist in China. [Chinese]. <i>Chinese Journal of Radiology (China)</i> . 2013;47(12):1153-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Liebgott B. Dental age: its relation to skeletal age and the time of peak circumpuberal growth in length of the mandible. <i>Dental journal</i> . 1978;44(5):223-7.	Less than 50 subjects in relevant age range.
Liljequist B, Lundberg M. Skeletal and tooth development. A methodologic investigation. <i>Acta Radiologica: Diagnosis</i> . 1971;11(2):97-112.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Liu J, Qi J, Liu Z, Ning Q, Luo X. Automatic bone age assessment based on intelligent algorithms and comparison with TW3 method. <i>Computerized Medical Imaging and Graphics</i> . 2008;32(8):678-84.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Liversidge HM, Buckberry J, Marquez-Grant N. Age estimation. <i>Annals of Human Biology</i> . 2015;42(4):299-301.	Not an empirical study published in full text format (abstracts, reviews, other).

Livshits G, Vainder M, Pavlovsky O, Kobyliansky E. Population biology of human aging: ethnic and climatic variation of bone age scores. <i>Human Biology</i> . 1996;68(2):293-314.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Londhe PS, Ahmed Sheikh N. A roentgen graphic study of wrist joint for age determination in sholapur region of South Maharashtra (India). <i>Indian Journal of Forensic Medicine and Toxicology</i> . 2010;4(1):38-40.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Low WD, Fung SH, Cerny EE. Biological age of southern Chinese boys. <i>Zeitschrift fur Morphologie und Anthropologie</i> . 1976;67(2):136-44.	The study population is not living persons between the ages 10-25 years.
Low WD, Fung SH, Cerny EE. Biological age of southern Chinese girls. <i>Zeitschrift fur Morphologie und Anthropologie</i> . 1976;67(3):264-74.	The study population is not living persons between the ages 10-25 years.
Low WD. Assessing skeletal maturity by inspection and bone-specific methods with the atlas of greulich & pyle. <i>Zeitschrift fur Morphologie und Anthropologie</i> . 1975;67(1):1-5.	The study population is not living persons between the ages 10-25 years.
Ludvigsson J. [Age determination is not racism]. <i>Lakartidningen</i> . 2015;112.	Not an empirical study published in full text format (abstracts, reviews, other)
Luo J, Lin Z, Lu R. A computer system for bone age measurement of hand radiographs. [Chinese]. <i>Zhonghua kou qiang yi xue za zhi = Zhonghua kouqiang yixue zazhi = Chinese journal of stomatology</i> . 1998;33(6):378-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Luo W, Li M, Yun P, Xiao H. [Design and appliation of a kind of bone age automatic analysis and assessment device]. [Chinese]. <i>Zhongguo yi liao qi xie za zhi = Chinese journal of medical instrumentation</i> . 2014;38(4):264-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Madhu S, Hegde AM, Munshi AK. The developmental stages of the middle phalanx of the third finger (MP3): a sole indicator in assessing the skeletal maturity? <i>Journal of Clinical Pediatric Dentistry</i> . 2003;27(2):149-56.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Mahmoodi S, Sharif BS, Chester EG, Owen JP, Lee R. Skeletal growth estimation using radiographic image processing and analysis. <i>IEEE transactions on information technology in biomedicine : a publication of the IEEE Engineering in Medicine and Biology Society</i> . 2000;4(4):292-7.	Less than 50 subjects in relevant age range.
Mainland D. Evaluation of the skeletal age method of estimating children's development. I. Systematic errors in the assessment of roentgenograms. <i>Pediatrics</i> . 1953;12(2):114-29.	The study population is not living persons between the ages 10-25 years
Mainland D. Evaluation of the skeletal age method of estimating children's development. II. Variable errors in the assessment of roentgenograms. <i>Pediatrics</i> . 1954;13(2):165-73.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Malcolm LA. Growth and development of the Kaiapit children of the Markham Valley, New Guinea. <i>American Journal of Physical Anthropology</i> . 1969;31(1):39-51.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Malcolm LA. Growth of the Asai child of the Madang district of New Guinea. <i>Journal of Biosocial Science</i> . 1970;2(3):213-26.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Malina RM, Chamorro M, Serratosa L, Morate F. TW3 and Fels skeletal ages in elite youth soccer players. <i>Annals of Human Biology</i> . 2007;34(2):265-72.	Less than 50 subjects in relevant age range.
Malina RM. Skeletal age and age verification in youth sport. <i>Sports medicine (Auckland, NZ)</i> . 2011;41(11):925-47.	Not an empirical study published in full text format (abstracts, reviews, other).
Manos GK, Cairns AY, Rickets IW, Sinclair D. Segmenting radiographs of the hand and wrist. <i>Computer Methods and Programs in Biomedicine</i> . 1994;43(3):227-37.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Marcondes E, Rumel A, Schvartsman S. [Determination of bone and tooth age by radiographic examination of children from a middle socio-economic class]. <i>Revista Da Faculdade de Odontologia Da Universidade de Sao Paulo</i> . 1965;3(1):185-91.	Less than 50 subjects in relevant age range.
Marshall D. Radiographic correlation of hand, wrist, and tooth development. <i>Dental Radiography & Photography</i> . 1976;49(3):51-72.	Not an empirical study published in full text format (abstracts, reviews, other).
Marshall WA. Individual variations in rate of skeletal maturation. <i>Nature</i> . 1969;221(5175):91.	Not an empirical study published in full text format (abstracts, reviews, other).

Martin DD, Heckmann C, Jenni OG, Ranke MB, Binder G, Thodberg HH. Metacarpal thickness, width, length and medullary diameter in children--reference curves from the First Zurich Longitudinal Study. <i>Osteoporosis International</i> . 2011;22(5):1525-36.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Martin DD, Heckmann C, Neuhof J, Jenni OG, Ranke MB, Binder G. Comparison of radiogrammetrical metacarpal indices in children and reference data from the First Zurich Longitudinal Study. <i>Pediatric Radiology</i> . 2012;42(8):982-91.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Martin DD, Neuhofer J, Jenni OG, Ranke MB, Thodberg HH. Automatic determination of left- and right-hand bone age in the first Zurich longitudinal study. <i>Hormone Research in Paediatrics</i> . 2010;74(1):50-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Martin DD, Sato K, Sato M, Thodberg HH, Tanaka T. Validation of a new method for automated determination of bone age in Japanese children. <i>Hormone research in paediatrics</i> . 2010;73(5):398-404.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Martin DD, Schittenhelm J, Ranke MB, Binder G, Thodberg HH. Validation of a new adult height prediction method based on automated determination of bone age in a French population. <i>Hormone Research in Paediatrics</i> . 2010;74:200.	Not an empirical study published in full text format (abstracts, reviews, other).
Martin DD, Schittenhelm J, Thodberg HH. Validation of adult height prediction based on automated bone age determination in the Paris Longitudinal Study of healthy children. <i>Pediatric Radiology</i> . 2016;46(2):263-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Martorell R, Yarbrough C, Himes JH, Klein RE. Sibling similarities in number of ossification centers of the hand and wrist in a malnourished population. <i>Human Biology</i> . 1978;50(1):73-81.	Another objective than to compare age estimation using x-ray of the hand with known chronological age. Less than 50 subjects in relevant age range.
Mc KT. Estimation of skeletal age from combined maturational activity. <i>American Journal of Physical Anthropology</i> . 1957;15(3):399-408.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
McCormack MK, Scarr-Salapatek S, Polesky H, Thompson W, Katz SH, Barker WB. A comparison of the physical and intellectual development of black children with and without sickle-cell trait. <i>Pediatrics</i> . 1975;56(6):1021-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Medicus H, Gron AM, Moorrees CF. Reproducibility of rating stages of osseous development. (Tanner-Whitehouse system). <i>American Journal of Physical Anthropology</i> . 1971;35(3):359-72.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Mellits ED, Dorst JP, Cheek DB. Bone age: its contribution to the prediction of maturational or biological age. <i>American Journal of Physical Anthropology</i> . 1971;35(3):381-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Memon N, Memon MU, Junejo MA, Memon J. Age determination by employing radiological technique in pediatric age groups. <i>Journal of the Liaquat University of Medical and Health Sciences</i> . 2011;10(2):53-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Memon N, Memon MU, Memon K, Junejo H, Memon J. Radiological indicators for determination of age of consent and criminal responsibility. <i>Journal of the Liaquat University of Medical and Health Sciences</i> . 2012;11(2):64-70.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Menzel HJ, Vilser C, Eulensteiner M, Schwartz T, Vogt S, Bottcher J, et al. Assessment of skeletal age at the wrist in children with a new ultrasound device. <i>Pediatric Radiology</i> . 2005;35(4):429-33.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Menzel HJ, Vogt S, Vilser C, Schwartz T, Eulensteiner M, Bottcher J, et al. Assessment of skeletal age using a new ultrasound method. [German]. <i>RoFo Fortschritte auf dem Gebiet der Rontgenstrahlen und der Bildgebenden Verfahren</i> . 2005;177(12):1699-705.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Milner GR, Levick RK, Kay R. Assessment of bone age: a comparison of the Greulich and Pyle, and the Tanner and Whitehouse methods. <i>Clinical Radiology</i> . 1986;37(2):119-21.	Less than 50 subjects in relevant age range.
Mimouni M, Shuper A, Mimouni F, Grunebaum M, Varsano I. Retarded skeletal maturation in children with primary enuresis. <i>European Journal of Pediatrics</i> . 1985;144(3):234-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Mohite DP, Chaudhary MS, Mohite PM, Patil SP. Age assessment from mandible: Comparison of radiographic and histologic methods. <i>Romanian Journal of Morphology and Embryology</i> . 2011;52(2):659-68.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Molinari L, Gasser T, Largo R. A comparison of skeletal maturity and growth. <i>Annals of Human Biology</i> . 2013;40(4):333-40.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Molinari L, Gasser T, Largo RH. TW3 bone age: RUS/CB and gender differences of percentiles for score and score increments. <i>Annals of Human Biology</i> . 2004;31(4):421-35.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Moore RN, Moyer BA, DuBois LM. Skeletal maturation and cranio-facial growth. <i>American Journal of Orthodontics & Dentofacial Orthopedics</i> . 1990;98(1):33-40.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Moore WM. Comparability in skeletal maturation research. <i>American Journal of Physical Anthropology</i> . 1971;35(3):411-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Morris JM, Park JH. Correlation of dental maturity with skeletal maturity from radiographic assessment. <i>Journal of Clinical Pediatric Dentistry</i> . 2012;36(3):309-14.	Not an empirical study published in full text format (abstracts, reviews, other).
Murray JR, Bock RD, Roche AF. The measurement of skeletal maturity. <i>American Journal of Physical Anthropology</i> . 1971;35(3):327-30.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Mustafa S, Raj AC, Anekar J, Divakar DD, Al Kheraif AA, Ramakrishnaiah R, et al. Evaluation of dental and skeletal maturity using digital panoramic radiographs and digital cephalograms. <i>Asian Biomedicine</i> . 2015;9(3):335-42.	Less than 50 subjects in relevant age range.
Nahhas RW, Sherwood RJ, Chumlea WC, Towne B, Duren DL. Predicting the timing of maturational spurts in skeletal age. <i>American Journal of Physical Anthropology</i> . 2013;150(1):68-75.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Nicoletti I, Cheli D. [Current methods for the evaluation of the skeletal maturation. 1. Adaptation of the T.W.2 method to the Italian population]. <i>Minerva Pediatrica</i> . 1983;35(21):1083.	Not an empirical study published in full text format (abstracts, reviews, other).
Niedzwiecka Z, Kopczynska-Sikorska J. Determination of the skeletal age of Warsaw children by the TW 2 method. [Polish]. <i>Pediatrica polska</i> . 1985;60(9):625-30.	The study population is not living persons between the ages 10-25 years
Ning G, Qu HB, Liu GJ, Wu KM, Xie SX, Chen XJ. [TW systems in estimation of carpal bone age and their potential applications in diagnosis of idiopathic precocious puberty in Chinese girls]. <i>Fa i Hsueh Tsa Chih Journal of Forensic Medicine</i> . 2007;23(2):97-100.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Nuhoglu B, Ayyildiz A, Fidan V, Cebeci O, Kosar U, Germiyanoğlu C. Do children with primary nocturnal enuresis have a retarded bone age? A cross-sectional study. <i>International Journal of Urology</i> . 2006;13(2):109-10.	Less than 50 subjects in relevant age range.
Ochoa BK, Nanda RS. Comparison of maxillary and mandibular growth. <i>American Journal of Orthodontics & Dentofacial Orthopedics</i> . 2004;125(2):148-59.	Less than 50 subjects in relevant age range.
Oestreich AE. Tanner-Whitehouse versus Greulich-Pyle in bone age determinations. <i>Journal of Pediatrics</i> . 1997;131(1):5-6.	Not an empirical study published in full text format (abstracts, reviews, other).
Omelka F, Bystry J, Pohlidal A, Tuma A. [Bone age in boys and adolescents and their growth (final report of a long-term study in Brno)]. <i>Ceskoslovenska Pediatrie</i> . 1985;40(3):152-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Omondi O. Age estimation in clinical and public health research. <i>East African Medical Journal</i> . 1985;62(12):861-76.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ortega-Pertuz AI, Haiter-Neto F, Costa ALF. Atlasbased methods or TW3?- Estimation of bone age in Brazilians. [Portuguese]. <i>Medicina (Brazil)</i> . 2015;48(2):129-37.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ostojic SM. Prediction of adult height by Tanner-Whitehouse method in young Caucasian male athletes. <i>Ojm</i> . 2013;106(4):341-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ottow C, Kramer JA, Olze A, Schmidt S, Schulz R, Wittschieber D, et al. Magnetic resonance tomography studies on age estimation of	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

unaccompanied minor refugees. [German]. Rechtsmedizin. 2015;25(1):12-20.	
Ousley S, Frazee K, Stull K. Assessing bone growth and development in modern American children. American Journal of Physical Anthropology. 2010;141:183.	Not an empirical study published in full text format (abstracts, reviews, other).
Pan FP. [A longitudinal study of the relationship between skeletal age and adolescence]. Chung-Hua Yu Fang i Hsueh Tsa Chih [Chinese Journal of Preventive Medicine]. 1985;19(2):79-82.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Park E. Radiological anthropometry of the hand in Turner's syndrome. American Journal of Physical Anthropology. 1977;46(3):463-70.	A study population with chronic disease or developmental disorders.
Pashkova VI, Burov SA. [Possibility of using standard indices of skeletal ossification for the forensic medical expertise of determining the age of children and adolescents living throughout the whole territory of the USSR]. Sudebno-Meditsinskaia Ekspertiza. 1980;23(3):22-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pelech L, Borsky M, Havlik J, Bleha P. Estimation of biological age by determining the degree of bone and tooth maturation (Czech). [Czech]. Ceskoslovenska Hygiena. 1976;21(4):180-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pelech L, Pechstein J. [Comparative studies of skeletal age in Czech and Soth German children]. Arztliche Jugendkunde. 1976;67(1):23-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pelech L. Skeletal maturation of children living in areas with a different contamination of the atmosphere. Czechoslovak Medicine. 1978;1(4):203-10.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pichai S, Rajesh M, Reddy N, Adusumilli G, Reddy J, Joshi B. A comparison of hand wrist bone analysis with two different cervical vertebral analysis in measuring skeletal maturation. Journal of International Oral Health. 2014;6(5):36-41.	Less than 50 subjects in relevant age range.
Pickett KE, Haas JD, Murdoch SM, Rivera JA, Martorell R. Early nutritional supplementation and skeletal maturation in Guatemalan adolescents. Journal of Nutrition. 1995;125(4):1097S-103S.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pietka E, Gertych A, Pospiech S, Cao F, Huang HK, Gilsanz V. Computer-assisted bone age assessment: image preprocessing and epiphyseal/metaphyseal ROI extraction. IEEE Transactions on Medical Imaging. 2001;20(8):715-29.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pietka E, Gertych A, Pospiech Euro Kurkowska S, Cao F, Huang HK, Gilzanz V. Computer-assisted bone age assessment: graphical user interface for image processing and comparison. Journal of digital imaging : the official journal of the Society for Computer Applications in Radiology. 2004;17(3):175-88.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pietka E, McNitt-Gray MF, Kuo ML, Huang HK. Computer-assisted phalangeal analysis in skeletal age assessment. IEEE Transactions on Medical Imaging. 1991;10(4):616-20.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Pietka E, Pospiech-Kurkowska S, Gertych A, Cao F. Integration of computer assisted bone age assessment with clinical PACS. Computerized Medical Imaging and Graphics. 2003;27(2):217-28.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Polito C, Di Toro A, Collini R, Cimmaruta E, D'Alfonso C, Del Giudice G. Advanced RUS and normal carpal bone age in childhood obesity. International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity. 1995;19(7):506-7.	Less than 50 subjects in relevant age range.
Porcu E, Venturoli S, Fabbri R, Paradisi R, Longhi M, Sganga E, et al. Skeletal maturation and hormonal levels after the menarche. Archives of Gynecology & Obstetrics. 1994;255(1):43-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Post GB, Kemper HC. Nutrient intake and biological maturation during adolescence. The Amsterdam growth and health longitudinal study. European Journal of Clinical Nutrition. 1993;47(6):400-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Poulton A, Evans R, Melzer E, Bui Q. Stimulant medication affects growth but not bone age. Archives of Disease in Childhood. 2014;99:A4-A5.	Not an empirical study published in full text format (abstracts, reviews, other).

Poulton AS, Bui Q, Melzer E, Evans R. Stimulant medication effects on growth and bone age in children with attention-deficit/hyperactivity disorder: a prospective cohort study. International Clinical Psychopharmacology. 2016;31(2):93-9.	The study population is not living persons between the ages 10-25 years, Less than 50 subjects in relevant age range.
Powell S, Ma D, Jones G. Determinants of skeletal age deviation in a cross-sectional study. Journal of Clinical Endocrinology and Metabolism. 2008;93(2):521-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Prakash S. Age and order of appearance of elbow and hand-wrist ossific centres in Punjabee children from Rohtak. Indian Journal of Medical Research. 1975;63(5):640-51.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Prasad CK, Reddy VN, Sreedevi G, Ponnada SR, Priya KP, Naik BR. Objective evaluation of cervical vertebral bone age' its reliability in comparison with hand-wrist bone age: by TW3 method. The journal of contemporary dental practice. 2013;14(5):806-13.	Less than 50 subjects in relevant age range.
Prasad M, Ganji VS, George SA, Talapaneni AK, Shetty SK. A comparison between cervical vertebrae and modified MP3 stages for the assessment of skeletal maturity. Journal of Natural Science Biology & Medicine. 2013;4(1):74-80.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Prokopec M, Pfeiferova K, Josifko M. Ossification of the sesamoid bone at the base of the first finger in Czech boys and girls. Central European Journal of Public Health. 1997;5(4):155-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Rachmiel M, Naugolny L, Mazor-Aronovitch K, Levin A, Koren-Morag N, Bistritzer T. Bone age assessment by a novel quantitative ultrasound based device, SonicBone, is comparable to the conventional Greulich and Pyle method. Hormone Research in Paediatrics. 2013;80:35.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Rajagopal R, Kansal S. A comparison of modified MP3 stages and the cervical vertebrae as growth indicators. Journal of Clinical Orthodontics. 2002;36(7):398-406.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Rajic MK, Brisson GR, Shephard RJ, Lavallee H, Jequier JC, Masse R, et al. Bone maturity and physical performance. [French]. Canadian journal of applied sport sciences. 1979;0(3):223-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Davila GH, Eyman SL. A comparison between Greulich-Pyle and Tanner-Whitehouse assessments of skeletal maturity. Radiology. 1971;98(2):273-80.	Less than 50 subjects in relevant age range.
Roche AF, Davila GH. Associations between the rates of maturation and growth in the short bones of the hand. American Journal of Physical Anthropology. 1970;33(3):349-55.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Davila GH. The reliability of assessments of the maturity of individual hand-wrist bones. Human Biology. 1976;48(3):585-97.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Eyman SL, Davila GH. Skeletal age prediction. Journal of Pediatrics. 1971;78(6):997-1003.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, French NY. Differences in skeletal maturity levels between the knee and hand. The American journal of roentgenology, radium therapy, and nuclear medicine. 1970;109(2):307-12.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Johnson JM. A comparison between methods of calculating skeletal age (Greulich-Pyle). American Journal of Physical Anthropology. 1969;30(2):221-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Rohmann CG, French NY, Davila GH. Effect of training on replicability of assessments of skeletal maturity (Greulich-Pyle). The American journal of roentgenology, radium therapy, and nuclear medicine. 1970;108(3):511-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF, Wainer H, Thissen D. The RWT method for the prediction of adult stature. Pediatrics. 1975;56(6):1027-33.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Roche AF. Assessment of skeletal maturation. Summary. American Journal of Physical Anthropology. 1971;35(3):467-9.	Not an empirical study published in full text format (abstracts, reviews, other).
Roche AF. Associations between the rates of maturation of the bones of the hand-wrist. American Journal of Physical Anthropology. 1970;33(3):341-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Roche AF. Fels section for the assessment of growth and maturity. <i>American Journal of Clinical Nutrition</i> . 1978;31(6):908.	Not an empirical study published in full text format (abstracts, reviews, other).
Rodriguez Larralde A, Arias Cazorla S. [Regression equations for age determination by the length of long bones of the hand in the Venezuelan population during growth]. <i>Acta Cientifica Venezolana</i> . 1980;31(5):468-74.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Rucci M, Coppini G, Nicoletti I, Cheli D, Valli G. Automatic analysis of hand radiographs for the assessment of skeletal age: A subsymbolic approach. <i>Computers and Biomedical Research</i> . 1995;28(3):239-56.	Less than 50 subjects in relevant age range.
Rudzinski BW. [Puberty and its indices in hand and wrist radiograms in boys from Warsaw schools]. <i>Pediatria Polska</i> . 1970;45(12):1447-61.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ruf S, Pancherz H. Can frontal sinus development be used for the prediction of skeletal maturity at puberty? <i>Acta Odontologica Scandinavica</i> . 1996;54(4):229-34.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ruf S, Pancherz H. Development of the frontal sinus in relation to somatic and skeletal maturity. A cephalometric roentgenographic study at puberty. <i>European Journal of Orthodontics</i> . 1996;18(5):491-7.	Less than 50 subjects in relevant age range.
Rundle AT, Dollimore J. Age-related bone change. <i>Journal of Mental Deficiency Research</i> . 1976;20(1):55-9.	Less than 50 subjects in relevant age range.
Sachan K, Sharma VP, Tandon P. A correlative study of dental age and skeletal maturation. <i>Indian Journal of Dental Research</i> . 2011;22(6):882.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Saint-Martin P, Rerolle C, Dedouit F, Rousseau H, Rouge D, Telmon N. Evaluation of an automatic method for forensic age estimation by magnetic resonance imaging of the distal tibial epiphysis--a preliminary study focusing on the 18-year threshold. <i>International Journal of Legal Medicine</i> . 2014;128(4):675-83.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
San Roman P, Palma JC, Oteo MD, Nevado E. Skeletal maturation determined by cervical vertebrae development. <i>European Journal of Orthodontics</i> . 2002;24(3):303-11.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sarria A, Moreno L, Bueno M. Analysis of the Greulich-Pyle atlas using the TW2 and TW2-A method. [Spanish]. <i>Anales espanoles de pediatria</i> . 1986;24(2):105-10.	Less than 50 subjects in relevant age range.
Sato K, Ashizawa K, Anzo M, Otsuki F, Kaneko S, Tanaka T, et al. Setting up an automated system for evaluation of bone age. <i>Endocrine Journal</i> . 1999;46:S97-100.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sato K, Mito T, Mitani H. An accurate method of predicting mandibular growth potential based on bone maturity. <i>American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics</i> . 2001;120(3):286-93.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Saxena S, Saxena NB. Skeletal maturation of hands and wrists in normal and malnourished children. <i>Indian Journal of Pediatrics</i> . 1980;47(386):187-91.	Less than 50 subjects in relevant age range.
Schmeling A, Olze A, Reisinger W, Geserick G. The impact of ethnicity on characteristics examined for age estimation in criminal law. [German]. <i>Rechtsmedizin</i> . 2001;11(3):78-81.	Not an empirical study published in full text format (abstracts, reviews, other).
Schmeling A, Reisinger W, Geserick G, Olze A. Age estimation of unaccompanied minors. Part I. General considerations. <i>Forensic Science International</i> . 2006;159:S61-4.	Not an empirical study published in full text format (abstracts, reviews, other).
Schmidt S, Baumann U, Schulz R, Reisinger W, Schmeling A. Study of age dependence of epiphyseal ossification of the hand skeleton. <i>International Journal of Legal Medicine</i> . 2008;122(1):51-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Schmidt S, Fracasso T, Pfeiffer H, Schmeling A. Skeletal age determination of the hand. [German]. <i>Rechtsmedizin</i> . 2010;20(6):475-82.	Not an empirical study published in full text format (abstracts, reviews, other).
Schmidt S, Nitz I, Ribbecke S, Schulz R, Pfeiffer H, Schmeling A. Skeletal age determination of the hand: a comparison of methods. <i>International Journal of Legal Medicine</i> . 2013;127(3):691-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Schmidt S, Schiborr M, Pfeiffer H, Schmeling A, Schulz R. Age dependence of epiphyseal ossification of the distal radius in ultrasound diagnostics. International Journal of Legal Medicine. 2013;127(4):831-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Schmidt S, Schramm D, Ribbecke S, Schulz R, Wittschieber D, Olze A, et al. [Forensic age estimation in juveniles and young adults: Reducing the range of scatter in age diagnosis by combining different methods]. Archiv fur Kriminologie. 2016;237(1):25-37.	The study population is not living persons between the ages 10-25 years
Schundeln MM, Marschke L, Bauer JJ, Hauffa PK, Schweiger B, Fuhrer-Sakel D, et al. A Piece of the Puzzle: The Bone Health Index of the BoneXpert Software Reflects Cortical Bone Mineral Density in Pediatric and Adolescent Patients. PLoS ONE [Electronic Resource]. 2016;11(3):e0151936.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Serinelli S, Panebianco V, Martino M, Battisti S, Rodacki K, Marinelli E, et al. Accuracy of MRI skeletal age estimation for subjects 12-19. Potential use for subjects of unknown age. International Journal of Legal Medicine. 2015;129(3):609-17.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Serinelli S, Panetta V, Pasqualetti P, Marchetti D. Accuracy of three age determination X-ray methods on the left hand-wrist: a systematic review and meta-analysis. Legal Medicine. 2011;13(3):120-33.	Not an empirical study published in full text format (abstracts, reviews, other).
Shakir A, Zaini S. Skeletal maturation of the hand and wrist of young children in Baghdad. Annals of Human Biology. 1974;1(2):189-99.	The study population is not living persons between the ages 10-25 years
Sharat S, Khanduja PC, Agarwal KN, Saha MM, Gupta S, Bhardwaj OP. Skeletal growth in school children. Indian Pediatrics. 1970;7(2):98-108.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Shin SM, Kim YI, Choi YS, Yamaguchi T, Maki K, Cho BH, et al. The skeletal maturation status estimated by statistical shape analysis: axial images of Japanese cervical vertebra. Dento-Maxillo-Facial Radiology. 2015;44(3):20140323.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sierra AM. Assessment of dental and skeletal maturity. A new approach. Angle Orthodontist. 1987;57(3):194-208.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Soegiharto BM, Moles DR, Cunningham SJ. Discriminatory ability of the skeletal maturation index and the cervical vertebrae maturation index in detecting peak pubertal growth in Indonesian and white subjects with receiver operating characteristics analysis. American Journal of Orthodontics and Dentofacial Orthopedics. 2008;134(2):227-37.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Somkantha K, Theera-Umporn N, Auephanwiriyakul S. Bone age assessment in young children using automatic carpal bone feature extraction and support vector regression. Journal of digital imaging : the official journal of the Society for Computer Applications in Radiology. 2011;24(6):1044-58.	The study population is not living persons between the ages 10-25 years
Souguir MK, Ben Dhiab M, Masmoudi T, Zemni M. Age determination among Tunisian young people, 200 cases related. [French]. Journal de Medecine Legale Droit Medical. 2002;45(2):89-91.	Not an empirical study published in full text format (abstracts, reviews, other).
Speed B. Limitations of the Greulich and Pyle atlas method of skeletal age estimation in South African children. Clinical Anatomy. 2012;25:926.	Not an empirical study published in full text format (abstracts, reviews, other).
Sproul A, Peritz E. Assessment of skeletal age in short and tall children. American Journal of Physical Anthropology. 1971;35(3):433-9.	The study population is not living persons between the ages 10-25 years
Stasiowska B, Benso L, Pastorin L, Giachino M, Nigro N. [Comparison of 2 methods of evaluation of skeletal maturation. 2. Analysis of the correlation with another biological variable]. Minerva Pediatrica. 1978;30(21):1665-77.	The study population is not living persons between the ages 10-25 years
Steel GH. A Correlation Study between Dental and Skeletal Maturity. Report of the Congress; European Orthodontic Society. 1964;40:83-91.	Not an empirical study published in full text format (abstracts, reviews, other).
Stiehl J, Muller B, Dibbets J. The development of the cervical vertebrae as an indicator of skeletal maturity: comparison with the classic method of hand-wrist radiograph. Journal of Orofacial Orthopedics. 2009;70(4):327-35.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Sugiura Y. Age determination by skeleton. [Japanese]. Nippon Seikeigeka Gakkai zasshi. 1970;44(11):1021-34.	Less than 50 subjects in relevant age range.
Suma G, Rao BB, Annigeri RG, Rao DJ, Goel S. Radiographic correlation of dental and skeletal age: Third molar, an age indicator. Journal of forensic dental sciences : JFDS. 2011;3(1):14-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sundick RI. Age and sex determination of subadult skeletons. Journal of Forensic Sciences. 1977;22(1):141-4.	Not an empirical study published in full text format (abstracts, reviews, other).
Surendran S, Thomas E. Tooth mineralization stages as a diagnostic tool for assessment of skeletal maturity. American Journal of Orthodontics & Dentofacial Orthopedics. 2014;145(1):7-14.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Suttapreyasri D. A standard of radiographic bone age of Thai children. Journal of the Medical Association of Thailand = Chotmaihet thangphaet. 1979;62(6):310-4.	Less than 50 subjects in relevant age range.
Suzuki Y. [Study of development of the carpal bones of children. 1. Normal developmental pattern]. Nippon Shonika Gakkai Zasshi [Journal of Japan Pediatric Society]. 1968;72(5):541-52.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sypniewski M, Mikusek J. The correlation between skeletal age assessed by the method of Tanner et al., and dental age assessed by means of tables of Lucas and Matjeka. [Polish]. Protetyka stomatologiczna. 1975;25(1):21-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Szabo S, Doka J, Apor P, Somogyvari K. [Relationship between bone age, functional anthropometric data and aerobic capacity]. Schweizerische Zeitschrift fur Sportmedizin. 1972;20(3):109-15.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Takai S. Velocities for the Tanner-Whitehouse 2 skeletal maturity in northwest Japanese children. Okajimas Folia Anatomica Japonica. 1993;70(2):119-26.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Tanner JM, Gibbons RD. A computerized image analysis system for estimating Tanner-Whitehouse 2 bone age. Hormone Research. 1994;42(6):282-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Tanner JM, Gibbons RD. Automatic bone age measurement using computerized image analysis. Journal of Pediatric Endocrinology. 1994;7(2):141-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Tanner JM, Oshman D, Lindgren G, Grunbaum JA, Elsouki R, Labarthe D. Reliability and validity of computer-assisted estimates of Tanner-Whitehouse skeletal maturity (CASAS): comparison with the manual method. Hormone Research. 1994;42(6):288-94.	Less than 50 subjects in relevant age range.
Thodberg HH, Jenni OG, Caflisch J, Ranke MB, Martin DD. Prediction of adult height based on automated determination of bone age. Journal of Clinical Endocrinology & Metabolism. 2009;94(12):4868-74.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg HH, Jenni OG, Ranke MB, Martin DD. Standardization of the Tanner-Whitehouse bone age method in the context of automated image analysis. Annals of Human Biology. 2012;39(1):68-75.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg HH, Kreiborg S, Juul A, Pedersen KD. The BoneXpert method for automated determination of skeletal maturity. IEEE Transactions on Medical Imaging. 2009;28(1):52-66.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg HH, Lomholt J, Juul A, Martin DD, Kreiborg S. Review and further validation of a new adult height prediction method based on automated determination of bone age. Hormone Research in Paediatrics. 2010;74:200.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg HH, Martin DD. Reference databases of radiogrammetric Bone Health Index from hand radiographs of 1000 normal children of four ethnicities. Bone. 2009;45:S110.	Not an empirical study published in full text format (abstracts, reviews, other).
Thodberg HH, Neuhof J, Ranke MB, Jenni OG, Martin DD. Validation of bone age methods by their ability to predict adult height. Hormone research in pediatrics. 2010;74(1):15-22.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg HH, Savendahl L. Validation and reference values of automated bone age determination for four ethnicities. Academic Radiology. 2010;17(11):1425-32.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thompson GG, Blanksby BA, Doran GA. Proceedings: Interrelationship between swimming performance and chronological age, dental age, skeletal age of age group competitive swimmers. Journal of Anatomy. 1973;116:483.	Not an empirical study published in full text format (abstracts, reviews, other).

Tian Z, Yan Y. Tooth mineralization stages as a diagnostic tool for assessment of skeletal maturity. <i>American Journal of Orthodontics and Dentofacial Orthopedics</i> . 2014;145(5):551-2.	Not an empirical study published in full text format (abstracts, reviews, other).
Triratana T, Phengpinij S, Clearpongse N. [Correlation between age and developmental stages of permanent teeth and developmental stages of hand-wrist skeletal]. <i>Journal of the Dental Association of Thailand</i> . 1989;39(2):57-65.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Tristan-Vega A, Arribas JI. A radius and ulna TW3 bone age assessment system. <i>IEEE Transactions on Biomedical Engineering</i> . 2008;55(5):1463-76.	Not an empirical study published in full text format (abstracts, reviews, other).
Van der Werff ten B. Skeletal maturation. <i>Folia Medica Nederlandica</i> . 1968;11(5):209-18.	Not an empirical study published in full text format (abstracts, reviews, other).
Van G. [Problems in Using the Greulich-Pyle Skeletal Atlas and the Bailey Tables with Dutch Children]. <i>Maandschrift voor Kinder-geneeskunde</i> . 1965;33:121-35.	A study population with chronic disease or developmental disorders.
van Rijn RR, Lequin MH, Thodberg HH. Automatic determination of Greulich and Pyle bone age in healthy Dutch children. <i>Pediatric Radiology</i> . 2009;39(6):591-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Van Teunenbroek A, De Waal W, Roks A, Chinafo P, Fokker M, Mulder P, et al. Computer-aided skeletal age scores in healthy children, girls with Turner syndrome, and in children with constitutionally tall stature. <i>Pediatric Research</i> . 1996;39(2):360-7.	A study population with chronic disease or developmental disorders.
Varkkola O, Ranta H, Metsaniitty M, Sajantila A. Age assessment by the Greulich and Pyle method compared to other skeletal X-ray and dental methods in data from Finnish child victims of the Southeast Asian Tsunami. <i>Forensic Science, Medicine, and Pathology</i> . 2011;7(4):311-6.	The study population is not living persons between the ages 10-25 years
Vejvoda M, Grant DB. Discordant bone maturation of the hand in children with precocious puberty and congenital adrenal hyperplasia. <i>Acta Paediatrica Scandinavica</i> . 1981;70(6):903-5.	Less than 50 subjects in relevant age range.
Vidal-Linhares R, De Oliveira-Matta M, Perrout-Lima J, Barros-Costa M, Fernandes-Filho J. [The relationship between adolescents' dermatoglyphic characteristics and skeletal maturation]. [Portuguese]. <i>Revista de salud publica (Bogota, Colombia)</i> . 2010;12(6):929-37.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Vijayaraghavan R, Pandiarajan V, Subramanian A, Viswanathan J. Effect of protein energy malnutrition and vitamin A deficiency on the bones of hand and wrist. <i>Indian Pediatrics</i> . 1982;19(9):753-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Viol M. Estimation of biological age by Medved's modified radiographic method. [German]. <i>Medizin und Sport</i> . 1977;17(1):15-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Von Harnack GA. Determination of skeletal maturation in childhood. [German]. <i>ZGeburtshPerinatologie</i> . 1974;178(4):237-44.	Not an empirical study published in full text format (abstracts, reviews, other).
Walker GF. Forensic medicine. <i>New Zealand Dental Journal</i> . 1969;65(299):46-7.	Not an empirical study published in full text format (abstracts, reviews, other).
Wang A, Yang F, Yu B, Shan Y, Gao L, Zhang X, et al. [Hand and wrist bone maturation in children with central precocious puberty and idiopathic short stature]. [Chinese]. <i>Zhejiang da xue xue bao</i> . 2013;0(4):411-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Wang P, Zhu GY, Wang YH, Fan LH, Zhang GZ, Ying CL, et al. Assessment of skeletal age in Chinese male adolescents. [Chinese]. <i>Journal of Forensic Medicine</i> . 2008;24(4):252-5+8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Wang Y, Wang Z, Wei H, Wan L, Ying C, Zhu G. Automated assessment of developmental levels of epiphysis by support vector machine. [Chinese]. <i>Fa yi xue za zhi</i> . 2014;30(6):422-6.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Wang YH, Ying CL, Wan L, Zhu GY. Long-term trend of bone development in the contemporary teenagers of Chinese Han nationality. <i>Journal of Forensic Medicine</i> . 2012;28(4):269-74.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Wang YH, Zhu GY, Wang P, Fan LH, Zhang GZ, Ying CL, et al. [Mathematical models of forensic bone age assessment of living subjects in Chinese Han female teenagers]. <i>Fa i Hsueh Tsa Chih Journal of Forensic Medicine</i> . 2008;24(2):110-3.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

Wenzel A, Melsen B. Replicability of assessing radiographs by the Tanner and Whitehouse-2 method. Human Biology. 1982;54(3):575-81.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Wenzel A, Thylstrup A, Melsen B, Fejerskov O. The relationship between water-borne fluoride, dental fluorosis and skeletal development in 11-15 year old Tanzanian girls. Archives of Oral Biology. 1982;27(12):1007-11.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Willich E. Skeletal age determination in adolescence and the 3d decade of life. [German]. Der Radiologe. 1986;26(5):227-9.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Winkielman J. Comparison of the standards method and the Tanner-Whitehouse method in the estimation of skeletal age for clinical purposes. [Polish]. Pediatria polska. 1981;56(5):527-30.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Ye LY, Li K, Ye K, Zhang QC, Liu ZH, Wen XM, et al. Trend of bone development of adolescent at 14 years old. [Chinese]. Journal of Forensic Medicine. 2008;24(3):182-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang GD. A study of skeletal age in 250 pairs of twins. [Chinese]. Zhonghua yu fang yi xue za zhi [Chinese journal of preventive medicine]. 1985;19(6):321-3.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang S, Zhang J, Liu L, Liu G. Nondiscernable epiphyseal line of radius and ulna as the indicator to estimate adolescent age of 18 years old. [Chinese]. Chinese Journal of Forensic Medicine. 2010;25(2):100-1.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang S, Zhang J. The age estimation for overweight, obesity and thinness in 7146 urban adolescents. [Chinese]. Chinese Journal of Forensic Medicine. 2010;25(4):228-31.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang SC. [A longitudinal study of bone age]. Chung-Hua Kou Chiang i Hsueh Tsa Chih Chinese Journal of Stomatology. 1991;26(2):99-102.	Less than 50 subjects in relevant age range.
Zhang SY, Liu G, Ma CG, Han YS, Shen XZ, Xu RL, et al. Automated determination of bone age in a modern chinese population. ISRN Radiology. 2013;2013:874570.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang SY, Tanaka T, Martin D, Liu G, Ma CG, Han YS, et al. Adult height prediction for Han children based on automated bone age determination. Hormone Research in Paediatrics. 2012;78:144.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang Y, Wang B. Observation of cervical vertebrae and estimation of their bone age. [Chinese]. Zhonghua kou qiang yi xue za zhi = Zhonghua kouqiang yixue zazhi = Chinese journal of stomatology. 1997;32(3):152-4.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhang Z, Li K, Yu RJ, Zhang Q. Preliminary study on the applying value of two measurements for bone age in the cases of minors. [Chinese]. Fa yi xue za zhi. 2004;20(4):212-4.	Less than 50 subjects in relevant age range.
Zhao H, Dong XA, Zheng T, Qing SH, Deng ZH, Zhu GY. Skeletal age estimation of sternal end of clavicle in Sichuan Han nationality youth using thin-section computed tomography. [Chinese]. Journal of Forensic Medicine. 2011;27(6):417-20.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zhu GY, Fan LH, Zhang GZ, Ying CL, Lu X, Xia WT, et al. [Staging methods of skeletal growth by X-ray in teenagers]. Fa i Hsueh Tsa Chih Journal of Forensic Medicine. 2008;24(1):18-24.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Zink P, Zink I, Reinhardt G. [Roentgen image of the hand as a principle for age determination in adolescents]. Archiv fur Kriminologie. 1986;178(1):15-24.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Aicardi, G., et al., Assessment of skeletal maturity of the hand-wrist and knee: A comparison among methods. American Journal of Human Biology, 2000. 12(5): p. 610-615.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Galescu, O.A., et al., Obesity and predicted height in children: A longitudinal analysis. Endocrine Reviews. Conference: 97th Annual Meeting and Expo of the Endocrine Society, ENDO, 2015. 36(no pagination).	Not an empirical study published in full text format (abstracts, reviews, other).
Hassan, N., et al., Age of fusion of the distal radial and ulnar epiphyses from hand radiographs-A study in Kashmiri population. Science & Justice, 2016. 56(6): p. 431-436.	Not relevant methods.

Sypek, S.A., et al., A holistic approach to age estimation in refugee children. <i>Journal of Paediatrics & Child Health</i> , 2016. 52(6): p. 614-20.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Thodberg, H.H., et al., Automated determination of bone age from hand X-rays at the end of puberty and its applicability for age estimation. <i>International Journal of Legal Medicine</i> , 2016. 18: p. 18.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Hsieh CW, Liu TC, Wang JK, Jong TL, Tiu CM. Simplified radius, ulna, and short bone-age assessment procedure using grouped-Tanner-Whitehouse method. <i>Pediatrics International</i> . 2011;53(4):567-75.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Johnston FE. The use of the Greulich-Pyle method in a longitudinal growth study. <i>American Journal of Physical Anthropology</i> . 1971;35(3):353-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Kadam SS, Viswanathan KG. Age estimation radiologically from epiphyseal union at wrist joint among subjects in Davanagere district. <i>Indian Journal of Forensic Medicine and Toxicology</i> . 2012;6(2):195-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Sahni D, Jit I. Time of fusion of epiphyses at the elbow and wrist joints in girls of northwest India. <i>Forensic Science International</i> . 1995;74(1):47-55.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Theintz GE, Howald H, Allemann Y, Sizonenko PC. Growth and pubertal development of young female gymnasts and swimmers: a correlation with parental data. <i>International Journal of Sports Medicine</i> . 1989;10(2):87-91.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Napoli MM, Saraiva PA. [Bone age]. <i>Revista do Hospital das Clinicas; Faculdade de Medicina Da Universidade de Sao Paulo</i> . 1985;40(5):210-5.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Geier M, Dahlmann H. [Differences in growth and development of patients with cleft lip and palate and normal subjects]. <i>Fortschritte der Kieferorthopadie</i> . 1988;49(5):444-53.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Fu CA, Henson T, Bethard JD. Validation study of the Ubelaker and London atlases. <i>American Journal of Physical Anthropology</i> . 2014;153:122.	Not an empirical study published in full text format (abstracts, reviews, other).
Pechnikova M, Gibelli D, De Angelis D, de Santis F, Cattaneo C. The "blind age assessment": applicability of Greulich and Pyle, Demirjian and Mincer aging methods to a population of unknown ethnic origin. <i>La Radiologia medica</i> . 2011;116(7):1105-14.	Less than 50 subjects in relevant age range.
Gindhart PS. Growth standards for the tibia and radius in children aged one month through eighteen years. <i>American Journal of Physical Anthropology</i> . 1973;39(1):41-8.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.
Giordano D, Leonardi R, Maiorana F, Scarciofalo G, Spampinato C. Epiphysis and metaphysis extraction and classification by adaptive thresholding and DoG filtering for automated skeletal bone age analysis. <i>Conference proceedings</i> : . 2007;0:6552-7.	Another objective than to compare age estimation using x-ray of the hand with known chronological age.

References that we did no consider because the full text articles were expensive to buy or difficult to procure:
Baransel Isir A, Buken B, Tokdemir M, Dulger HE, Erel O, Fedakar R. Assessing the age determination cases which have been analyzed at forensic medicine departments of 5 different region's universities in Turkey between years 1998-2005. [Turkish]. <i>Turkiye Klinikleri Journal of Medical Sciences</i> . 2009;29(2):304-13.
Berchel C. [Bone growth and skeletal maturation in children]. <i>Soins; La Revue de Reference Infirmiere</i> . 1977;22(9):3-6.
Bleha P, Havlik J, Borsky M, Pelech L. [Relationship between dental and bone maturation]. <i>Ceskoslovenska Stomatologie</i> . 1976;76(1):15-24.
Fragoso I, Massuca LM, Ferreira J. Effect of birth month on physical fitness of soccer players (Under-15) according to biological maturity. <i>International Journal of Sports Medicine</i> . 2015;36(1):16-21.
Groell R, Lindbichler F, Riepl T, Gherra L, Roposch A, Fotter R. The reliability of bone age determination in central European children using the Greulich and Pyle method. <i>British Journal of Radiology</i> . 1999;72(857):461-4.
Jain S. Estimation of age from 13 to 21 years. <i>Journal of Forensic Medicine and Toxicology</i> . 1999;16(1):27-30.
Jungmann A. [the Estimation of Age]. <i>Harefuah</i> . 1964;67:178-81.
Kangne RN, Sami SA, Deshpande VL. Age estimation of adolescent girls by radiography. <i>Journal of Forensic Medicine and Toxicology</i> . 1999;16(1):20-6.



Labrune M. [Bone age determination in children]. Soins - Gynecologie, Obstetrique, Puericulture, Pediatrie. 1984;0(36):9-12.
Sabino Prates N, Peters CF, Lopes E. Bone maturity of the hand and wrist and mandibular growth. [Portuguese]. Rgo. 1988;36(5):318-24.
Senguttuvan R, Dahdal DG, Chin C, Desoky S, Wheeler M. Study of concordance in bone age estimation between pediatric endocrinologists vs. Pediatric radiologists. Pediatric Radiology. 2015;45:S227-S8.
Taranger J, Karlberg J, Bruning B, Engstrom I. Standard deviation score charts of skeletal maturity and its velocity in Swedish children assessed by the Tanner-Whitehouse method (TW2-20). Annals of Human Biology. 1987;14(4):357-65.
van Venrooij-Ysselmuizen ME, van Ipenburg A. Mixed longitudinal data on skeletal age from a group of Dutch children living in Utrecht and surroundings. Annals of Human Biology. 1978;5(4):359-80.
Wenzel A, Droschl H, Melsen B. Skeletal maturity in Austrian children assessed by the GP and the TW-2 methods. Annals of Human Biology. 1984;11(2):173-7.
Wenzel A, Melsen B. Skeletal maturity in 6-16-year-old Danish children assessed by the Tanner-Whitehouse-2 method. Annals of Human Biology. 1982;9(3):277-81.
Xiong XX, Li BY, Guo H, Feng ZH. Hand-wrist bone age assessment of Han and Uygur adolescents and children in Urumqi. [Chinese]. Chinese Journal of Tissue Engineering Research. 2015;19(15):2341-5.
Ye YY, Wang CX, Cao LZ. Skeletal maturity of the hand and wrist in Chinese children in Changsha assessed by TW2 method. Annals of Human Biology. 1992;19(4):427-30.
Yogesh S, Bhavesh B, Sanjeev B. A prospective roentgenological study in Mewar region of Rajasthan to establish age group 16 to 18 years. Medico-Legal Update. 2013;13(2):17-20.
Zhen OY, Baolin L. Skeletal maturity of the hand and wrist in Chinese school children in Harbin assessed by the TW2 method. Annals of Human Biology. 1986;13(2):183-7.


Appendix 6: All analysed results



Difference in chronological age and skeletal age (Greulich & Pyle development stages) by chronological age intervals in boys

Figures A1-A6: The difference in chronological age (CA) and skeletal age (SA) (development stages from the Greulich & Pyle atlas) in the included studies for boys 14 to 19 years respectively, grouped according to year of chronological age.

CI: Confidence interval of the mean difference










Difference in chronological age and skeletal age (Greulich & Pyle development stages) by chronological age intervals in girls

Figures A7-A11: The difference in chronological age (CA) and skeletal age (SA) (development stages from the Greulich & Pyle atlas) in the included studies for girls 14 to 18 years respectively, grouped according to year of chronological age.

CI: Confidence interval of the mean difference








Chronological age for boys in the development stages 10 years to 19 years from the Greulich & Pyle atlas


Figures A12-A23: Mean chronological age (CA) in the included studies for boys in development stages from 10 years to 19 years from the Greulich & Pyle atlas.

CI: Confidence interval of the mean difference












Chronological age for girls in the development stages 10 years to 18 years from the Greulich & Pyle atlas

Figures A24-A33: Mean chronological age (CA) in the included studies for girls in development stages from 10 years to 18 years from the Greulich & Pyle atlas.

CI: Confidence interval of the mean difference











Mean chronological age by GP skeletal age categories alongside age distribution of included study participants, boys.



Figures A34-A37: Graphical representation of results for boys from the four studies presenting mean chronological age in GP skeletal age categories. The top part of the chart shows each GP skeletal age category, with the number of observations (left), mean chronological age (middle point) 95% confidence intervals for population means (brackets) and one standard deviation on each side (end of lines). The lower part of the figure shows the number of participants in each year of chronological age.







Mean chronological age by GP skeletal age categories alongside age distribution of included study participants, girls.

Figures A34-A37: Graphical representation of results for girls from the four studies presenting mean chronological age in GP skeletal age categories. The top part of the chart shows each GP skeletal age category, with the number of observations (left), mean chronological age (middle point) 95% confidence intervals for population means (brackets) and one standard deviation on each side (end of lines). The lower part of the figure shows the number of participants in each year of chronological age.





Utgitt av Folkehelseinstituttet
Mars 2017
Postboks 4404 Nydalen
NO-0403 Oslo
Telefon: 21 07 70 00
Rapporten kan lastes ned gratis fra
Folkehelseinstituttets nettsider www.fhi.no