

RESEARCH

Open Access



Does pre-COVID impulsive behaviour predict adherence to hygiene and social distancing measures in youths following the COVID-19 pandemic onset? Evidence from a South African longitudinal study.

Katharina Haag^{1,2}, Stefani Du Toit³, Nace Mikus^{4,5}, Sarah Skeen^{3,6}, Kathryn Steventon Roberts¹, Marguerite Marlow³, Vuyolwethu Notholi³, Akhona Sambudla³, Yeukai Chideya³, Lorraine Sherr¹ and Mark Tomlinson^{3,7*}

Abstract

Background Engagement in protective behaviours relating to the COVID-19 pandemic has been proposed to be key to infection control. This is particularly the case for youths as key drivers of infections. A range of factors influencing adherence have been identified, including impulsivity and risk taking. We assessed the association between pre-COVID impulsivity levels and engagement in preventative measures during the COVID-19 pandemic in a longitudinal South African sample, in order to inform future pandemic planning.

Methods Data were collected from $N=214$ youths (mean age at baseline: $M=17.81$ ($SD=.71$), 55.6% female) living in a South African peri-urban settlement characterised by high poverty and deprivation. Baseline assessments were taken in 2018/19 and the COVID follow-up was conducted in June–October 2020 via remote data collection. Impulsivity was assessed using the Balloon Analogue Task (BART), while hygiene and social distancing behaviours were captured through self-report. Stepwise hierarchical regression analyses were performed to estimate effects of impulsivity on measure adherence.

Results Self-rated engagement in hygiene behaviours was high (67.1–86.1% “most of the time”, except for “coughing/sneezing into one’s elbow” at 33.3%), while engagement in social distancing behaviours varied (22.4–57.8% “most of the time”). Higher impulsivity predicted lower levels of hygiene ($\beta=.14$, $p=.041$) but not social distancing behaviours ($\beta=-.02$, $p=.82$). This association was retained when controlling for a range of demographic and COVID-related factors ($\beta=.14$, $p=.047$) and was slightly reduced when including the effects of a life-skills interventions on hygiene behaviour ($\beta=-.13$, $p=.073$).

*Correspondence:

Mark Tomlinson
markt@sun.ac.za

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusions Our data indicate that impulsivity may predict adolescent engagement in hygiene behaviours post COVID-19 pandemic onset in a high risk, sub-Saharan African setting, albeit with a small effect size. For future pandemics, it is important to understand predictors of engagement, particularly in the context of adversity, where adherence may be challenging. Limitations include a small sample size and potential measure shortcomings.

Keywords COVID, Impulsivity, Adolescence, LMIC, Longitudinal

Background

Following the outbreak of the COVID-19 pandemic, social distancing and hygiene practices were suggested as key measures for limiting infections and deaths, and to prevent health systems from becoming overwhelmed [1]. Recommended social distancing practices included an avoidance of close physical contact to individuals outside the household [2], while hygiene practices comprised hand and respiratory hygiene measures (e.g., hand sanitising, mask wearing). Early evidence from initial lockdowns (approximately March–June 2020) across sub-Saharan Africa indicates moderate to high adherence levels to protective behaviours [3, 4], but also that these decline over time [5]. Factors predicting adherence were demographic (including age, education, poverty, living situation), psychological (perceived COVID-risk and adherence barriers, self-efficacy), and/or COVID-related (e.g., knowledge, dissatisfaction with government measures) in nature [5–7]. However, costs of engagement in such measures may be particularly high for youths, who have a lower risk of severe illness, but for whom social distancing in particular may require forgoing income or social opportunities [8]. In accordance, it was proposed from mid-2020 and onwards that young adults may be key drivers of COVID-19 infections across the globe, including in African settings [9, 10]. Even though the world is reopening, the need to prevent COVID-19 remains. Therefore, it is important to understand predictors of engagement in young people living in Sub-Saharan Africa, particularly in contexts of poverty or high population density, where adherence may be challenging [11, 12]. This information can also be valuable for future pandemic planning.

Impulsive individuals have been shown to struggle with anticipating future consequences of behaviours, and to be more likely to act prematurely, take risks and be easily distracted [13]. Since COVID-related hygiene and social distancing measures commonly require deviations from habitual and automatic responses, impulsive individuals may struggle to maintain them [14, 15]. In HICs, impulsivity, alongside factors such as opportunities to break rules, has been linked to poorer engagement in COVID-related protective behaviours both cross-sectionally [16, 17] and longitudinally [18]. Furthermore, in a Turkish sample, impulsivity was

associated with both lower H1N1- and COVID-related hygiene behaviours [19, 20]. Similarly, higher impulsivity was linked to poorer health-behaviours and higher COVID infection rates during the first Mexican lockdown [14]. There is also a range of indirect evidence, suggesting lower hygiene behaviours in individuals with conditions associated with increased impulsivity, such as attention deficit hyperactivity disorder [21] and anti-social personality disorder [22]. However, there is a lack of longitudinal data and many of the above-mentioned studies use questionnaire rather than behavioural measures of impulsivity and risk-taking. Longitudinal data are valuable as they allow investigating whether population baseline levels of impulsivity/risk-taking predict pandemic behaviours, and whether intervention is warranted for future pandemic planning. Furthermore, previous studies were predominantly conducted in adults, with youths having been proposed to engage more frequently in risky behaviours [23].

Within sub-Saharan Africa, impulsivity has often been studied in the context of youths' health risk behaviours, such as risky sex, gambling, or substance use [24, 25]. To the best of our knowledge, impulsivity has not been explored as a potential risk factor for poor COVID measure compliance but may an important target behaviour in future pandemic planning and readiness. We also had a secondary aim of utilising the current data to derive recommendations for public health communication and implementation of protective measures for future pandemics. We utilised data from a longitudinal study conducted from birth until current ages 19–21 years in an impoverished neighbourhood of Cape Town, South Africa, to gain a first impression of such associations. Before the current COVID-assessments, data were most recently collected following a randomised control trial of a life skills intervention specifically designed for adolescents living in adverse contexts and aimed at teaching youths' skills such as future planning and reflection on their behaviours.

We hypothesized that youths' risk taking in a behavioural task measured before onset of the COVID-19 pandemic (i.e., their "baseline" behaviours) would be associated with lower engagement in hygiene and social distancing behaviours post COVID-19 pandemic onset.

We controlled for and performed secondary analyses on the effects of a pre-COVID adolescent life skills intervention on any relationships found between impulsivity and protective behaviour engagement. With the life skills intervention having previously been found to affect impulsivity in young men in particular, we also expected that it may serve as a mediator of the found associations.

Methods

Study design and setting

The sample was drawn from a longitudinal intervention study conducted in peri-urban Khayelitsha, South Africa, which followed children and their families from before birth until current age (19–21 years; see Fig. 1

for a CONSORT flowchart of assessments). From the antenatal period until 6 months after birth, expectant mothers received either a parenting intervention ('Thula Sana'; $n = 220$) aimed at improving parenting skills and attachment, or maternal services as usual (control group, $n = 229$). All mothers in the community who were eligible for study participation were invited, and group assignment was randomized. Families were followed up several times over the first 18 months of the child's life [26] and again at 13 years of child age [27]. No effects of the early intervention on adolescent outcomes were identified [27] and we subsequently do not control for receipt of this intervention in the current study. The youths then underwent a second intervention ('Zifune'; for details, see




Fig. 1 CONSORT Flow Diagram for Cohort Studies

below), aimed at teaching life skills to improve pro-sociality and reduce violence behaviours, at ages 16–19 years ($n=319$; re-randomized based on early intervention group allocation) (Skeen S, Du Toit S, Marlow M, Stewart J, Rabie S, Melendez-Torrez GJ, et al: Zifune: Does a second wave intervention delivered to former recipients of an early mother-infant attachment intervention reduce interpersonal violence during adolescence? A re-randomized controlled trial, in preparation). Data collection took place in 2018/2019, at three time-points: pre-intervention, directly post-interventions ($n=314$) and at a 3-month follow-up ($n=307$). At the post-intervention assessment, participants completed several behavioural tasks to investigate whether the intervention had led to any changes in risk taking and moral behaviours ($n=280$).

Life skills intervention

The Zifune life skills intervention was developed for use with youths living in high adversity contexts in low- and middle-income countries (LMIC) (Skeen S, Du Toit S, Marlow M, Stewart J, Rabie S, Melendez-Torrez GJ, et al: Zifune: Does a second wave intervention delivered to former recipients of an early mother-infant attachment intervention reduce interpersonal violence during adolescence? A re-randomized controlled trial, in preparation). An adolescent advisory board provided feedback to ensure applicability and acceptability of its contents. The intervention utilises a collaborative approach, incorporates principles of cognitive behaviour therapy, and employs creative and fun methods to allow youths to reflect on their relationships and behaviours and to devise future plans. Eight group-based sessions for groups of approximately 20 youths each were provided by trained facilitators from the local community. Sessions covered six main themes: vision for the future, time management, financial planning, mindfulness, risk-taking behaviour and interpersonal violence, with sessions about long-term planning and risk-taking behaviours in particular potentially affecting impulsivity levels. An intervention facilitator remained in regular contact with and provided support to the youths throughout the course of the study via phone calls.

Data collection during the COVID-19 pandemic

Following the outbreak of the COVID-19 pandemic, South Africa went into a strict lockdown in March 2020. Brief telephonic interviews were conducted with participants ($n=237$) in June to October 2020 through remote-working data collectors. During this time, South Africa's

first large case wave took place (July–August 2020), followed by a strong decline in cases. Participants were assessed on a range of COVID-related variables, including social distancing and hygiene behaviours, household food security, mental health, and schooling outcomes. We utilize data from those who took part in the behavioural tasks at the post-intervention assessment of the Zifune study and completed the COVID-related questionnaire after the pandemic outbreak ($n=214$).

Consent and procedure

All participants provided written consent at each wave of the data collection. Assessments were conducted in the participants' language of choice, predominantly isiXhosa. All data were collected by trained and supervised data collectors, with at least a high school diploma and with prior experience in working with vulnerable populations. For the current phase of the study, ethical approval was obtained from the Health Research Ethics Committee (HREC) from Stellenbosch University (Ref: N17/10/094).

Measures

Demographic variables Information on the gender, age, level of education (utilised in the form of a "correct grade for age" variable), housing (formal vs informal housing); number of household members the individual was living with during the COVID-19 pandemic, HIV status, and household receipt of any form of government-provided cash grants was collected.

Impulsivity/risk taking - balloon analogue risk task (BART) The BART [28] is a naturalistic computer task measuring impulsive and risk-taking behaviours. Participants are presented with a balloon, which they can enlarge in a step-wise fashion by pressing a button. Each pump increases the reward pay-off that the participant receives, but also the chance of the balloon popping, which leads to no rewards for the trial. Participants have the choice to step away after each button press, and collect the already accrued rewards for the trial, or to keep pumping. In the current study, all participants were asked to complete 30 trials. They were told that one trial would be chosen at random in the end, for which they would receive the earned monetary reward. To even out expectations, all participants observed 12 balloons being inflated to their bursting point before commencing the task. The bursting point was set to be identical in each trial between participants. The overall number of pumps was used as a predictor of interest, with a higher number of pumps reflecting higher risk taking.

Hygiene Behaviours The extent to which participants engaged in each of four hygiene behaviours (hand washing, hand sanitising, coughing/sneezing into one’s elbow, and wearing a face mask) during the past week was measured on a scale from 0 “never” to 3 “most of the time” for each item (see Additional file 1: Appendix 1 for full item list and rating scale). A total score (0–12) was calculated. An exploratory factor analysis revealed that the items did not load well onto a single underlying factor, potentially due to participants picking and choosing certain behaviours or adhering less stringently to measures as the pandemic situation in South Africa relaxed towards September/October 2020. As a result, we decided to investigate the total score, reflecting the overall extent of hygiene behaviours each participant engaged in, but also analysed the four behaviours separately to see whether any effects found were driven by high scores on particular items.

Social distancing Behaviours The extent to which participants engaged in five social distancing practices during the past week was assessed: keeping a 1–2 m distance, and avoiding public transport, going to the shops/pharmacy, public spaces and going for a walk in the neighbourhood. Items were rated from 0 “never” to 3 “most of the time” and summed up into a total score (0–15). Exploratory factor analysis suggested that the latter three items loaded onto a potential “avoidance of public outings” factor, though individual item loadings were small. Therefore, we chose to investigate the total score, indexing the extent of overall social distancing behaviours, and to additionally explore single-item effects.

Potential confounders We added age and sex to the analyses, since risk behaviours in the BART have been shown to be influenced by both factors. We furthermore controlled for education (being in the correct grade for age) and timing of the assessment, since the COVID situation changed substantially in South Africa throughout our data collection, from the first case wave in June/July 2020 to level 1 restrictions in September 2020. In terms of COVID-related factors that could have influenced participants’ abilities to engage in hygiene and social distancing behaviours, we adjusted our analyses for household food security as a measure of deprivation (Household Food Insecurity Access Scale (HFIAS, [29])), and the number of individuals living in the participant’s household, which could have desensitized participants to being around large groups of people, or heightened worries and subsequent measure engagement, especially in multi-generational households. Finally, we controlled for receipt of the life skill intervention at ages 16–19 years, as it was found to influence risk taking in males particularly (Mikus N, Skeen, S, Stewart J,

Marlow M, DuToit S, Rabie S, Mendelez Torres GJ, et al: Psychosocial intervention improved self-control in adolescents, in preparation).

Statistical analysis

Analyses were conducted using StataSE 16 and R 4.1.1. In a first step, we investigated descriptive characteristics of the sample and compared it to participants who had completed the BART impulsivity measure and were not included in the COVID follow-up on relevant demographic factors, using t-tests and χ^2 tests as appropriate. We then performed Pearson’s correlation analyses between the key variables. Finally, based on findings from the correlation analyses, a hierarchical linear regression analysis was performed, with hygiene behaviours as the key outcome. In the first step, impulsivity was added as a predictor, with higher pumps on the BART indexing higher impulsivity/ risk taking. Secondly, the demographic factors of age, sex and correct class for age were included. In a third step, COVID-related factors (food security, number of people living in the household, time to level 1 restrictions) were added to the model. In a last step, receipt of the life skills intervention was added, to see whether any effects found may be explained by exposure to its contents. Finally, since the intervention was found to affect BART-measured impulsivity in a previous study (Mikus N, Skeen, S, Stewart J, Marlow M, DuToit S, Rabie S, Mendelez Torres GJ, et al: Psychosocial intervention improved self-control in adolescents, in preparation) and showed close to significant predictions of hygiene behaviours ($\beta = .09$, $p = .190$) in the current study, we conducted secondary exploratory causal mediation analyses, using the “mediation” package in R 4.1.1 [30]. The aim was to investigate whether the life skills intervention may be able

Table 1 Baseline and COVID-19 pandemic sample characteristics (N = 214)

Baseline sample	Statistic
Mean Age in Years	M = 17.81 (SD = .71)
Sex (1 = female)	119 (55.6%)
Living in Informal Housing	21 (9.9%)
HIV Positive	9 (4.6%)
Correct Class for Age	97 (45.5%)
Household Cash Grant Receipt	117 (54.9%)
COVID-19 pandemic sample	
Mean Age in Years	M = 19.53, (SD = .60)
Number of People in Household	M = 3.79, (SD = 1.91)
Food Insecurity Score (0–27)	M = 7.68 (SD = 5.95; range 0–22)

For assessment of baseline characteristics, data from the 3-months follow-up were used, since this was when the BART was completed




Fig. 2 Frequency of individual hygiene and social distancing behaviours in the week before the COVID-19 pandemic follow-up

to buffer potential associations between higher impulsivity and lower protective behaviour engagement. For this, we explored whether any indirect effects of exposure to the life skills intervention on hygiene behaviours through impulsivity would be found. However, we acknowledge limited power due to a small sample size.

Results

Descriptive information

The mean age of the included sample was $M=17.81$ ($SD=.71$; range: 16–20) years at the time of the BART data collection; and $M=19.52$ ($SD=.60$; range 19–21) years at the COVID follow-up. 55.6% ($N=119$) of participants were female. Further descriptive information on the sample is provided in Table 1. The average number of pumps on the BART across the 30 trials was $M=21.13$ ($SD=6.68$; range: 6.82–37.17). Participant responses to

the hygiene and social distancing behaviour items are illustrated in Fig. 2. For hygiene behaviours, the mode answers showed participants adhered to them “most of the time”, except for “coughing and sneezing into one’s elbow”, which only 33.3% of participants engaged in “most of the time” and 40.7% never or rarely engaged in. Twenty-two percent ($n=47$) showed adherence to all four hygiene behaviours “most of the time”, with average total scores lying at $M=10.02$ ($SD=1.55$, range: 6–12). For all social distancing behaviours, the mode was “some of the time”, except for keeping a 1–2-m distance, which a majority of participants complied with “most of the time”. 4.2% ($n=9$) participants engaged in all social distancing behaviours “most of the time” over the past 7 days; the average total score was $M=10.62$ ($SD=2.41$, range: 3–15).

Comparing the sub-samples that were and were not followed up as part of the COVID-19 pandemic assessment

Table 2 Bivariate correlation between impulsivity, hygiene and social distancing behaviours, and potential confounders

	1.	2.	3.	4.	5.	6.	7.	8.	9.
1. Impulsivity (BART pumps)	–								
2. Hygiene Behaviours Total Score	–.14*	–							
3. Social Distancing Behaviour Total Score	–.02	.21*	–						
4. Age	–.03	–.12	.06	–					
5. Sex (1 = female)	–.16*	.04	.22*	.09	–				
6. Correct class for age	–.09	–.05	–.07	.10	–.08	–			
7. Intervention (1 = Received)	–.15*	.11	.11	.04	.20	–.05	–		
8. Time to Level 1 Restrictions	.03	–.13	–.08	.03	.02	–.04	.03	–	
9. Number of Household Members	–.09	.11	–.02	.03	.06	.03	.03	–.08	–
10. COVID Food Security	.06	–.03	–.08	.01	–.02	.12	–.04	.13	.01

(loss to follow-up: 24.5%), we found that females (55.6% versus 44.4%, $\chi^2 = 4.37, p = .037$) and those receiving the life skills intervention (56.1% versus 43.9%, $\chi^2 = 5.62, p = .018$) were more likely to have taken part in the follow-up study. Those followed up had somewhat lower average BART pumps, indexing lower impulsivity ($M_{excluded} = 22.88, SD = 7.67, M_{included} = 21.13, SD = 6.68; p = .007$). No group differences were found in terms of mean age, if the participants lived in formal versus informal housing and were in the correct school class for their age, and whether the household the adolescent lived in received any social grants (all $p > 0.55$).

Correlation analyses between protective behaviours, impulsivity, and potential confounders

Correlation analyses (Table 2) indicated that higher impulsivity (indexed by the number of BART pumps) was negatively associated at a small effect size ($r = -.14, p = .041$) with the extent of hygiene behaviours, but not with social distancing behaviours. It was also negatively linked with being of female sex ($r = -.16, p = .021$) and having received the life skills intervention ($r = -.15, p = .026$). Hygiene and social distancing behaviours were positively correlated at a small effect size ($r = .21, p = .002$). The extent of hygiene behaviours was also marginally negatively associated with age ($r = -.12, p = .082$) and time to level 1 COVID restrictions ($r = -.13, p = .057$) and positively with the number of household members ($r = .11, p = .097$). The extent of social distancing behaviours was positively associated with female sex ($r = .22, p = .001$). Since the hygiene and social distancing items did not load onto the same underlying factors, we also conducted analyses using single items (Additional file 2: Appendix 2). Associations between impulsivity as measured by the BART and individual hygiene behaviours followed the same direction as for the total score for three out of the four hygiene items (except for

Table 3 Step-wise regression model predicting COVID hygiene behaviours during the past 7 days

	B	95% CI	β	p
Step 1				
Impulsivity (BART pumps)	–.03	–.06; –.00	–.14	.041
<i>F, p, adjusted R²</i>	4.21; .041; .015			
Step 2				
Impulsivity (BART pumps)	–.03	–.07; .00	–.15	.035
Sex (1 = female)	.05	–.38; .47	.01	.830
Age	–.30	–.65; .05	–.12	.094
Correct Class for Age	–.16	–.58; .26	–.05	.461
<i>F, p, adjusted R²</i>	2.01; .095; .019			
Step 3				
Impulsivity (BART pumps)	–.03	–.06; .00	–.14	.045
Sex (1 = female)	.02	–.41; .45	.01	.927
Age	–.28	–.63; .07	–.11	.113
Correct Class for Age	–.27	–.71; .16	–.09	.214
Food Insecurity	.00	–.04; .03	.00	.992
Number of Household Members	.09	–.02; .21	.12	.097
Time to Level 1 Restrictions	–.01	–.02; .00	–.13	.066
<i>F, p, adjusted R²</i>	2.12; .043; .036			
Step 4				
Impulsivity (BART pumps)	–.03	–.06; .00	–.13	.073
Sex (1 = female)	.00	–.43; .43	.00	.999
Age	–.29	–.63; .06	–.11	.104
Correct Class for Age	–.27	–.71; .15	–.08	.204
Food Insecurity	.00	–.03; .04	.00	.955
Number of Household Members	.09	–.02; .20	.11	.103
Time to Level 1 Restrictions	–.01	–.02; .00	–.13	.058
Intervention (1 = yes)	.28	–.13; .71	.09	.180
<i>F, p, adjusted R²</i>	2.09; .038; .039			

B Unstandardized regression coefficient, CI Confidence interval, β Standardized regression coefficient

mask-wearing), though at somewhat smaller effect sizes. For the social distancing items, associations were small/non-significant and in mixed directions, reflecting the overall lack of relationship found between impulsivity (BART pumps) and the total social distancing behaviour score. Resultingly, we report the results of multiple regression analysis only with the total hygiene behaviour score as an outcome. However, equivalent analyses for social distancing behaviours are reported in Additional file 3: Appendix 3.

Multivariable models predicting hygiene behaviours

In the first step, higher impulsivity (i.e., a higher number of BART pumps) negatively predicted the total amount of hygiene behaviours at a small effect size ($\beta = -.14$, $p = .045$), explaining 1.5% of their variance (Table 3). This effect was retained once the demographic confounders were included ($\beta = .15$; $p = .035$), though the overall model ($F(4,208) = 2.01$, $p = .095$) was not significant at level $p < .05$. Once COVID-related variables were controlled for, the model regained significance ($F(7,205) = 2.12$, $p = .04$), and impulsivity continued to predict the total hygiene behaviour score at a small effect size ($\beta = .14$, $p = .047$). Since the life skills intervention was by itself found to predict lower impulsivity in males, we examined how the effect of impulsivity on hygiene behaviours changed when including the life skills intervention as a predictor variable. We found that the degree to which impulsivity predicted hygiene behaviours was slightly reduced ($\beta = .13$, $p = .073$), when the receipt of the life skills intervention variable was included in the model.

Mediation analysis for the life skills intervention

This prompted us to explore whether the intervention itself had an effect on hygiene behaviours and whether it was mediated by impulsivity. We found an average causal mediation effect (ACME) of .03 (95% Quasi-Bayesian Confidence Interval (QBCI) = $[-.00, .10]$; $p = .098$), and an average direct effect (ADE) of .18 (95% QBCI = $[-.10, .46]$, $p = .218$). The total estimated effect was .21 (95% QBCI = $[-.06, .49]$, $p = .135$) and the proportion mediated .13 (95% QBCI = $[-.99, 1.46]$, $p = .221$). Effects lay at identical sizes when a moderation by sex was included (ACME = .03, 95% QBCI = $[-.00, .10]$, ADE = .18, 95% QBCI = $[-.10, .46]$).

Discussion

In a South African sample, we found that pre-COVID impulsivity levels predicted hygiene but not social distancing behaviours following the outbreak of the COVID-19 pandemic at a small but significant effect size. This effect was retained when controlling for a range of demographic variables and factors measured post

COVID-19 pandemic onset. When we included the life skills intervention in the model, which aimed to increase prosociality and reduce violence behaviour, the effects of impulsivity were slightly reduced. This suggests that the total effect of impulsivity on hygiene behaviours is partially due to the effects of the intervention on hygiene behaviours. When investigating the effect of the intervention on the main outcome variable, we found some evidence that the life skills intervention itself had a positive (but insignificant) effect on hygiene behaviours, that was mediated by the effects of the intervention on impulsivity with a trend ($p = 0.09$).

This study was not specifically set up to investigate COVID-related changes and was somewhat limited in sample size and resulting analytic power. Thus, findings should not be over-interpreted. However, given that longitudinal data predicting hygiene and social distancing behaviours post COVID-19 pandemic onset and data from LMICs in particular is lacking, it still offers some relevant insights that can be applied to future pandemic planning:

Firstly, in accordance with studies from high income countries (HIC) [15–18], we found that pre-COVID impulsivity predicted lower engagement in hygiene behaviours, though at a small effect size. This supports the notion that challenges such as lack of behavioural inhibition, difficulties to anticipate long-term consequences and altered reward processes may make it more difficult for impulsive individuals to engage in health-related protective behaviours [31, 32]. The BART has been proposed to measure impulsive choice and decision making naturalistically and with a higher objectivity and external validity than questionnaires [28]. While it may be a concern that the BART predominantly examines risk-taking relating to small monetary rewards, it has previously also been linked to health-behaviours, such as smoking and alcohol use [33, 34] and may work similarly for sexual rather than monetary rewards [35], suggesting a degree of generalizability. However, other studies have found relatively low correlations of the BART with multiple facets relating to real-life impulsivity and risk taking [36], which means we may be under-estimating associations. Future studies should aim to investigate more specific underlying processes (e.g., low inhibition versus reward-responsiveness) and potential mediators (e.g., altered risk perceptions) to identify specific behaviours and pathways that can be targeted by interventions. Given the small associations found, it may also be important to investigate specific subgroups that showed high impulsivity during the pandemic and associations of impulsivity/risk-taking during the pandemic with both, baseline impulsivity levels pre-pandemic and measure engagement post pandemic onset.

Second, life skills interventions can reduce impulsivity levels in the general population. The life skills intervention investigated here aimed to teach a range of skills, such as thinking through actions before engaging in them (Skeen S, Du Toit S, Marlow M, Stewart J, Rabie S, Melendez-Torrez GJ, et al: Zifune: Does a second wave intervention delivered to former recipients of an early mother-infant attachment intervention reduce interpersonal violence during adolescence? A re-randomized controlled trial, in preparation) and was found to affect risk taking measures in males in particular (Mikus N, Skeen, S, Stewart J, Marlow M, DuToit S, Rabie S, Mendelez Torres GJ, et al: Psychosocial intervention improved self-control in adolescents, in preparation). Despite not acting as a mediator in our analysis, the trends found suggest that modification of behaviours may be possible and could affect real life outcomes and increase hygiene behaviour adherence. With impulsivity also affecting a range of other key health behaviours [37] and better self-regulation potentially acting as a buffer for healthy development in challenging environments [38] there is a need for public health policies that address its variable levels, ideally through utilising integrative programmes and interventions that focus on its early developmental precursors [38–43].

Third, in accordance with our secondary aim of providing relevant data for future pandemic planning, we found that in a LMIC context characterized by poverty and deprivation, self-reported engagement in hygiene measures was high, but as found in other studies, reduced somewhat as the pandemic situation in South Africa relaxed [5, 44]. With mask wearing being mandated from May 1st, 2020, and the government and NGOs strongly advertising the other three hygiene measures assessed here [2, 45], our study suggests that individuals can both extend and adjust known health behaviours such as hand washing and adopt novel behaviours such as mask wearing if targeted by effective public health messaging. Importantly however, observational evidence indicates that real-life adherence may lie substantially below self-reported levels [46], highlighting potential social desirability and self-selection effects in surveys, and the limitations of self-report. Thus, our study may also be prone to over-reporting engagement. Our results also show that very few individuals consistently engaged in all four recommended hygiene behaviours, mirroring similar findings from South Africa indicating that only 35% of survey respondents followed a high impact set of preventative behaviours [47]. Together, these findings suggest that despite high existing knowledge and awareness of the efficacy of hygiene measures [48], these may ultimately not be adopted consistently. However, part of these findings may also be explained by certain measures being incompatible (e.g., mask wearing and sneezing into

one's elbow), or mask wearing being mandated, which may explain why no associations were found between mask wearing and impulsivity on an individual level.

Of note, self-reported engagement in social distancing behaviours was lower than for hygiene behaviours, potentially due to fewer targeted health messages, different perceived social norms [49], or a lower ability to engage in some of the behaviours (e.g., avoiding going to the shops or using public transport to get to work). Alternatively, it may be that the phrasing of the items (asking for instance how frequently a behaviour was avoided, rather than engaged in) could have affected reported frequencies and thus the validity of the results. Varying compliance with different types of measures has also been established in the Argentinian context [50], with divergent factors predicting engagement in different measures. Overall, there will be a need to better understand and address intersecting vulnerabilities (race, gender, poverty [reflected for instance in high population density, large households, communal water sources], and geopolitics) in sub-Saharan Africa contexts [48, 51, 52] to increase rates of compliance with protective measures in future pandemics.

Fourth, female sex predicted higher engagement in social distancing but not hygiene behaviours. This is of interest, since the former measures were less strongly mandated by official sources and may require higher voluntary engagement. Our findings are in accordance with other studies suggesting lower rates of male engagement in hygiene and social distancing behaviours during the COVID-19 pandemic, but also previous pandemics such as SARS and MERS [53, 54]. One potential explanatory factor is male socialization, which may lead to men being more likely to mask fear, downplay risks and/ or engage in high-risk behaviours [55]. Studies from HIC contexts also suggest that females may have higher individual risk perceptions [56], may be more willing to cooperate with mandated health measures [57] and may be overall more health conscious [58]. Furthermore, while both men and women are employed in keyworker positions, men may be more likely to be the breadwinners for families in the sub-Saharan African context, and thus more likely to struggle to adhere to social distancing (e.g., during transport to work, at the workplace). Future studies will need to further disentangle sex effects, and targeted interventions for males in particular may be required.

Overall, our study highlights the need to apply targeted approaches for both health messaging during a pandemic and future intervention planning. Efforts should be directed towards specific subgroups of individuals that may be at particular risk of not engaging in protective behaviours [54, 55]. Our findings add to a range of

studies investigating potential demographic, psychological, personality and structural predictors [59] that can guide such efforts.

The study has several strengths, including the utilization of longitudinal data from a sub-Saharan African, high-risk context. However, it also has several limitations. Firstly, the measures used to assess hygiene and social distancing behaviours had not previously been validated and did not load onto the same underlying factors. The behaviours assessed by the social distancing items may also have been affected by unassessed secondary factors such as the need to use public transport for work. However, the items still capture a range of key behaviours recommended by governmental and non-governmental agencies. Secondly, given that only a subset of youths had completed the behavioural tasks and that it was challenging to reach all participants remotely, the sample size was somewhat low, limiting our analytical power. Thirdly, the follow-up study was designed to get a rapid overview of COVID-related outcomes and questionnaires were kept brief to prevent drop-out in telephonic interviews. Thus, some constructs potentially relevant to the study such as risk perceptions or being unable to socially distance due to workplace conditions were not assessed, which means effects may be over- or underestimated. Fourth, COVID assessments were collected within 1–1.5 years post-intervention assessment. While substantially different in content, participants could have considered the COVID follow-up as another part of the intervention evaluation. This may have made the intervention group more prone to social desirability, self-selection and continuation effects, potentially leading to higher levels of self-reported engagement in protective behaviours and thus skewing intervention effects.

Conclusion

Overall, our analyses provide a first, tentative indication that impulsivity may predict engagement in hygiene behaviours after the COVID-19 pandemic onset in a high risk, sub-Saharan African setting. Impulsivity therefore may require consideration within future pandemic planning. Interventions addressing behavioural factors may be of benefit and, health messaging may need targeting according to different population needs.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-023-15310-w>.

Additional file 1: Appendix 1. Hygiene and social distancing behaviour items.

Additional file 2: Appendix 2. Correlations between the BART and individual hygiene and social distancing items.

Additional file 3: Appendix 3. Step-wise regression model predicting COVID social distancing behaviours during the past 7 days.

Acknowledgements

The authors are particularly grateful to the families from Khayelitsha who participated in this project, as well as the data collectors involved in this study.

Authors' contributions

KH performed the data analysis, drafted the manuscript and incorporated feedback. SDT was responsible for study management, data collection and data cleaning. NM was responsible for methodology and data analysis. SS was responsible for funding acquisition, methodology, supervision and study management. KR, MM, VN, AS and YC were responsible for data collection, management and cleaning. LS was responsible for supervision and study management. MT was responsible for funding acquisition, conceptualization, supervision and study management. The author(s) read and approved the final manuscript.

Funding

The "Zifune" project was funded by Wellspring Philanthropic Fund. The previous follow up study (age 13 years) was funded by Grand Challenges Canada. The current analyses were also supported through the GCRF "Accelerating Achievement for Africa's Adolescents" hub, through a grant from the UK Economic and Social Research Council. Founders did not have any influence on study design and collection, analysis and interpretation of data, and on writing the manuscript.

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to their sensitive nature but are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Participants aged 18 or older provided informed written consent at each wave of the data collection. For participants younger than 18 years, informed written (for in-person assessments) or verbal (audio-recorded for COVID telephone assessments) consent was obtained from a parent and/or legal guardian and assent was provided by the youths. Assessments were conducted in the participants' language of choice, predominantly isiXhosa. For the current phase of the study, ethical approval was obtained from the Health Research Ethics Committee (HREC) from Stellenbosch University (Ref: N17/10/094). All research was conducted in accordance with the rules and guidelines outlined in the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Institute for Global Health, University College London, London, UK. ²Present affiliation: Department of Child Health and Development, Norwegian Institute for Public Health, Oslo, Norway. ³Institute for Life Course Health Research, Department of Global Health, Stellenbosch University, Cape Town, South Africa. ⁴Department of Cognition, Emotion, and Methods in Psychology, Faculty of Psychology, University of Vienna, Vienna, Austria. ⁵Interacting Minds Centre, Aarhus University, Aarhus, Denmark. ⁶Amsterdam Institute for Social Science Research, Faculty of Social and Behavioural Sciences, University of Amsterdam, Amsterdam, Netherlands. ⁷School of Nursing and Midwifery, Queens University, Belfast, UK.

Received: 8 August 2022 Accepted: 22 February 2023

Published online: 20 March 2023

References

- Nyabadza F, Chirove F, Chukwu CW, Visaya MV. Modelling the potential impact of social distancing on the COVID-19 epidemic in South Africa. Alcaraz R, editor. *Comput Math Methods Med*. 2020;2020:1–12. <https://doi.org/10.1155/2020/5379278>.

2. World Health Organization. Coronavirus disease (COVID-19) advice for the public. 2020. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>. Accessed 20 October 2021.
3. Reddy SP, Sewpaul R, Mabaso M, Parker S, Naidoo I, Jooste S, et al. South Africans' understanding of and response to the COVID-19 outbreak: an online survey. *South Afr Med J*. 2020;110(9):894–902. <https://doi.org/10.7196/SAMJ.2020.v110i9.14838>.
4. Shewasinad Yehualashet S, Asefa KK, Mekonnen AG, Gemedo BN, Shiferaw WS, Aynalem YA, et al. Predictors of adherence to COVID-19 prevention measure among communities in north Shoa zone, Ethiopia based on health belief model: a cross-sectional study. *PLoS One*. 2021;16(1):e0246006. <https://doi.org/10.1371/journal.pone.0246006>.
5. Kollamparambil U, Oyenubi A. Behavioural response to the Covid-19 pandemic in South Africa. Goli S, editor. *PLOS ONE*. 2021;16(4):e0250269. <https://doi.org/10.1371/journal.pone.0250269>.
6. Austrian K, Pinchoff J, Tidwell JB, White C, Abuya T, Kangwana B, et al. COVID-19 related knowledge, attitudes, practices and needs of households in informal settlements in Nairobi, Kenya. *Bull World Health Organ*. 2020. <https://doi.org/10.2471/BLT.20.260281>.
7. Kabamba Nzaji M, Ngoie Mwamba G, Mbidi Miema J, Kilolo Ngoy Umba E, Kangulu IB, Banza Ndala DB, et al. Predictors of non-adherence to public health instructions during the COVID-19 pandemic in the Democratic Republic of the Congo. *J Multidiscip Healthc*. 2020;13:1215–21. <https://doi.org/10.2147/JMDH.S274944>.
8. Nivette A, Ribeaud D, Murray A, Steinhoff A, Bechtiger L, Hepp U, et al. Non-compliance with COVID-19-related public health measures among young adults in Switzerland: insights from a longitudinal cohort study. *Soc Sci Med*. 2021;268:113370. <https://doi.org/10.1016/j.socscimed.2020.113370>.
9. Diop BZ, Ngom M, Pougué Biyong C, Pougué Biyong JN. The relatively young and rural population may limit the spread and severity of COVID-19 in Africa: a modelling study. *BMJ Glob Health*. 2020;5(5):e002699. <https://doi.org/10.1136/bmjgh-2020-002699> Erratum in: *BMJ Glob Health* 2020 Jul;5(7): PMID: 32451367; PMCID: PMC7252974.
10. Zuo C, Meng Z, Zhu F, Zheng Y, Ling Y. Assessing vaccination prioritization strategies for COVID-19 in South Africa based on age-specific compartment model. *Front Public Health*. 2022;10:876551. <https://doi.org/10.3389/fpubh.2022.876551> PMID: 35784231; PMCID: PMC9240634.
11. Gibson L, Rush D. Novel coronavirus in Cape Town informal settlements: feasibility of using informal dwelling outlines to identify high risk areas for COVID-19 transmission from social distancing perspective. (Preprint). *JMIR Public Health Surveill*. 2020;6(2). <https://doi.org/10.2196/18844>.
12. Mueller V, Sherif G, Keeler C, Jehn M. COVID – 19 policy modeling in sub-Saharan Africa. *Appl Econ Perspect Policy*. 2020;43(1):24–38. <https://doi.org/10.1002/aep.13078>.
13. Moeller FG, Barratt ES, Dougherty DM, Schmitz JM, Swann AC. Psychiatric aspects of impulsivity. *Am J Psychiatr*. 2001;158(11):1783–93. <https://doi.org/10.1176/appi.ajp.158.11.1783>.
14. Frías-Armenta M, Corral-Frías NS, Corral-Verdugo V, Lucas MY. Psychological predictors of precautionary behaviors in response to COVID-19: a structural model. *Front Psychol*. 2021;12(12):559289. <https://doi.org/10.3389/fpsyg.2021.559289>.
15. Wismans A, Letina S, Wennberg K, Thurik R, Baptista R, Burke A, et al. The role of impulsivity and delay discounting in student compliance with COVID-19 protective measures. *Personal Individ Differ*. 2021;179:110925. <https://doi.org/10.1016/j.paid.2021.110925>.
16. Byrne KA, Six SG, Anaraky RG, Harris MW, Winterlind EL. Risk-taking unmasked: using risky choice and temporal discounting to explain COVID-19 preventative behaviors. *PLoS One*. 2021;16(5):e0251073. <https://doi.org/10.1371/journal.pone.0251073>.
17. Kuiper ME, de Bruijn AL, Reinders Folmer C, Olthuis E, Brownlee M, Kooistra EB, et al. The intelligent lockdown: compliance with COVID-19 mitigation measures in the Netherlands: Amsterdam Law School Research paper; 2020. <https://doi.org/10.31234/osf.io/5wdb3>.
18. Reinders Folmer CP, Brownlee MA, Fine AD, Kooistra EB, Kuiper ME, Olthuis EH, et al. Social distancing in America: Understanding long-term adherence to COVID-19 mitigation recommendations. Tu W-J, editor. *PLOS ONE*. 2021;16(9):e0257945. <https://doi.org/10.1371/journal.pone.0257945>.
19. Alper S, Bayrak F, Yilmaz O. Psychological correlates of COVID-19 conspiracy beliefs and preventive measures: evidence from Turkey. *Curr Psychol*. 2020;40(11):5708–17. <https://doi.org/10.1007/s12144-020-00903-0>.
20. Gaygisiz Ü, Gaygisiz E, Özkan T, Lajunen T. Individual differences in behavioral reactions to H1N1 during a later stage of the epidemic. *J Infect Public Health*. 2012;5(1):9–21. <https://doi.org/10.1016/j.jiph.2011.09.008>.
21. Merzon E, Manor I, Rotem A, Schneider T, Vinker S, Golan Cohen A, et al. ADHD as a risk factor for infection with Covid-19. *J Atten Disord*. 2021;25(13):1783–90. <https://doi.org/10.1177/1087054720943271>.
22. Miguel FK, Machado GM, Pianowski G, Carvalho LF. Compliance with containment measures to the COVID-19 pandemic over time: do antisocial traits matter? *Pers Individ Dif*. 2021;168:110346. <https://doi.org/10.1016/j.paid.2020.110346> Epub 2020 Aug 21. PMID: 32863507; PMCID: PMC7441860.
23. Romer D. Adolescent risk taking, impulsivity, and brain development: implications for prevention. *Dev Psychobiol*. 2010;52(3):263–76. <https://doi.org/10.1002/dev.20442>.
24. Ssewanyana D, Newton CR, van Baar A, Hassan AS, Stein A, Taylor HG, et al. Beyond their HIV status: the occurrence of multiple health risk behavior among adolescents from a rural setting of sub-Saharan Africa. *Int J Behav Med*. 2020;27(4):426–43. <https://doi.org/10.1007/s12529-020-09877-6>.
25. Ssewanyana D, Bitanihirwe B. Problem gambling among young people in sub-Saharan Africa. *Front Public Health*. 2018;6:23. <https://doi.org/10.3389/fpubh.2018.00023>.
26. Cooper PJ, Tomlinson M, Swartz L, Landman M, Molteno C, Stein A, et al. Improving quality of mother-infant relationship and infant attachment in socioeconomically deprived community in South Africa: randomised controlled trial. *BMJ*. 2009;338:b974. <https://doi.org/10.1136/bmj.b974>.
27. Tomlinson M, Skeen S, Melendez-Torres GJ, Hunt X, Desmond C, Morgan B, et al. First 1,000 days: enough for mothers but not for children? Long-term outcomes of an early intervention on maternal depressed mood and child cognitive development: follow-up of a randomised controlled trial. *J Child Psychol Psychiatry*. 2022;63(3):261–72. <https://doi.org/10.1111/jcpp.13482>.
28. Lejuez CW, Read JP, Kahler CW, Richards JB, Ramsey SE, Stuart GL, et al. Evaluation of a behavioral measure of risk taking: the balloon analogue risk task (BART). *J Exp Psychol Appl*. 2002;8(2):75–84. <https://doi.org/10.1037/1076-898x.8.2.75>.
29. Swindale A, Bilinsky P. Development of a universally applicable household food insecurity measurement tool: process, current status, and outstanding issues. *J Nutr*. 2006;136(5):1449S–52S. <https://doi.org/10.1093/jn/136.5.1449S>.
30. Tingley D, Yamamoto T, Hirose K, Keele L, Imai K. Mediation: RPackage for causal mediation analysis. *J Stat Softw*. 2014;59(5):1–38. <https://doi.org/10.18637/jss.v059.i05>.
31. Evenden JL. (1999). Varieties of impulsivity. *Psychopharmacology*. 1999 Oct;146(4):348–61. <https://doi.org/10.1007/pl0000548>.
32. Stautz K, Pechey R, Couturier D-L, Deary IJ, Marteau TM. Do executive function and impulsivity predict adolescent health behaviour after accounting for intelligence? Findings from the ALSPAC cohort. *PLoS One*. 2016;11(8):e0160512. <https://doi.org/10.1371/journal.pone.0160512>.
33. Canning JR, Schallert MR, Larimer ME. A systematic review of the balloon analogue risk task (BART) in alcohol research. *Alcohol Alcohol*. 2022;57(1):85–103. <https://doi.org/10.1093/alcal/agab004>.
34. Lejuez CW, Aklin WM, Jones HA, Richards JB, Strong DR, Kahler CW, et al. The balloon analogue risk task (BART) differentiates smokers and non-smokers. *Exp Clin Psychopharmacol*. 2003;11(1):26–33. <https://doi.org/10.1037/1064-1297.11.1.26>.
35. Lawyer SR. Risk taking for sexual versus monetary outcomes using the balloon analogue risk task. *Psychol Rec*. 2013;63(4):803–20. <https://doi.org/10.1037/1064-1297.11.1.26>.
36. Frey R, Pedroni A, Mata R, Rieskamp J, Hertwig R. Risk preference shares the psychometric structure of major psychological traits. *Sci Adv*. 2017;3(10):e1701381. <https://doi.org/10.1126/sciadv.1701381>.
37. Ssewanyana D, Mwangala PN, van Baar A, Newton CR, Abubakar A. Health risk behaviour among adolescents living with HIV in sub-Saharan Africa: a systematic review and Meta-analysis. *Biomed Res Int*. 2018;2018:7375831. <https://doi.org/10.1155/2018/7375831>.
38. Haslam D, Mejia A, Thomson D, Betancourt T. Self-regulation in low- and middle-income countries: challenges and future directions. *Clin Child Fam Psychol Rev*. 2019;22(1):104–17. <https://doi.org/10.1007/s10567-019-00278-0> PMID: 30725308.
39. Olson SL, Bates JE, Sandy JM, Schilling EM. Early developmental precursors of impulsive and inattentive behavior: from infancy to middle childhood. *J Child Psychol Psychiatry*. 2002;43(4):435–47. <https://doi.org/10.1111/1469-7610.00035>.

40. Anderson AC, Youssef GJ, Robinson AH, Lubman DI, Verdejo-Garcia A. Cognitive boosting interventions for impulsivity in addiction: a systematic review and meta-analysis of cognitive training, remediation and pharmacological enhancement. *Addiction*. 2021;116(12):3304–19. <https://doi.org/10.1111/add.15469>.
41. Barry MM, Clarke AM, Jenkins R, Patel V. A systematic review of the effectiveness of mental health promotion interventions for young people in low- and middle-income countries. *BMC Public Health*. 2013;13:835. <https://doi.org/10.1186/1471-2458-13-835>.
42. Singla DR, Waqas A, Hamdani SU, Suleman N, Zafar SW, Zill-e-Huma, et al. Implementation and effectiveness of adolescent life skills programs in low- and middle-income countries: a critical review and meta-analysis. *Behav Res Ther*. 2020;130:103402. <https://doi.org/10.1016/j.brat.2019.04.010>.
43. Musci RJ, Bradshaw CP, Maher B, Uhl GR, Kellam SG, Ialongo NS. Reducing aggression and impulsivity through school-based prevention programs: a gene by intervention interaction. *Prev Sci*. 2014;15(6):831–40. <https://doi.org/10.1007/s11121-013-0441-3> Erratum in: *Prev Sci* 2014 Dec;15(6):841. PMID: 24178584; PMCID: PMC4007396.
44. MacIntyre CR, Nguyen P-Y, Chughtai AA, Trent M, Gerber B, Steinhofel K, et al. Mask use, risk-mitigation behaviours and pandemic fatigue during the COVID-19 pandemic in five cities in Australia, the UK and USA: a cross-sectional survey. *Int J Infect Dis*. 2021;106:199–207. <https://doi.org/10.1016/j.ijid.2021.03.056>.
45. National Department of Health (NDoH). COVID-19 Risk Adjusted Strategy. 2020. <https://sacoronavirus.co.za/covid-19-risk-adjusted-strategy/>. Accessed 24 Oct 2021.
46. Jakubowski A, Egger D, Nekesa C, Lowe L, Walker M, Miguel E. Self-reported vs directly observed face mask use in Kenya. *JAMA Netw Open*. 2021;4(7):e2118830. <https://doi.org/10.1001/jamanetworkopen.2021.18830>.
47. Burger R, Christian C, Maughan-Brown B, Rensburg R, Rossouw L. COVID-19 risk Perception, knowledge and behaviour. Cape Town: University of Cape Town; 2020. <https://cramsurvey.org/wp-content/uploads/2020/07/Burger-COVID19-risk-perception-knowledge-and-behaviour-.pdf>. Accessed 16 Oct 2021
48. Gittings L, Toska E, Medley S, Cluver L, Logie CH, Ralayo N, et al. "Now my life is stuck!": experiences of adolescents and young people during COVID-19 lockdown in South Africa. *Glob Public Health*. 2021;16(6):947–63. <https://doi.org/10.1080/17441692.2021.1899262>.
49. Allen J IV, Mahumane A, Riddell J IV, Rosenblat T, Yang D, Yu H. Correcting perceived social distancing norms to combat COVID-19: National Bureau of Economic Research; 2021. <https://doi.org/10.3386/w28651>.
50. Freidin E, Acera Martini L, Senci CM, Duarte C, Carballo F. Field observations and survey evidence to assess predictors of mask wearing across different outdoor activities in an Argentine city during the COVID-19 pandemic. *Appl Psychol Health Well-Being*. 2022;14(1):81–100. <https://doi.org/10.1111/aphw.12292>.
51. Kelley M, Ferrand RA, Muraya K, Chigudu S, Molyneux S, Pai M, et al. An appeal for practical social justice in the COVID-19 global response in low-income and middle-income countries. *Lancet Glob Health*. 2020;8(7):e888–9. [https://doi.org/10.1016/S2214-109X\(20\)30249-7](https://doi.org/10.1016/S2214-109X(20)30249-7).
52. Sewpaul R, Mabaso M, Cloete A, Dukhi N, Naidoo I, Davids AS, et al. Social distancing behaviour: avoidance of physical contact and related determinants among South Africans: twelve days into the COVID-19 lockdown. *Psychol Health Med*. 2022;1–19. <https://doi.org/10.1080/13548506.2022.2075020>.
53. Baker P, White A, Morgan R. Men's health: COVID-19 pandemic highlights need for overdue policy action. *Lancet*. 2020;395:1886–8. [https://doi.org/10.1016/S0140-6736\(20\)31303-9](https://doi.org/10.1016/S0140-6736(20)31303-9).
54. Bish A, Michie S. Demographic and attitudinal determinants of protective behaviours during a pandemic: a review. *Br J Health Psychol*. 2010;15:797–824. <https://doi.org/10.1348/135910710X485826>.
55. Griffith DM, Sharma G, Holliday CS, Enyia OK, Valliere M, Semlow AR, et al. Men and COVID-19: a biopsychosocial approach to understanding sex differences in mortality and recommendations for practice and policy interventions. *Prev Chronic Dis*. 2020;17:200247. <https://doi.org/10.5888/pcd17.200247>.
56. Brug J, Aro AR, Oenema A, de Zwart O, Richardus JH, Bishop GD. SARS Risk Perception, Knowledge, Precautions, and information sources, the Netherlands. *Emerg Infect Dis*. 2004;10(8):1486–9. <https://doi.org/10.3201/eid1008.040283>.
57. Eastwood K, Durrheim D, Francis JL, Tursan d'Espaignet E, Duncan S, Islam F, et al. Knowledge about pandemic influenza and compliance with containment measures among Australians. *Bull World Health Organ*. 2009;87(8):588–94. <https://doi.org/10.2471/blt.08.060772>.
58. Furnham A, Kirkcaldy B. Age and sex differences in health beliefs and Behaviours. *Psychol Rep*. 1997;80(1):63–6. <https://doi.org/10.2466/pr0.1997.80.1.63>.
59. Coetzee BJ, Kagee A. Structural barriers to adhering to health behaviours in the context of the COVID-19 crisis: considerations for low- and middle-income countries. *Glob Public Health*. 2020;15(8):1093–102. <https://doi.org/10.1080/17441692.2020.1779331>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

