

1 **TITLE**

2 Prevalence and Burden of Health Problems in Competitive Adolescent Distance Runners: A
3 6-Month Prospective Cohort Study

4

5 **Corresponding Author:**

6 Alan R. Barker.

7 Address: Children's Health and Exercise Research Centre, Sport and Health Sciences,
8 College of Life and Environmental Sciences, University of Exeter, St Luke's Campus,
9 Exeter, UK, EX1 2LU.

10 Contact Details: +44 (0)1392 722766, A.R.Barker@exeter.ac.uk.

11

12 **CO-AUTHOR LIST**

13 Robert H. Mann., Children's Health and Exercise Research Centre, University of Exeter.

14 Benjamin M. Clarsen., Norwegian School of Sport Sciences & Norwegian Institute of Public
15 Health.

16 Carly D. McKay., Centre for Motivation and Health Behaviour Change, Department for
17 Health, University of Bath.

18 Bryan C. Clift., Department for Health and Centre for Qualitative Research, University of
19 Bath.

20 Craig A. Williams., Children's Health and Exercise Research Centre, University of Exeter.

21

22 **MAIN TEXT WORD COUNT:** 4,739

23

24 **ABSTRACT**

25 **Objectives:** To describe all health problems (injuries and illnesses) in relation to type, location,
26 incidence, prevalence, time loss, severity, and burden, in competitive adolescent distance
27 runners in England. **Design:** Prospective observational study. **Methods:** 136 competitive
28 adolescent distance runners (73 female athletes) self-reported all health problems for 24-weeks
29 between May and October 2019. Athletes self-reported health problems using the Oslo Sports
30 Trauma Research Center Questionnaire on Health Problems. **Results:** The incidence of
31 running-related injury per 1,000 hours of exposure was markedly higher, compared to previous
32 research. At any time, 24% [95% Confidence Intervals (CI):21-26%] of athletes reported a
33 health problem, with 11% [95% CI:9-12%] having experienced a health problem that had
34 substantial negative impact on training and performance. Female athletes reported noticeably
35 more illnesses, compared to male athletes, including higher prevalence, incidence, time loss,
36 and severity. The most burdensome health problems, irrespective of sex, included lower leg,
37 knee, and foot/toes injuries, alongside upper respiratory illnesses. The mean weekly prevalence
38 of time loss was relatively low, regardless of health problem type or sex. **Conclusion:**
39 Competitive adolescent distance runners are likely to be training and competing whilst
40 concurrently experiencing health problems. These findings will support the development of
41 injury and illness prevention measures.

42

43 **WORD COUNT:** 200

44

45 **KEY WORDS:** Athlete Health Protection, Prospective Study, Epidemiology, Injury
46 Prevention, Illness, Performance, Athletics, Track and Field.

47 INTRODUCTION

48 Distance running is one of the most popular sports among children and adolescents around the
49 world.¹ In England, distance running has been reported to be the second most prevalent sport
50 among adolescents.² Although distance running is associated with multiple health benefits in
51 later life,³ adult-based research indicates that participation is also associated with negative
52 health outcomes, such as injury.⁴ In adolescent distance runners, there is a lack of research that
53 has investigated such outcomes.⁵ This population is often included as a sub-group within larger
54 multi-sport samples of adolescent athletes,⁶⁻¹¹ whereby sex differences have been investigated
55 within a heterogenous population rather than at sport-specific levels.^{6 7 9 10} Regardless, in those
56 studies that include adolescent distance runners, the reported running-related injury (RRI)
57 incidence ranges from 0.84 to 17.0 per 1,000 hours of exposure,^{7 8 10 12 13} and injury prevalence
58 ranges from 15 to 32%.^{6 9 11} While these studies used different methodologies,¹⁴ which may
59 account for these differences, data suggest that the most frequently injured anatomical body
60 region is the lower limb, with the knee,^{6 8 10-13 15} lower leg,^{6 8 10 11 15} and ankle^{6 7 12 13 15} being the
61 most commonly affected body areas. However, due to small sample sizes and narrow age
62 ranges, there is limited opportunity to generalise these findings to other distance running
63 populations.

64 Another limitation of the existing distance running literature (adult and paediatric populations)
65 is that numerous epidemiological studies use a time loss or medical attention injury definition,
66 and often do not account for illness within their study design.¹⁶ Therefore, these studies may
67 have underestimated the total number of health problems (injuries and illnesses),^{14 16-18} while
68 ignoring the potential impact of illnesses. For example, injuries that do not result in time loss,
69 and allow athletes to continue to participate regardless of the injury, may be missed. This “loss
70 of detail” is exacerbated when studying adolescent athletes and not examining sex differences
71 within sub-groups of broader sporting populations (i.e., focusing upon track and field athletes,

72 instead of distance runners). This is an important consideration given that the growth and
73 maturation of adolescent athletes differs according to sex.^{19 20} Therefore, any sex differences
74 related to the burden of health problems, defined as the cross-product of severity and
75 incidence,²¹ may require further attention, with the possibility of developing sex- and event-
76 specific injury and illness prevention measures.

77 The purpose of this study was to describe the prevalence, incidence, severity, and burden of
78 health problems within a population of competitive adolescent distance runners in England,
79 using a prospective cohort study design. Specifically, the study aimed to (1) describe all health
80 problems in relation to type, location, incidence, prevalence, time loss, severity, and burden,
81 and (2) describe sex differences related to these outcomes.

82 **METHODS**

83 **Study Design:**

84 This was a 24-week prospective cohort study based on weekly completion of an online
85 questionnaire. Data collection took place between May and October 2019. This timeframe was
86 chosen to reflect the international and domestic outdoor track and field season (approx. April
87 until September) and the start of the cross-country season (approx. October to March).

88 **Participants:**

89 A total of 644 distance runners (athletes) from 210 England Athletics affiliated athletics clubs,
90 aged between 13 and 18 years, were invited to participate in this study. These athletes were
91 selected to take part based on achieving a Top-50 performance in their given age-group during
92 2018, according to the publicly-available *Power of 10* database, for all distance running events
93 from 800 m up to 10,000 m, including the steeplechase.²² The Top-50 performances for each
94 distance running event were collated according to the age-groups used in the *Power of 10*
95 database: 13-14 years (U15), 15-16 years (U17), and 17-19 years (U20). Data extracted from

96 the *Power of 10* database included: event ranking, performance time, name of athlete, year in
97 age-group, name of coach, and name of athletics club. These data were not retained for analysis.
98 Athletes that had achieved a Top-50 performance in their third year as an U20 were excluded
99 due to being over the age of 18 years. Once exported, any duplicate data were identified (i.e.,
100 the same athlete achieving a Top 50 performance for multiple distance running events) and
101 athletes were grouped according to their athletics club affiliation. Once collated, each athletics
102 club was contacted by letter and email with study information and which athletes were eligible
103 to take part. Each athletics club was actively encouraged to share this study information with
104 eligible athletes, their coach, and guardians. If interested, these athletes were able to enrol onto
105 the study by contacting the primary author (RM) via email or telephone. Athletes were
106 excluded from the study if they were injured at the time of study enrolment, not aged between
107 13 and 18 years old, unable to fully understand the study procedure, and/or failed to complete
108 the consent/assent forms and/or baseline questionnaire. Both consent and assent were obtained
109 before an athlete completed the baseline questionnaire. A flow diagram of the recruitment
110 process is presented in Figure 1. Ethics approval was granted by the institutional ethics
111 committee (180801/B/02).

112 Athletes provided data on a rolling basis. During the first 4 weeks, the sample size (n) increased
113 by the following amount: 98 (week 1), 16 (week 2), 19 (week 3), and 3 (week 4). The final
114 study sample consisted of 136 athletes (73 female). Regardless of the athletes' given week of
115 enrolment, data were collected up to week 24. In relation to internal validity, the sex split within
116 this study sample was 54% female, compared with 46% male. Within the total available sample
117 (n = 644), the sex split was 48% female, compared with 52% male.

118 **Data Collection Procedures**

119 **Baseline Questionnaire:**

120 Before starting weekly data collection, each athlete completed a baseline questionnaire via
121 Qualtrics XM (Provo, Utah, USA), an online platform that is compatible with computers and
122 mobile devices. The questionnaire included sections on background demographics (e.g. date of
123 birth), performance history (e.g. event preferences), training practices (e.g. sessions per week),
124 and medical information (e.g. injury history). This questionnaire was based on previous
125 research,^{6 15 23} and developed for a prior study (unpublished). Key stakeholders were involved
126 in the development of this questionnaire to ensure that it was appropriate for the target audience
127 (face validity). This included adolescent distance runners, parents, athletics coaches, and sports
128 physiotherapists (n = 12). Please see supplementary material for a copy of this questionnaire.
129 Participant characteristics were calculated from these questionnaire responses. Chronological
130 age (decimal age) was calculated, before being categorised according to age-group: 13-14 years
131 (U15), 15-16 years (U17), and 17-18 years (U19). Training ages (i.e., number of years
132 participating in distance running); stature (cm), body mass (kg), current performance level (i.e.,
133 club, county, regional, national, or international), and injury history were all self-reported. Each
134 athlete's age at peak height velocity (PHV) was determined by applying sex-specific maturity
135 offset equations,²⁴ and used to estimate maturity timing and tempo.²⁵

136 **Weekly Data Collection:**

137 Injury and illness data were collected using the Oslo Sports Trauma Research Center
138 questionnaire on health problems (OSTRC-H).^{17 26 27} The questionnaire has demonstrated good
139 validity and reliability in samples including runners.²⁶ It consists of four questions about athlete
140 participation in sport, training volume, sports performance, and symptoms of health problems
141 during the previous 7 days.¹⁷ The response to each of these questions is given a value between
142 0 and 25, with 0 (minimum value) representing "no problems" and 25 (maximum value)
143 representing "severe problems". The four values were summed to calculate a severity score
144 from 0 to 100 for each recorded health problem. If the athlete answered all four questions with

145 the minimum value (full participation without health problems, no reduction in training volume
146 or sports performance, and no symptoms), the OSTRC-H was completed for that week. If
147 athletes reported a health problem, they were asked to self-report whether it was an injury or
148 an illness. Athletes were asked to record the anatomical location of all reported injuries, and
149 the main symptoms experienced for all reported illnesses. For all recorded health problems,
150 athletes were asked to record the number of days of complete time loss from training and
151 competition, whether the health problem had previously been recorded, and who the health
152 problem had been reported to (i.e., nobody, medical doctor, or physiotherapist). Athletes were
153 able to report multiple health problems per week. Alongside the OSTRC-H, athletes were also
154 asked to self-report a weekly training diary, having been encouraged to record this throughout
155 the week. Each weekly training diary allowed athletes to detail the type, total duration, distance
156 covered, and rating of perceived exertion related to all of their running-related training sessions
157 or competitions. Athletes also completed the adolescent version of the Profile of Mood States.²⁸
158 The OSTRC-H was sent to athletes on a weekly basis (every Sunday) by email from 5th May
159 until 13th October 2019 (24 weeks) and was completed via Qualtrics XM. If athletes did not
160 complete the questionnaire, email reminders were sent on the following day (Monday), after
161 two days (Wednesday), and after four days (Friday). The athlete's parents or legal guardians
162 were copied into the email reminders after two and four days, respectively. If a response had
163 still not been received after five days (Saturday), the principal investigator would send an SMS
164 reminder to non-responders. If the questionnaire remained unanswered by the time the
165 subsequent weekly questionnaire was distributed, the athlete was categorised as a "non-
166 responder" for that specific week and recorded as missing data.

167 **Definition and Classification of Health Problems**

168 Aligned with recent consensus statements,^{16 29-31} a "broad" definition of health problems was
169 used, recording all health problems regardless of time loss and/or the need for medical

170 attention. Health problems were classified as an injury if they affected the musculoskeletal
171 system and were classified as an illness if they affected a specific organ system or represented
172 general symptoms. Athletes did not classify injuries as having an acute or overuse mechanism.
173 Instead, the primary author (RM) classified injury onset as gradual or sudden. Health problems
174 were defined as “substantial” if they caused moderate or severe reductions in training volume,
175 moderate or severe reductions in performance, or complete inability to participate in distance
176 running, according to the OSTRC-H scoring guide.^{17 26} Health problems were classified as
177 having caused time loss if the injury or illness led to the athlete being unable to participate fully
178 in distance running training and competition the day after the incident occurred.^{16 29}

179 **Prevalence Calculations**

180 The following prevalence measures were calculated on a weekly basis: all health problems,
181 substantial health problems, time loss health problems, all injuries, substantial injuries, time
182 loss injuries, all illnesses, substantial illnesses, and time loss illnesses. The mean prevalence
183 and 95% confidence intervals (CI) were calculated for the entire study period and stratified by
184 sex. To avoid potential overreporting of health problems, each athlete’s first week of data were
185 excluded from analyses.¹⁷

186 **Incidence and Relative Burden of Health Problems**

187 After reviewing each athlete’s questionnaire responses for the entire season, a list of cases was
188 compiled that included the following details: type of health problem, body region and area (for
189 injuries) or main organ system affected (for illnesses), number of weeks reported, cumulative
190 time loss days, and cumulative severity score. To identify the main organ system affected for
191 illnesses, the athletes’ self-reported symptoms were independently reviewed and classified by
192 the first author and a medical doctor, using recommended categories.¹⁶ Once classified,
193 differences were discussed and the main affected organ system was subsequently agreed upon
194 (percentage agreement = 89%). The severity of each case was also based on its cumulative time

195 loss, reported as: *none* (0 days), *slight* (1 day), *minimal* (2-3 days), *mild* (4-7 days), *moderately*
196 *serious* (8-28 days), *serious* (>28 days-6 months), or *long-term* (>6 months).²⁹ The incidence
197 of each type of health problem was expressed as both the number of cases per athlete per year
198 (52 weeks) and per 1,000 hours of exposure. Exposure was calculated from the weekly training
199 diary data.

200 To reflect the relative burden of injuries and illnesses as a proportion of the total health burden,
201 severity scores for each health problem were summed and divided by the cumulative severity
202 score for all health problems.²¹ A risk matrix was created based on the severity and incidence
203 of health problems in all affected injury body areas and illness organ systems, stratified by sex.

204 **Statistical Analysis**

205 For the participant characteristics, the statistical software SPSS (version 26.0; IBM., Chicago,
206 USA) was used to calculate means and standard deviations (SD) for continuous variables. Also,
207 solely in relation to participant characteristics, percentages (%) were calculated for categorical
208 variables, while sex differences were analysed using independent samples t-tests for continuous
209 variables and Chi-squared tests (X^2) for categorical variables. Statistical significance was set
210 at an alpha level of 0.05 and effect sizes (ES) for mean comparisons were described using
211 Cohen's thresholds (small = 0.2, medium = 0.5, large = 0.8).³² For the incidence and prevalence
212 data, the statistical software R was used (version 3.6.1; The R Foundation for Statistical
213 Computing., Vienna, Austria). 95% confidence intervals reported for incidence and prevalence
214 data were used to indirectly infer differences between male and female athletes.

215 **RESULTS**

216 **Response Rate and Participant Characteristics**

217 A total of 136 (73 female) adolescent distance runners participated in this study. Participant
218 characteristics are shown in Table 1. Throughout the study, a total of 2969 questionnaires were
219 distributed, and 2774 responses were received (mean weekly response rate, 91% (range: 85-

220 99%)). During the follow-up period, 97 of the 136 (71%) athletes enrolled in the study
221 completed every weekly questionnaire, while seven athletes dropped out of the study (Figure
222 1). The data collected for these athletes until the time they dropped out were included in the
223 analysis. Responses to the questionnaire were generally received on the Sunday (47%) or
224 Monday (30%) and the median questionnaire completion time was 8 min.

225 **Number, Incidence, and Severity of Health Problems**

226 In total, 136 athletes reported 213 injuries and 150 illnesses. This translated to 4.0 new injuries
227 and 2.8 new illnesses/athlete/year. The incidence for all health problems (both sexes combined)
228 was 42.6 per 1,000 hours (95% CI, 38.4-47.1). The mean time loss was 4 days/athlete/year
229 (95% CI, 3-5 days), with a mean of five days for injuries (95% CI, 3-7 days) and three days for
230 illnesses (95% CI, 2-4 days) (Table 2).

231 The most frequent injury locations were the lower leg (27%), knee (19%), and foot/toes (13%).
232 For illnesses, the most frequently affected organ systems were upper respiratory (65%), lower
233 respiratory (11%), and non-specific illness (10%). The number and severity of injuries (body
234 region and area) and illnesses (organ system) are summarised in Table 3. 61% of injuries had
235 a gradual onset and 39% had a sudden onset. The most frequent injury locations for gradual
236 onset injuries were the lower leg (38%), knee (17%), and thigh (13%). In comparison, the most
237 frequent injury locations for sudden onset injuries were the knee (22%), foot/toes (20%), lower
238 leg (11%), and ankle (11%).

239 **Prevalence of Health Problems**

240 The weekly mean prevalence of all health problems, substantial health problems, and time loss
241 health problems are presented in Table 4. When compared to all health problems, the mean
242 weekly prevalence was reduced for substantial health problems (approx. 50%), and again for
243 time loss health problems (approx. 33%) across the sample.

244 **Burden of Health Problems**

245 Using the total number of time loss days as the basis for injury severity when calculating
246 relative burden (Table 2), injuries represented 80% of the total burden of health problems, with
247 illnesses representing 20%. This was 66% and 34% for female athletes, compared to 85% and
248 15% for male athletes, respectively. Using cumulative severity score as the basis for injury
249 severity (Table 2), injuries represented 70% of the total burden of health problems, with
250 illnesses representing 30%. This was 61% and 39%, and 82% and 18% for female and male
251 athletes, respectively.

252 Figure 2 illustrates the relationship between severity and incidence for the five most commonly
253 affected body areas (injuries) and organ systems (illnesses), stratified by sex, with
254 supplementary data provided for all other health problems.

255 Regardless of sex differences, the body areas representing the highest burden of injuries were
256 the lower leg, knee, and foot/toes. For affected organ systems, the highest burden of illnesses
257 was caused by upper respiratory illness, non-specific illness, and lower respiratory illnesses.

258 **DISCUSSION**

259 To the authors' knowledge, this is the first study to record all injuries and illnesses, including
260 those that did not result in time loss and/or medical attention, exclusively in a population of
261 competitive adolescent distance runners. The key findings were that: 1) the incidence of RRI
262 per 1,000 hours of exposure was markedly higher when compared to previous research; 2) at
263 any time, 24% of athletes reported a health problem, with 11% having experienced a health
264 problem that had substantial negative impact on training and performance; 3) female athletes
265 reported noticeably more illnesses compared with male athletes, including higher prevalence,
266 incidence, time loss, and severity; 4) the most burdensome health problems, regardless of sex,
267 included lower leg, knee, and foot/toes injuries, alongside upper respiratory illnesses; and 5)

268 the mean weekly prevalence of time loss was relatively low, regardless of health problem type
269 or sex.

270 The first key finding was that the incidence of RRI per 1,000 hours of exposure was markedly
271 higher when compared to previous research. For example, the reported RRI per 1,000 hours
272 for all injuries, including male and female athletes, within this study (25.0) was higher than
273 that reported in similar cohorts of adolescent endurance athletes (range: 4.0-13.1), when using
274 a prospective study design.^{8 10 12} These differences remain apparent when sex-specific analyses
275 are made. The data from the present study is also higher than that previously reported in novice
276 adult distance runners.³³ Differences between the aforementioned studies may be explained by
277 the fact that the present study included data from the outdoor Track and Field season, whereby
278 athletes regularly reduced their training volume in order to perform to their best ability in races.
279 Likewise, a period of rest (i.e., training break) was usually taken following athletes' final track
280 race of the season, before transitioning into the cross-country season. When combined, this
281 highlights that the reported exposure may have been lower than if the study had captured data
282 throughout an entire calendar year. Further to this, the use of a broad definition of recordable
283 health problems, capturing 'all health problems,' may inflate the reported incidence per 1,000
284 of exposure.

285 The mean weekly prevalence of all health problems reported within this study (24%) was lower
286 than that reported in cohorts of adolescent endurance athletes (range: 32.7-38%), as part of sub-
287 group analyses in studies that used similar methods.^{9 11} Likewise, the reported mean weekly
288 prevalence of substantial health problems within this study (11%) was lower than that reported
289 in comparable cohorts (range: 17.6-22%).^{9 11} These studies,^{9 11} as well as the current study,
290 demonstrate a pattern that approximately half of all health problems are substantial. When only
291 focussing on injuries, the mean weekly prevalence reported within this study (16%) is both
292 similar to (range: 15-19.4%),^{9 10} and lower than (range: 25.9-32.4%),^{8 11} that reported in similar

293 cohorts of adolescent endurance athletes. For illnesses, the mean weekly prevalence reported
294 within this study (8%) is predominantly lower than that reported in the comparable studies
295 (range: 14-23%),⁸⁻¹⁰ with the exception being a cohort of elite Irish adolescent endurance
296 athletes (6.9%).¹¹ Differences between these studies may be explained by the longer follow-up
297 period (52-weeks) used in two of the studies,^{8 10} thus being representative of a full calendar
298 year, in addition to the possibility that the smaller sample sizes (range: 25 to 76) used in these
299 studies overestimate the prevalence of these health problems.⁸⁻¹¹ The fundamental
300 methodological differences between other studies make any further comparison difficult.

301 The third key finding was that female athletes reported more illnesses (109 illnesses, 73
302 participants), compared to male athletes (41 illnesses, 63 participants). They also reported more
303 injuries (118 injuries, 73 participants) than male athletes (95 injuries, 63 participants) too,
304 although this is a less noticeable difference compared to illnesses. In this study, this resulted in
305 higher prevalence, incidence, time loss, and severity data relating to illnesses in female athletes.
306 In the two available studies that report sex differences specific to adolescent distance runners,⁸
307 ¹¹ this pattern is consistent. However, in studies that combine sport sub-samples when analysing
308 sex-differences,^{9 10} this pattern is not identified. Also, the difference between female and male
309 athletes, in relation to weekly illness prevalence data (8%), is more pronounced in the present
310 study, when compared to others (~3-4%).^{8 11} Nonetheless, this identified sex difference in self-
311 reported illness (and wider health problems) is consistent across general adolescent populations
312 in Europe and North America,³⁴ and elite adult athletes.³¹ When trying to explain this sex
313 difference, it is apparent that female athletes self-report upper and lower respiratory illnesses,
314 and non-specific illnesses, more often than male athletes do. While the data related to
315 respiratory illnesses are contrary to those sex differences reported in non-athletic populations,
316 including adults and adolescents,³⁵ it does align with research in adult endurance athletes.³⁶ In
317 relation to non-specific illnesses, the higher number self-reported by female athletes is difficult

318 to explain without aetiological information, derived from medical diagnoses. Therefore, future
319 research should look to describe and analyse this sex difference according to specific diagnosis
320 and aetiology.¹⁶

321 In relation to the burden of health problems (Figure 2), results were similar regardless of sex.
322 For example, the body region resulting in the greatest burden from injuries was the lower limb,
323 with the greatest burden according to body area being to the lower leg, knee, and foot/toes.
324 Although comparison to previous research is problematic, these reported body areas are largely
325 consistent with previous adult- and adolescent-based research, irrespective of mode of onset.⁴
326 ^{6 8 10-13 15} When combined with the prevalence and incidence data, these results indicate that
327 injury and illness prevention measures for competitive adolescent distance runners should
328 focus on reducing the risk of these specific injuries. Also, as overuse is the usual mode of onset
329 within distance running, any measures should attempt to address this problem. In relation to
330 illnesses, the greatest burden was related to upper respiratory illnesses, in both male and female
331 athletes. While this finding is consistent across the majority of sports,³⁷ the development of
332 prevention measures within this population may also want to consider this illness system. When
333 combined, these findings demonstrate that a holistic approach to injury and illness prevention
334 is required, whereby a range of different prevention strategies may need to be applied.

335 As a pattern identified in the data, the mean weekly prevalence of time loss health problems,
336 regardless of type or sex, was relatively low. For example, the mean weekly prevalence of all
337 health problems was 24%, compared to 4% when employing a time loss definition. This means
338 that a large proportion of self-reported health problems did not cause athletes to miss training
339 and competition. Although this could be interpreted as a positive finding in relation to athlete
340 availability, it also worryingly highlights that competitive adolescent distance runners are
341 likely to be training and competing whilst also experiencing a health problem. The potentially
342 adverse consequences of this practice are concerning, representing a “silent issue” in the sport

343 that is largely overlooked by youth sport consensus statements^{19 38 39} and long-term athlete
344 development models.⁴⁰ However, this finding may be aligned to the nature of endurance sports,
345 whereby athletes are required to sustain consistent and monotonous training intensities,
346 durations, and frequencies,⁴¹ regardless of health problems. Therefore, the potentially negative
347 consequences of training and competing when concurrently experiencing a health problem
348 warrants further investigation, while improved access to medical support at the time of initial
349 injury may act to limit this pattern.

350 **Methodological Considerations**

351 Data collection was reliant on athlete self-report outcomes, without any dedicated support from
352 medical professionals. Although this is normal for adolescent distance running in England, it
353 means that recording specific diagnoses for injuries and illnesses was not possible¹⁶ and, as
354 discussed elsewhere,¹⁷ using an “all health problems” definition can result in overreporting of
355 minor and transient problems (i.e., non-specific symptoms). However, within a homogenous
356 population of distance runners, it is more likely that differences in reporting introduce ‘random
357 noise,’ rather than systematic bias into the results, whereby some athletes may under-report
358 and others may over-report. Nonetheless, to account for the potential issue of over-reporting,
359 the “substantial health problems” definition provides additional information on the full impact
360 of injuries and illnesses in this population of adolescent athletes. Also, injuries were not
361 classified based on their mechanism.¹⁶ However, based on previous studies,^{6 7 9 11 15 38} and the
362 nature of the sport, it is likely that most injuries in this population have a repetitive mechanism,
363 irrespective of whether the onset was sudden or gradual.

364 An additional study limitation is the extent to which these findings are generalisable to more
365 recreational adolescent distance runners and different periods of the calendar year. With the
366 emphasis being on competitive athletes, future studies may wish to focus their attention on the

367 wider population of distance runners, allowing for comparison to these data. Likewise, a longer
368 follow-up period (i.e., one year) may better capture seasonal variations related to the incidence,
369 prevalence, and burden of health problems within this population. As internal validity is a
370 prerequisite for generalisability,⁴² it is also important to highlight that the proportionately low
371 sample size (representing 22% of the total possible sample), coupled with the rolling enrolment
372 of participants, may have unintentionally made the potential for bias greater. However, this
373 form of baseline self-selection resulted in a group of highly motivated participants, evidenced
374 by the high mean weekly response rate (91%) and small number of participants who dropped
375 out of the study (n = 7). This can be upheld as a methodological strength of this study and, in
376 turn, can be seen to decrease selection bias.

377 **Practical Implications**

378 Future injury and illness prevention measures within this population should be aimed at
379 reducing the risk of lower limb injuries, with an emphasis on the lower leg, knee, and foot/toes
380 – supported by previous research.^{6 8 10-13 15} The development of prevention measures should
381 also consider how to address the possibility that adolescent distance runners are training and
382 competing whilst concurrently experiencing health problems, including attempting to improve
383 initial access to medical support. This is important to consider in relation to safeguarding the
384 long-term health and wellbeing of these athletes, whereby excelling as an adolescent athlete is
385 unlikely to be necessary for, nor a guarantee of, success as a senior athlete.⁴³ An additional
386 practical implication is that sex differences in the self-reporting of respiratory and non-specific
387 illnesses should be incorporated into the debate surrounding youth athletic development,¹⁹ with
388 further evidence required to explain this difference. Based on the findings of this study, future
389 descriptive epidemiological studies including adolescent athletes should present data for male
390 and female athletes separately.

391 From a methodological perspective, it is important to reiterate that the response rate during the
392 study was high (91%), with a large proportion of athletes (71%) responding to every weekly
393 questionnaire. Therefore, this study indicates that prospective self-report surveillance methods
394 are feasible in this population, while the questionnaire distribution method can also be
395 advocated for future studies. Finally, the application of the OSTRC-H questionnaire can be
396 recommended, based on its simplicity and capacity to record all health problems.^{17 26} However,
397 future studies should adopt the updated questionnaire,²⁷ include medical diagnoses, and, where
398 appropriate, extend the length of follow-up.

399 **PERSPECTIVES**

400 This study provides an important insight into the extent of health problems within a population
401 of competitive adolescent distance runners. The incidence of RRI per 1,000 hours of exposure
402 was markedly higher when compared to previous research. At any time throughout the follow-
403 up period, 24% of athletes had a health problem, with 11% having a substantial problem with
404 a negative impact on their training and performance. Regardless of sex, lower leg, knee, and
405 foot/toes injuries were the most burdensome health problems, alongside upper respiratory
406 illnesses, which were a particular problem for female athletes. This study also shows that
407 competitive adolescent distance runners are likely to be training and competing whilst
408 concurrently experiencing health problems, whereby initial access to medical support needs to
409 be improved. Therefore, appropriate management strategies for athletes and coaches should be
410 developed (i.e., return-to-play decision making) for when health problems do occur. These data
411 also support the development of holistic injury and illness prevention measures, that should
412 aim to safeguard the long-term health and wellbeing of competitive adolescent distance
413 runners.

414

415 **DECLARATION OF INTEREST STATEMENT**

416 No competing interests (financial or otherwise) declared.

417

418 **DATA AVAILABILITY STATEMENT**

419 Data are available from the corresponding author upon reasonable request.

420

421

422 **REFERENCES**

- 423 1. Hulteen RM, Smith JJ, Morgan PJ, et al. Global participation in sport and leisure-time
424 physical activities: A systematic review and meta-analysis. *Preventive medicine*
425 2017;95:14-25. doi: <https://doi.org/10.1016/j.ypmed.2016.11.027>
- 426 2. Sport England. Active Lives Survey Data Tables London: Sport England; 2019 [Available
427 from: https://www.sportengland.org/know-your-audience/data/active-lives/active-lives-data-tables#children_and_young_people_surveys2020.
428
- 429 3. Pedisic Z, Shrestha N, Kovalchik S, et al. Is running associated with a lower risk of all-
430 cause, cardiovascular and cancer mortality, and is the more the better? A systematic
431 review and meta-analysis. *British Journal of Sports Medicine* 2019;54(15):898-905.
432 doi: 10.1136/bjsports-2018-100493
- 433 4. van Gent RN, Siem D, van Middelkoop M, et al. Incidence and determinants of lower
434 extremity running injuries in long distance runners: A systematic review. *British*
435 *Journal of Sports Medicine* 2007;41(8):469-80. doi: 10.1136/bjism.2006.033548
- 436 5. Steffen K, Engebretsen L. More data needed on injury risk among young elite athletes.
437 *British Journal of Sports Medicine* 2010;44(7):485-89. doi:
438 10.1136/bjism.2010.073833

- 439 6. Jacobsson J, Timpka T, Kowalski J, et al. Prevalence of musculoskeletal injuries in
440 Swedish elite track and field athletes. *American Journal of Sports Medicine*
441 2012;40(1):163-69. doi: 10.1177/0363546511425467 [published Online First:
442 2011/11/05]
- 443 7. Pierpoint LA, Williams CM, Fields SK, et al. Epidemiology of injuries in United States
444 High School track and field: 2008-2009 through 2013-2014. *American Journal of*
445 *Sports Medicine* 2016;44(6):1463-68. doi: 10.1177/0363546516629950 [published
446 Online First: 2016/02/28]
- 447 8. von Rosen P, Floström F, Frohm A, et al. Injury patterns in adolescent elite endurance
448 athletes participating in running, orienteering, and cross-country skiing. *Int J Sports*
449 *Phys Ther* 2017;12(5):822-32.
- 450 9. Moseid CH, Myklebust G, Fagerland MW, et al. The prevalence and severity of health
451 problems in youth elite sports: A 6-month prospective cohort study of 320 athletes.
452 *Scandinavian Journal of Medicine & Science in Sports* 2018;28(4):1412-23. doi:
453 10.1111/sms.13047
- 454 10. von Rosen P, Heijne A, Frohm A, et al. High injury burden in elite adolescent athletes: A
455 52-week prospective study. *Journal of athletic training* 2018;53(3):262-70. doi:
456 10.4085/1062-6050-251-16
- 457 11. Carragher P, Rankin A, Edouard P. A one-season prospective study of illnesses, acute,
458 and overuse injuries in elite youth and junior Track and Field athletes. *Frontiers in*
459 *Sports and Active Living* 2019;1(13) doi: 10.3389/fspor.2019.00013
- 460 12. Rauh MJ, Margherita AJ, Rice SG, et al. High school cross country running injuries: A
461 longitudinal study. *Clinical Journal of Sport Medicine* 2000;10(2):110-16. doi:
462 10.1097/00042752-200004000-00005 [published Online First: 2000/05/08]

- 463 13. Rauh MJ, Koepsell TD, Rivara FP, et al. Epidemiology of musculoskeletal injuries
464 among high school cross-country runners. *American Journal of Epidemiology*
465 2006;163(2):151-59. doi: 10.1093/aje/kwj022 [published Online First: 2005/11/25]
- 466 14. Tabben M, Whiteley R, Wik E, et al. Methods may matter in injury surveillance: “How”
467 may be more important than “what, when or why”. *Biology of sport* 2020;37(1):3-5.
468 doi: 10.5114/biol sport.2020.89935
- 469 15. Huxley DJ, O'Connor D, Healey PA. An examination of the training profiles and injuries
470 in elite youth track and field athletes. *European Journal of Sport Science*
471 2014;14(2):185-92. doi: 10.1080/17461391.2013.809153 [published Online First:
472 2013/06/20]
- 473 16. Bahr R, Clarsen B, Derman W, et al. International Olympic Committee consensus
474 statement: Methods for recording and reporting of epidemiological data on injury and
475 illness in sport 2020 (including STROBE Extension for Sport Injury and Illness
476 Surveillance (STROBE-SIIS)). *British Journal of Sports Medicine* 2020;54(7):372-
477 89. doi: 10.1136/bjsports-2019-101969
- 478 17. Clarsen B, Myklebust G, Bahr R. Development and validation of a new method for the
479 registration of overuse injuries in sports injury epidemiology: The Oslo Sports
480 Trauma Research Centre (OSTRC) overuse injury questionnaire. *British Journal of*
481 *Sports Medicine* 2013;47(8):495-502. doi: 10.1136/bjsports-2012-091524 [published
482 Online First: 2012/10/06]
- 483 18. Clarsen B, Bahr R. Matching the choice of injury/illness definition to study setting,
484 purpose and design: One size does not fit all! *British Journal of Sports Medicine*
485 2014;48(7):510-2. doi: 10.1136/bjsports-2013-093297 [published Online First:
486 2014/03/13]

- 487 19. Bergeron MF, Mountjoy M, Armstrong N, et al. International Olympic Committee
488 consensus statement on youth athletic development. *British Journal of Sports*
489 *Medicine* 2015;49(13):843-51. doi: 10.1136/bjsports-2015-094962
- 490 20. Wik EH, Martinez-Silvan D, Farooq A, et al. Skeletal maturation and growth rates are
491 related to bone and growth plate injuries in adolescent athletics. *Scandinavian Journal*
492 *of Medicine & Science in Sports* 2020;30(5):894-903. doi: 10.1111/sms.13635
493 [published Online First: 2020/02/09]
- 494 21. Bahr R, Clarsen B, Ekstrand J. Why we should focus on the burden of injuries and
495 illnesses, not just their incidence. *British Journal of Sports Medicine*
496 2018;52(16):1018-21. doi: 10.1136/bjsports-2017-098160
- 497 22. Power of 10. Power of 10: Rankings 2020 [Available from:
498 <https://www.thepowerof10.info/rankings/2020>.
- 499 23. Woollings KY, McKay CD, Kang J, et al. Incidence, mechanism and risk factors for
500 injury in youth rock climbers. *British Journal of Sports Medicine* 2015;49(1):44-50.
501 doi: 10.1136/bjsports-2014-094067 [published Online First: 2014/11/12]
- 502 24. Moore SA, McKay HA, Macdonald H, et al. Enhancing a somatic maturity prediction
503 model. *Medicine & Science in Sports & Exercise* 2015;47(8):1755-64. doi:
504 10.1249/mss.0000000000000588 [published Online First: 2014/11/26]
- 505 25. Baxter-Jones ADG, Eisenmann JC, Sherar LB. Controlling for Maturation in Pediatric
506 Exercise Science. *Pediatric Exercise Science* 2005;17(1):18-30. doi:
507 10.1123/pes.17.1.18
- 508 26. Clarsen B, Ronsen O, Myklebust G, et al. The Oslo Sports Trauma Research Center
509 questionnaire on health problems: A new approach to prospective monitoring of
510 illness and injury in elite athletes. *British Journal of Sports Medicine* 2014;48(9):754-
511 60. doi: 10.1136/bjsports-2012-092087 [published Online First: 2013/02/23]

- 512 27. Clarsen B, Bahr R, Myklebust G, et al. Improved reporting of overuse injuries and health
513 problems in sport: an update of the Oslo Sport Trauma Research Center
514 questionnaires. *British Journal of Sports Medicine* 2020;54(7):390-96. doi:
515 10.1136/bjsports-2019-101337
- 516 28. Terry PC, Lane AM, Lane HJ, et al. Development and validation of a mood measure for
517 adolescents. *Journal of Sports Sciences* 1999;17(11):861-72.
- 518 29. Timpka T, Alonso JM, Jacobsson J, et al. Injury and illness definitions and data collection
519 procedures for use in epidemiological studies in Athletics (track and field): Consensus
520 statement. *British Journal of Sports Medicine* 2014;48(7):483-90. doi:
521 10.1136/bjsports-2013-093241 [published Online First: 2014/03/13]
- 522 30. Soligard T, Schweltnus M, Alonso JM, et al. How much is too much? (Part 1)
523 International Olympic Committee consensus statement on load in sport and risk of
524 injury. *British Journal of Sports Medicine* 2016;50(17):1030-41. doi:
525 10.1136/bjsports-2016-096581 [published Online First: 2016/08/19]
- 526 31. Schweltnus M, Soligard T, Alonso JM, et al. How much is too much? (Part 2)
527 International Olympic Committee consensus statement on load in sport and risk of
528 illness. *British Journal of Sports Medicine* 2016;50(17):1043-52. doi:
529 10.1136/bjsports-2016-096572 [published Online First: 2016/08/19]
- 530 32. Cohen J. A power primer. *Psychological Bulletin Journal* 1992;112(1):155-59. doi:
531 10.1037//0033-2909.112.1.155 [published Online First: 1992/07/01]
- 532 33. Videbaek S, Bueno AM, Nielsen RO, et al. Incidence of running-related injuries per 1000
533 h of running in different types of runners: A systematic review and meta-analysis.
534 *Sports Medicine* 2015;45(7):1017-26. doi: 10.1007/s40279-015-0333-8 [published
535 Online First: 2015/05/09]

- 536 34. Torsheim T, Ravens-Sieberer U, Hetland J, et al. Cross-national variation of gender
537 differences in adolescent subjective health in Europe and North America. *Social*
538 *Science & Medicine* 2006;62(4):815-27. doi:
539 <https://doi.org/10.1016/j.socscimed.2005.06.047>
- 540 35. Falagas ME, Mourtzoukou EG, Vardakas KZ. Sex differences in the incidence and
541 severity of respiratory tract infections. *Respiratory medicine* 2007;101(9):1845-63.
542 doi: 10.1016/j.rmed.2007.04.011 [published Online First: 2007/06/05]
- 543 36. He CS, Bishop NC, Handzlik MK, et al. Sex differences in upper respiratory symptoms
544 prevalence and oral-respiratory mucosal immunity in endurance athletes. *Exercise*
545 *Immunology Review* 2014;20:8-22. [published Online First: 2014/07/01]
- 546 37. Walsh NP. Recommendations to maintain immune health in athletes. *European Journal*
547 *of Sport Science* 2018;18(6):820-31. doi: 10.1080/17461391.2018.1449895
- 548 38. DiFiori JP, Benjamin HJ, Brenner JS, et al. Overuse injuries and burnout in youth sports:
549 A position statement from the American Medical Society for Sports Medicine. *British*
550 *Journal of Sports Medicine* 2014;48(4):287-88. doi: 10.1136/bjsports-2013-093299
- 551 39. Mountjoy M, Rhind DJA, Tiiwas A, et al. Safeguarding the child athlete in sport: A
552 review, a framework, and recommendations for the IOC youth athlete development
553 model. *British Journal of Sports Medicine* 2015;49(13):883-86. doi: 10.1136/bjsports-
554 2015-094619
- 555 40. Lloyd RS, Oliver JL. The Youth Physical Development Model: A New Approach to
556 Long-Term Athletic Development. *Strength & Conditioning Journal* 2012;34(3):61-
557 72. doi: 10.1519/SSC.0b013e31825760ea
- 558 41. Seiler S. What is best practice for training intensity and duration distribution in endurance
559 athletes? *International Journal of Sports Physiology and Performance* 2010;5(3):276-
560 91. doi: 10.1123/ijsp.5.3.276 [published Online First: 2010/09/24]

- 561 42. Rothman KJ, Greenland S, Lash TL. Validity in epidemiological studies. In: Rothman KJ,
562 Greenland S, Lash TL, eds. *Modern epidemiology*. 3rd ed. Philadelphia, USA:
563 Lippincott Williams & Wilkins, pp. 128-147 2008:128-47.
- 564 43. Kearney PE, Hayes PR. Excelling at youth level in competitive track and field athletics is
565 not a prerequisite for later success. *Journal of Sport Sciences* 2018;36(21):2502-09.
566 doi: 10.1080/02640414.2018.1465724

567

DRAFT

APPENDICES

Appendix 1: Baseline Questionnaire

This questionnaire has been resubmitted as a PDF, as explained in the response to reviewers.

Appendix 2: Supplementary Data

A supplementary file has been provided for access to original data used to create the risk matrix, excluding means and 95% confidence intervals for health problems with less than three cases.

DRAFT

TABLES

Table 1. Participant characteristics (data presented as mean and SD, unless otherwise stated)

Characteristic	Overall (n = 136)	Female athletes (n = 73)	Male athletes (n = 63)	p-Value	Effect Size
Chronological age, years	15.9 (1.3)	15.8 (1.3)	16.1 (1.2)	0.15	0.25
Training age, years	5.2 (2.1)	5.6 (2.1)	4.8 (1.9)	0.04	0.36
Age-group (n, %):				X ² = 0.67	
13-14 years	26 (19%)	19 (26%)	7 (11%)		
15-16 years	72 (53%)	37 (51%)	35 (56%)		
17-18 years	38 (28%)	17 (23%)	21 (33%)		
Stature, cm	171.0 (8.7)	166.1 (6.8)	176.6 (7.1)	<0.01	1.52
Body mass, kg	54.3 (9.1)	50.2 (6.9)	59.0 (9.1)	<0.01	1.10
Maturity timing (n, %)				X ² = 0.08	
Pre-PHV	0 (0%)	0 (0%)	0 (0%)		
At-PHV	7 (5%)	6 (8%)	1 (2%)		
Post-PHV	129 (95%)	67 (92%)	62 (98%)		
Maturity tempo (n, %)				X ² = 0.26	
Early	1 (1%)	1 (1%)	0 (0%)		
Average	128 (94%)	70 (96%)	58 (92%)		
Late	7 (5%)	2 (3%)	5 (8%)		
Injury <12 months				X ² = 0.24	
Yes	100 (74%)	57 (78%)	43 (68%)		
No	36 (27%)	16 (22%)	20 (32%)		
Current performance level (n, %):				X ² = 0.98	
Club	10 (7%)	6 (8%)	4 (6%)		
County	43 (32%)	22 (30%)	21 (33%)		
Regional	16 (12%)	9 (12%)	7 (11%)		
National	60 (44%)	32 (44%)	28 (44%)		
International	7 (5%)	4 (6%)	3 (5%)		

Abbreviations: n, number; cm, centimetres; kg, kilograms; PHV, peak height velocity.

NB: Due to rounding, not all numbers add up to stated N.

Table 2. Incidence, total time loss, and cumulative severity score of all health problems, all injuries, and all illnesses (split by sex)

	Incidence				Total time loss (d)	Cumulative severity score (AU)
	Cases/athlete/year	95% CI	Cases/1,000 hours of exposure	95% CI		
All health problems (n = 363)	6.8	6.13-7.53	42.6	38.4-47.1	1433	30218
Female athletes (n = 227)	4.3	3.7-4.8	50.5	44.1-57.5	813	17623
Male athletes (n = 136)	2.5	2.1-3.0	33.8	28.3-40.0	620	12595
All Injuries (n = 213)	4.0	3.5-4.6	25.0	21.8-28.6	1058	21121
Female athletes (n = 118)	4.0	3.3-4.8	26.2	21.7-31.4	533	10785
Male athletes (n = 95)	4.0	3.2-4.9	23.6	19.1-28.9	525	10336
All Illnesses (n = 150)	2.8	2.4-3.3	17.6	14.9-20.7	375	9097
Female athletes (n = 109)	3.7	3.0-4.4	24.2	19.9-29.2	280	6838
Male athletes (n = 41)	1.7	1.3-2.3	10.2	7.3-13.8	95	2259

Abbreviations: d, days; AU, arbitrary unit; %, percentage; CI, confidence interval; n, number.

Table 3. Severity of time loss of all health problems, all injuries (body region and area), and all illnesses (organ system).

Classification	Cases (number)										
	Body region Body area / organ system	Female athletes					Male athletes				
		0 days	1-7 days	8-28 days	>28 days	Total Time Loss (days)	0 days	1-7 days	8-28 days	>28 days	Total Time Loss (days)
All health problems	98	157	11	7	810	50	73	13	6	623	
All injuries	49	52	5	6	530	36	50	9	6	528	
Lower limb	41	45	5	5	490	31	41	8	6	488	
Foot/toes	2	9	2	1	132	5	4	3	1	84	
Ankle	5	2	0	0	6	4	5	2	1	76	
Lower leg	11	12	1	3	224	10	18	1	2	172	
Knee	12	12	1	1	92	5	7	0	2	116	
Thigh	6	8	0	0	20	4	5	1	0	27	
Hip/groin	5	2	1	0	16	3	2	1	0	13	
Trunk	6	7	0	1	40	4	6	0	0	18	
Abdomen	0	1	0	0	2	0	0	0	0	0	
Lumbosacral	4	4	0	1	36	2	6	0	0	18	
Thoracic spine	2	0	0	0	0	0	0	0	0	0	
Chest	0	2	0	0	2	2	0	0	0	0	
Upper limb	2	0	0	0	0	1	3	1	0	22	
Wrist	0	0	0	0	0	0	1	0	0	4	
Elbow	1	0	0	0	0	0	0	0	0	0	
Shoulder	1	0	0	0	0	1	2	1	0	18	
All illnesses	49	53	6	1	280	14	23	4	0	95	
Upper respiratory	33	28	6	0	142	9	18	4	0	80	
Lower respiratory	8	5	0	0	9	2	1	0	0	2	
Gastrointestinal	2	4	0	0	7	0	1	0	0	2	
Neurological	1	1	0	0	1	0	0	0	0	0	
Psychological	0	2	0	0	9	1	0	0	0	0	
Dermatological	0	0	0	0	0	0	1	0	0	7	
Non-specific illness	4	9	0	1	101	0	1	0	0	1	
Energy, load management and nutrition	1	4	0	0	11	2	1	0	0	3	

Table 4. Weekly prevalence of all health problems, substantial health problems, and time loss health problems (in percentages)

	All		Female athletes		Male athletes	
	Mean	95% CI	Mean	95% CI	Mean	95% CI
All health problems	24	21-26	27	24-30	20	16-23
All injuries	16	14-18	16	15-16	16	13-18
All illnesses	8	7-10	12	9-14	4	3-6
Substantial health problems	11	9-12	10	9-12	11	9-13
All injuries	7	6-9	6	5-7	9	7-11
All illnesses	4	3-4	4	3-6	2	1-3
Time loss health problems	4	3-4	3	3-4	4	3-5
All injuries	3	2-4	3	2-4	3	2-5
All illnesses	0	0-1	1	0-1	0	0-1

Abbreviations: %, percentage; CI, confidence interval.

FIGURES

Figure 1





Figure 2



DRAFT

FIGURE CAPTIONS

Figure 1. Study flow chart illustrating participant recruitment, enrolment, and dropout. N.B. Due to the nature of data collection, it is not possible to confirm whether all 443 athletes received study information. Only the athletics clubs confirmed receipt of this information.

Figure 2. Risk matrices illustrating the relationship between severity (consequence) and incidence (likelihood) of all injuries (areas) and illnesses (systems) with three or more reported cases in a population of competitive adolescent distance runners, stratified by sex. The five most commonly affected health problems are labelled. Shading illustrates the relative importance of each health problem; the darker the colour, the greater the overall burden, and the greater the priority should be given to prevention. A supplementary file can be downloaded for access to original data, excluding means and 95% confidence intervals for health problems with less than three cases.