

Supplementary Table. Hazard ratios (95% confidence interval) for the association between mortality and self-reported smoking status in 51 417 men and women

	Smoking status			
	Non-smokers	Current smokers		
		<10 cigarettes/day	10-19 cigarettes/day	20+ cigarettes/day
N	38 555	3707	5137	4018
Total mortality				
No. of deaths	2542	210	361	399
HR (95 % CI)	1 (Ref.)	1.93 (1.67-2.22)	2.12 (1.89-2.37)	2.47 (2.21-2.75)
Cardiovascular disease mortality				
No. of deaths	837	61	111	94
HR (95 % CI)	1 (Ref.)	1.77 (1.36-2.30)	2.19 (1.79-2.68)	1.95 (1.56-2.42)
Cancer mortality				
No. of deaths	841	66	113	153
HR (95 % CI)	1 (Ref.)	1.70 (1.32-2.19)	1.76 (1.44-2.16)	2.49 (2.08-2.99)
Smoking-related cancer mortality				
No. of deaths	317	36	64	99
HR (95 % CI)	1 (Ref.)	2.52 (1.78-3.57)	2.70 (2.05-3.56)	4.22 (3.32-5.36)

Hazard ratios are adjusted for sex, age, survey, social class, longstanding illness and alcohol intake. The non-smoker comprises never and former smokers