



Cochrane
Library

Cochrane Database of Systematic Reviews

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Palmer MJ, Henschke N, Bergman H, Villanueva G, Maayan N, Tamrat T, Mehl GL, Glenton C, Lewin S, Fønhus MS, Free C

Palmer MJ, Henschke N, Bergman H, Villanueva G, Maayan N, Tamrat T, Mehl GL, Glenton C, Lewin S, Fønhus MS, Free C.
Targeted client communication via mobile devices for improving maternal, neonatal, and child health.
Cochrane Database of Systematic Reviews 2020, Issue 8. Art. No.: CD013679.
DOI: [10.1002/14651858.CD013679](https://doi.org/10.1002/14651858.CD013679).

www.cochranelibrary.com

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Copyright © 2020 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	3
SUMMARY OF FINDINGS	5
BACKGROUND	19
OBJECTIVES	20
METHODS	20
Figure 1.	25
RESULTS	28
Figure 2.	31
DISCUSSION	40
AUTHORS' CONCLUSIONS	42
ACKNOWLEDGEMENTS	44
REFERENCES	45
CHARACTERISTICS OF STUDIES	55
DATA AND ANALYSES	125
Analysis 1.1. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 1: Health behaviour change – exclusive breastfeeding in short term (up to 3 months)	128
Analysis 1.2. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 2: Health behaviour change – receiving postpartum help with breastfeeding (3 months postpartum)	128
Analysis 1.3. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 3: Health behaviour change – taking iron and folate tablets during pregnancy	128
Analysis 1.4. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 4: Health behaviour change – contraceptive use (3 months postpartum)	129
Analysis 1.5. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 5: Health behaviour change – smoked in the last 30 days	129
Analysis 1.6. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 6: Health behaviour change – smoking cessation (objectively verified continuous abstinence) (36 weeks' gestation)	129
Analysis 1.7. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 7: Health behaviour change – no alcohol consumption during pregnancy	130
Analysis 1.8. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 8: Service utilisation – attendance at ≥ 4 antenatal care appointments	130
Analysis 1.9. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 9: Service utilisation – attendance for antenatal vaccination	130
Analysis 1.10. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 10: Service utilisation – attendance at antenatal preventive treatment for malaria	131
Analysis 1.11. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 11: Service utilisation – skilled attendant at birth	131
Analysis 1.12. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 12: Service utilisation – newborn postpartum care	131
Analysis 1.13. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 13: Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)	132
Analysis 1.14. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 14: Service utilisation – attendance for newborn vaccination	132
Analysis 1.15. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 15: Health status and well-being – maternal mortality and morbidity	133
Analysis 1.16. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 16: Health status and well-being – maternal mortality and morbidity	133
Analysis 1.17. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 17: Health status and well-being – neonatal mortality and morbidity	134
Analysis 1.18. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 18: Health status and well-being – neonatal mortality and morbidity	134

Analysis 1.19. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 19: Health status and well-being – neonatal health	135
Analysis 1.20. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 20: Health status and well-being – preterm birth	135
Analysis 1.21. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 21: Sensitivity analysis (cluster-RCTs: health status and well-being – preterm birth	135
Analysis 2.1. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 1: Health behaviour change – exclusive breastfeeding (9 weeks postpartum)	136
Analysis 2.2. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 2: Health behaviour change – contraceptive use (9 weeks postpartum)	137
Analysis 2.3. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 3: Service utilisation – newborn postpartum care (10 days after delivery)	137
Analysis 2.4. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 4: Service utilisation – attendance for newborn vaccination	137
Analysis 2.5. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 5: Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)	138
Analysis 2.6. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 6: Health status and well-being – maternal mortality and morbidity	138
Analysis 2.7. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 7: Health status and well-being – neonatal mortality and morbidity	138
Analysis 3.1. Comparison 3: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant and postpartum women), Outcome 1: Service utilisation – attendance for antenatal influenza vaccination	139
Analysis 3.2. Comparison 3: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant and postpartum women), Outcome 2: Service utilisation – birth at health facility	139
Analysis 4.1. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 1: Health behaviour change – mother taking any type of antiretroviral (ARV) (34–36 weeks' gestation)	140
Analysis 4.2. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 2: Health behaviour change – mother taking any type of ARV (6–8 weeks postpartum)	141
Analysis 4.3. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 3: Health behaviour change – infant ARV/prevention of mother-to-child transmission treatment adherence (6 weeks postpartum)	141
Analysis 4.4. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 4: Health behaviour change – infant HIV tested (6–8 weeks postpartum)	141
Analysis 4.5. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 5: Service utilisation – postnatal care: attendance at postpartum care appointment (6–8 weeks postpartum) ...	142
Analysis 4.6. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 6: Service utilisation – intrapartum care: birth in health facility	142
Analysis 4.7. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 7: Service utilisation – antenatal care: mean number of face-to-face or mobile communications with healthcare workers	142
Analysis 4.8. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 8: Health status and well-being – neonatal health: neonatal death/stillbirth	143
Analysis 4.9. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 9: Health status and well-being – neonatal health: infant HIV test positive (6–8 weeks postpartum)	143
Analysis 5.1. Comparison 5: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant women with HIV), Outcome 1: Health behaviour – infant antiretroviral/prevention of mother-to-child transmission adherence (6 weeks after delivery)	144
Analysis 5.2. Comparison 5: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant women with HIV), Outcome 2: Service utilisation – postnatal care: attendance at postpartum care appointment (10 weeks postpartum)	144

Analysis 6.1. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 1: Attendance for necessary healthcare	145
Analysis 6.2. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 2: Timeliness of vaccination	145
Analysis 6.3. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 3: Service utilisation – no emergency department attendance (6 months)	146
Analysis 7.1. Comparison 7: Digital targeted client communication (TCC) compared to non-digital targeted client communication (parents of children aged under five years), Outcome 1: Health behaviour change – oral health in children (Visible Plaque Index, [0–100%], low = good)	146
Analysis 7.2. Comparison 7: Digital targeted client communication (TCC) compared to non-digital targeted client communication (parents of children aged under five years), Outcome 2: Service utilisation – attendance for vaccinations at 14 weeks	147
Analysis 8.1. Comparison 8: Digital targeted client communication (TCC) compared to digital non-targeted client communication (parents of children aged under five years), Outcome 1: Service utilisation – attendance for vaccinations at 7 months	147
ADDITIONAL TABLES	147
APPENDICES	156
HISTORY	174
CONTRIBUTIONS OF AUTHORS	174
DECLARATIONS OF INTEREST	175
SOURCES OF SUPPORT	175
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	175
NOTES	176
INDEX TERMS	176

[Intervention Review]

Targeted client communication via mobile devices for improving maternal, neonatal, and child health

Melissa J Palmer¹, Nicholas Henschke², Hanna Bergman², Gemma Villanueva², Nicola Maayan³, Tigest Tamrat⁴, Garrett L Mehl⁴, Claire Glenton⁵, Simon Lewin^{5,6}, Marita S Fønhus⁵, Caroline Free¹

¹Department of Population Health, London School of Hygiene and Tropical Medicine, London, UK. ²Cochrane Response, Cochrane, London, UK. ³Independent consultant, London, UK. ⁴Department of Sexual and Reproductive Health, World Health Organization, Geneva, Switzerland. ⁵Norwegian Institute of Public Health, Oslo, Norway. ⁶Health Systems Research Unit, South African Medical Research Council, Cape Town, South Africa

Contact address: Melissa J Palmer, Melissa.palmer@lshtm.ac.uk.

Editorial group: Cochrane Consumers and Communication Group.

Publication status and date: New, published in Issue 8, 2020.

Citation: Palmer MJ, Henschke N, Bergman H, Villanueva G, Maayan N, Tamrat T, Mehl GL, Glenton C, Lewin S, Fønhus MS, Free C. Targeted client communication via mobile devices for improving maternal, neonatal, and child health. *Cochrane Database of Systematic Reviews* 2020, Issue 8. Art. No.: CD013679. DOI: [10.1002/14651858.CD013679](https://doi.org/10.1002/14651858.CD013679).

Copyright © 2020 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution-Non-Commercial Licence](https://creativecommons.org/licenses/by-nc/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

ABSTRACT

Background

The global burden of poor maternal, neonatal, and child health (MNCH) accounts for more than a quarter of healthy years of life lost worldwide. Targeted client communication (TCC) via mobile devices (MD) (TCCMD) may be a useful strategy to improve MNCH.

Objectives

To assess the effects of TCC via MD on health behaviour, service use, health, and well-being for MNCH.

Search methods

In July/August 2017, we searched five databases including The Cochrane Central Register of Controlled Trials, MEDLINE and Embase. We also searched two trial registries. A search update was carried out in July 2019 and potentially relevant studies are awaiting classification.

Selection criteria

We included randomised controlled trials that assessed TCC via MD to improve MNCH behaviour, service use, health, and well-being. Eligible comparators were usual care/no intervention, non-digital TCC, and digital non-targeted client communication.

Data collection and analysis

We used standard methodological procedures recommended by Cochrane, although data extraction and risk of bias assessments were carried out by one person only and cross-checked by a second.

Main results

We included 27 trials (17,463 participants). Trial populations were: pregnant and postpartum women (11 trials conducted in low-, middle- or high-income countries (LMHIC); pregnant and postpartum women living with HIV (three trials carried out in one lower middle-income country); and parents of children under the age of five years (13 trials conducted in LMHIC). Most interventions (18) were delivered via text

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Copyright © 2020 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

messages alone, one was delivered through voice calls only, and the rest were delivered through combinations of different communication channels, such as multimedia messages and voice calls.

Pregnant and postpartum women

TCCMD versus standard care

For behaviours, TCCMD may increase exclusive breastfeeding in settings where rates of exclusive breastfeeding are less common (risk ratio (RR) 1.30, 95% confidence intervals (CI) 1.06 to 1.59; low-certainty evidence), but have little or no effect in settings where almost all women breastfeed (low-certainty evidence). For use of health services, TCCMD may increase antenatal appointment attendance (odds ratio (OR) 1.54, 95% CI 0.80 to 2.96; low-certainty evidence); however, the CI encompasses both benefit and harm. The intervention may increase skilled attendants at birth in settings where a lack of skilled attendants at birth is common (though this differed by urban/rural residence), but may make no difference in settings where almost all women already have a skilled attendant at birth (OR 1.00, 95% CI 0.34 to 2.94; low-certainty evidence). There were uncertain effects on maternal and neonatal mortality and morbidity because the certainty of the evidence was assessed as very low.

TCCMD versus non-digital TCC (e.g. pamphlets)

TCCMD may have little or no effect on exclusive breastfeeding (RR 0.92, 95% CI 0.79 to 1.07; low-certainty evidence). TCCMD may reduce 'any maternal health problem' (RR 0.19, 95% CI 0.04 to 0.79) and 'any newborn health problem' (RR 0.52, 95% CI 0.25 to 1.06) reported up to 10 days postpartum (low-certainty evidence), though the CI for the latter includes benefit and harm. The effect on health service use is unknown due to a lack of studies.

TCCMD versus digital non-targeted communication

No studies reported behavioural, health, or well-being outcomes for this comparison. For use of health services, there are uncertain effects for the presence of a skilled attendant at birth due to very low-certainty evidence, and the intervention may make little or no difference to attendance for antenatal influenza vaccination (RR 1.05, 95% CI 0.71 to 1.58), though the CI encompasses both benefit and harm (low-certainty evidence).

Pregnant and postpartum women living with HIV

TCCMD versus standard care

For behaviours, TCCMD may make little or no difference to maternal and infant adherence to antiretroviral (ARV) therapy (low-certainty evidence). For health service use, TCC mobile telephone reminders may increase use of antenatal care slightly (mean difference (MD) 1.5, 95% CI -0.36 to 3.36; low-certainty evidence). The effect on the proportion of births occurring in a health facility is uncertain due to very low-certainty evidence. For health and well-being outcomes, there was an uncertain intervention effect on neonatal death or stillbirth, and infant HIV due to very low-certainty evidence. No studies reported on maternal mortality or morbidity.

TCCMD versus non-digital TCC

The effect is unknown due to lack of studies reporting this comparison.

TCCMD versus digital non-targeted communication

TCCMD may increase infant ARV/prevention of mother-to-child transmission treatment adherence (RR 1.26, 95% CI 1.07 to 1.48; low-certainty evidence). The effect on other outcomes is unknown due to lack of studies.

Parents of children aged less than five years

No studies reported on correct treatment, nutritional, or health outcomes.

TCCMD versus standard care

Based on 10 trials, TCCMD may modestly increase health service use (vaccinations and HIV care) (RR 1.21, 95% CI 1.08 to 1.34; low-certainty evidence); however, the effect estimates varied widely between studies.

TCCMD versus non-digital TCC

TCCMD may increase attendance for vaccinations (RR 1.13, 95% CI 1.00 to 1.28; low-certainty evidence), and may make little or no difference to oral hygiene practices (low-certainty evidence).

TCCMD versus digital non-targeted communication

TCCMD may reduce attendance for vaccinations, but the CI encompasses both benefit and harm (RR 0.63, 95% CI 0.33 to 1.20; low-certainty evidence).

No trials in any population reported data on unintended consequences.

Authors' conclusions

The effect of TCCMD for most outcomes is uncertain. There may be improvements for some outcomes using targeted communication but these findings were of low certainty. High-quality, adequately powered trials and cost-effectiveness analyses are required to reliably ascertain the effects and relative benefits of TCCMD. Future studies should measure potential unintended consequences, such as partner violence or breaches of confidentiality.

PLAIN LANGUAGE SUMMARY

Communicating to pregnant woman and parents through their mobile devices to improve maternal, neonatal, and child health

Aim of this review

We assessed the effect of sending targeted messages by mobile devices to pregnant women and parents of young children about health and healthcare services.

Key messages

There are gaps in the evidence regarding the effects of targeted messages by mobile devices to pregnant women and parents of young children about health and healthcare services. Some of these messages may improve some people's health and their use of health services, but others may make little or no difference. The existing evidence is mostly of low or very low certainty.

What was studied in the review?

Targeted client communication (TCC) is an intervention in which the health system sends information to particular people, based on their health status or other factors specific to that population group. Common types of TCC are text messages reminding people to attend appointments or that offer healthcare information and support. Our review assessed whether TCC can change pregnant women's and parents' behaviour, health service use, health, and well-being.

What happens when pregnant women receive targeted messages by mobile device?

Compared to women who get no messages

Women may breastfeed more in settings where exclusive breastfeeding is not common. They may also go to more antenatal care appointments. They may use skilled birth attendants more where this is less common. We do not know if the messages affect women's or babies' health because the certainty of the evidence is very low.

Compared to women who get messages sent in other ways

Women and newborns may have fewer health problems during the first 10 days after birth. The messages may make little or no difference to the number of women who breastfeed. We do not know if they make women use more health services.

Compared to women who get untargeted messages

The messages may make little or no difference to whether women get influenza vaccines during pregnancy. We do not know if the messages affect women's or babies' health or lead women to use skilled birth attendants more because the evidence is lacking or of very low certainty.

What happens when pregnant women living with HIV receive targeted messages by mobile device?

Compared to women who get no messages

Women may go to slightly more antenatal care appointments. We do not know whether the messages lead more women to give birth in a health facility or improve babies' health because the evidence is of very low certainty. The messages may make little or no difference to whether pregnant women and babies follow antiretroviral (ARV) treatment (used to treat HIV) according to plan. We do not know if the messages affect women's health because the evidence is missing.

Compared to women who get messages sent in other ways

We do not know what the effect of these messages is because we lack evidence.

Compared to women who get untargeted messages

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

More parents may follow their babies' ARV treatment according to plan. We do not know if the messages improve women's or babies' health or their use of services because the evidence is missing.

What happens when parents of young children receive targeted messages by mobile device?

Compared to parents who get no messages

More parents may take their children to healthcare services such as vaccination appointments. But we do not know if the messages improve children's health or their health behaviour because the evidence is missing.

Compared to parents who get messages sent in other ways

Slightly more parents may take their children to vaccination appointments. The messages may make little or no difference to children's toothbrushing habits. We do not know if the messages affect children's health because the evidence is missing.

Compared to parents who get untargeted messages

Fewer parents may take their children to vaccination appointments, but this evidence is mixed. We do not know if the messages affect children's health due to lack of evidence.

How up-to-date is this review?

We searched for studies published up to August 2017. We carried out a search update in July 2019 and relevant studies are reported in the 'Characteristics of studies awaiting classification' section.

SUMMARY OF FINDINGS

Summary of findings 1. Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women
Setting: community and healthcare settings
Intervention: digital targeted client communication via mobile devices
Comparison: standard care or no intervention

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with standard care	Risk with digital targeted client communication				
Health status and well-being – maternal mortality and morbidity Follow-up: up to 3 months	4 RCTs reported on 5 morbidity outcomes and the CIs for most outcomes encompassed both benefit and harm: severe obstetric complications up to 6 weeks postpartum (RR 0.86, 95% CI 0.70 to 1.07); 'any maternal health problem' up to 10 days postpartum (RR 0.50, 95% CI 0.90 to 2.76); breast pain up to 3 months postpartum (RR 0.28, 95% CI 0.09 to 0.80); breast engorgement up to 3 months postpartum (RR 0.58, 95% CI 0.31 to 1.10); number of acute maternal episodes requiring a clinic visit up to 3 months postpartum (MD 0.06, 95% CI -0.19 to 0.31). 1 study also reported on maternal mortality (RR 2.86, 95% CI 0.30 to 27.40), but this study only recorded 5 events.			3073 (4 RCTs)	⊕⊕⊕⊕ Very low a,b,c	We are uncertain of the effect of the intervention on maternal mortality and morbidity because the certainty of the evidence was very low.
Health status and well-being – neonatal mortality and morbidity Follow-up: up to 3 months	The results of pooled analysis of 3 RCTs for neonatal morbidity and mortality combined was consistent with no effect but CIs encompassed benefit and harm (OR 1.00, 95% CI 0.61 to 1.64). 1 RCT reported that the intervention reduced number of acute neonatal episodes requiring a clinic visit up to 3 months postpartum (MD -0.53, 95% CI -0.92 to -0.14).			3005 (4 RCTs)	⊕⊕⊕⊕ Very low a,b,c	We are uncertain of the effect of the intervention on neonatal mortality and morbidity because the certainty of the evidence was very low.
Health behaviour change: breastfeeding – exclusive breastfeeding	Low-risk setting			40 (1 RCT)	⊕⊕⊕⊕ Low a,d	In settings where 100% of women report exclusive breastfeeding, the intervention may make little or no difference to exclusive breastfeeding (though an increase would not be possible). The CI encompasses no effect and harm.
	1000 per 1000	920 per 1000 (790 to 1000)	RR 0.92 (0.79 to 1.08)			

Follow-up: up to 3 months	Moderate-risk setting					The intervention may increase short-term exclusive breastfeeding in settings where rates of exclusive breastfeeding are lower.
	667 per 1000	867 per 1000 (707 to 1000)	RR 1.30 (1.06 to 1.59)	135 (1 RCT)		
Service utilisation – attendance at ≥ 4 antenatal care appointments	311 per 1000	410 per 1000 (265 to 572)	OR 1.54 (0.80 to 2.96)	2550 (1 RCT)	⊕⊕⊕⊕ Low a,b	The intervention may increase attendance for antenatal care appointments, but the CI includes both an increase and a decrease.
Follow-up: during antenatal period						
Service utilisation: intrapartum care – skilled attendant at birth	Low-risk setting				⊕⊕⊕⊕ Low a,b	The intervention may make little or no difference to the proportion of births occurring with skilled attendance in settings where skilled attendance at is common.
	988 per 1000	988 per 1000 (978 to 998)	OR 1.00 (0.34 to 2.94)	1743 (1 RCT)		
	High-risk setting					
Follow-up: at delivery	Results were presented separately for urban and rural populations. The study reported intervention benefit in urban populations (cluster-adjusted OR 4.45, 95% CI 1.36 to 14.51), but not in rural populations (cluster adjusted OR 0.83, 95% CI 0.36 to 1.92)			2550 (1 RCT)		In settings where skilled attendance at birth is less common, the intervention may increase the proportion of births occurring with skilled attendance in urban areas. The intervention may slightly reduce skilled attendance in rural settings, but the CI encompasses both benefit and harm.
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence identified.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **MD:** mean difference; **OR:** odds ratio; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for risk of bias: all studies had unclear sequence generation or allocation concealment, or both.

^bDowngraded one level for imprecision: 95% confidence intervals that encompass both harmful and beneficial effects of the intervention.

^cDowngraded one level due to inconsistency: effect estimates vary in terms of direction and magnitude of effect.
^dDowngraded one level for imprecision: few events.

Summary of findings 2. Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women
Setting: community and healthcare settings
Intervention: digital targeted client communication via mobile devices
Comparison: non-digital targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with non-digital targeted client communication	Risk with digital targeted client communication				
Health status and well-being – maternal mortality and morbidity ('any maternal health problem' up to 10 days postpartum)	333 per 1000	63 per 1000 (13 to 263)	RR 0.19 (0.04 to 0.79)	59 (1 RCT)	⊕⊕⊕⊖ Low a,b	The intervention may reduce the proportion of women reporting a maternal health problem up to 10 days after birth.
Health status and well-being – neonatal mortality and morbidity ('any newborn health problem' up to 10 days postpartum)	481 per 1000	250 per 1000 (120 to 510)	RR 0.52 (0.25 to 1.06)	59 (1 RCT)	⊕⊕⊕⊖ Low a,b	The intervention may reduce the proportion of women reporting a newborn health problem up to 10 days after birth. But the CI includes both an increase and a decrease in reporting.
Health behaviour change: breastfeeding (exclusive breastfeeding at 9 weeks)	1000 per 1000	920 per 1000 (790 to 1000)	RR 0.92 (0.79 to 1.07)	42 (1 RCT)	⊕⊕⊕⊖ Low a,b	The intervention may make little or no difference to the proportion who breastfeed exclusively (though an increase would not be possible due to 100% breastfeeding in the control arm). The CI encompasses no beneficial effect and harm. It is possible the effect of the intervention may be different in populations with different levels of baseline risk.

Service utilisation – attendance antenatal care appointments	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on attendance for antenatal care is unknown as there was no direct evidence identified.
Service utilisation: intrapartum care – skilled attendant at birth	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on intrapartum care is unknown as there was no direct evidence identified.
Unintended consequences	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence identified.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for risk of bias: unclear risk of bias for allocation concealment and incomplete outcome data.

^bDowngraded one level for imprecision: few events.

Summary of findings 3. Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women

Setting: community and healthcare settings

Intervention: digital targeted client communication via mobile devices

Comparison: digital non-targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with digital non-targeted	Risk with digital targeted client communication				

	communi- cation					
Health status and well-being – maternal mortality and morbidity	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on maternal morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – neonatal mortality and morbidity	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on neonatal morbidity and mortality is unknown as there was no direct evidence.
Health behaviour change: breastfeeding	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on breastfeeding is unknown as there was no direct evidence.
Service utilisation – attendance antenatal care appointments (attendance for antenatal influenza vaccination)	310 per 1000	326 per 100 (220 to 490)	RR 1.05 (0.71 to 1.58)	204 (1 RCT)	⊕⊕○○ Low a,b	The intervention may make little or no difference to attendance for antenatal influenza vaccination, but the CI includes both an increase and a decrease in attendance.
Service utilisation: intrapartum care – skilled attendant at birth	875 per 1000	875 per 1000 (604 to 1000)	RR 1.00 (0.69 to 1.45)	16 (1 RCT)	⊕○○○ Very low b,c	We are uncertain of the effect of the intervention on the proportion of women having a skilled attendant at birth because the certainty of the evidence was very low.
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for imprecision: 95% confidence intervals that encompass a potential harmful effect and a potential beneficial effect of the intervention.

^bDowngraded one level for risk of bias: trial at unclear risk of bias for several domains.

^cDowngraded two levels for risk of bias: trial at unclear or high risk of bias across all domains.

Summary of findings 4. Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women with HIV
Setting: community and healthcare settings
Intervention: digital targeted client communication via mobile devices
Comparison: standard care or no intervention

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with standard care	Risk with digital targeted client communication				
Health status and well-being: maternal morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on maternal morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – neonatal morbidity and mortality combined Follow-up: up to 4 weeks	32 per 1000	36 per 1000 (12 to 105)	RR 1.12 (0.39 to 3.28)	381 (1 RCT)	⊕⊕⊕⊕ Low ^a	The intervention may make little or no difference to neonatal mortality and morbidity. However, the CI includes both an increase and a decrease in neonatal mortality and morbidity.
Service utilisation – antenatal care: communications with HCWs Follow-up: during antenatal period	Mean number of face-to-face or mobile communications with HCW for antenatal care was 6	Mean number of face-to-face or mobile communications with HCW for antenatal care was 7.5 (5.64 to 9.36)	MD 1.50 (–0.36 to 3.36)	297 (1 RCT)	⊕⊕⊕⊕ Low ^b	The intervention may slightly increase attendance for antenatal care appointments.
Service utilisation – intrapartum care: birth in health facility Follow-up: at delivery	600 per 1000	510 per 1000 (372 to 690)	RR 0.85 (0.62 to 1.15)	134 (1 RCT)	⊕⊕⊕⊕ Very low b,c	We are uncertain of the effect of the intervention on the proportion of women giving birth in a health facility because certainty of the evidence was very low.
Health status and well-being – neonatal health: infant HIV test positive	13 per 1000	7 per 1000 (1 to 33)	RR 0.54 (0.11 to 2.56)	852 (2 RCTs)	⊕⊕⊕⊕ Very low d,e	We are uncertain of the effect of the intervention on the proportion of infants testing positive for HIV because the certainty of the evidence was very low.

Follow-up: 8 weeks postpartum					
Health behaviour change: adherence to ARV therapy	1 study reported maternal antenatal ARV usage (RR 1.04, 95% CI 0.91 to 1.19), maternal postnatal ARV usage (RR 0.87, 95% CI 0.61 to 1.24), and infant ARV/PMTCT treatment adherence at 6–8 weeks postpartum (RR 1.01, 95% CI 0.98 to 1.04).	503 (1 RCT)	⊕⊕○○ Low ^b		The intervention may make little or no difference to maternal and infant ARV treatment adherence. However, the CI includes benefit and harm.
Follow-up: up to 6–8 weeks postpartum					
Unintended consequences	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

ARV: antiretroviral; **CI:** confidence interval; **HCW:** healthcare worker; **MD:** mean difference; **PMTCT:** prevention of mother-to-child transmission; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded two levels for imprecision: few events and a 95% confidence intervals that encompass a potential large harmful effect and a potential large beneficial effect of the intervention.

^bDowngraded two levels for risk of bias: more women were newly diagnosed with HIV in the control arm (55% with intervention versus 66% with control; $P = 0.015$), randomisation procedures and allocation concealment were not described, lack of blinding of participants, and only per-protocol analysis reported with unexplained dropouts (Kassaye 2016).

^cDowngraded one level for imprecision: 95% confidence intervals that encompass a harmful effect and a potential beneficial effect of the intervention.

^dDowngraded one level for risk of bias: for one trial randomisation procedures and allocation concealment were not described, lack of blinding of participants, and only per-protocol analysis reported with unexplained dropouts.

^eDowngraded two levels for imprecision: few events and a 95% confidence intervals that encompass a potential large harmful effects and a potential large beneficial effect of the intervention.

Summary of findings 5. Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women with HIV

Setting: community and healthcare settings

Intervention: digital targeted client communication via mobile devices

Comparison: non-digital targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with non-digital targeted client communication	Risk with digital targeted client communication				
Health status and well-being: maternal morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on maternal morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – neonatal morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on neonatal morbidity mortality is unknown as there was no direct evidence.
Service utilisation – antenatal care	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on attendance for antenatal care is unknown as there was no direct evidence.
Service utilisation – intrapartum care	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on intrapartum care is unknown as there was no direct evidence.
Health status and well-being – neonatal health: infant HIV status	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on the proportion of infants testing positive for HIV is unknown as there was no direct evidence.
Health behaviour change: adherence to ARV therapy	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on adherence to ARV therapy is unknown as there was no direct evidence.
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

ARV: antiretroviral; **CI:** confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

Summary of findings 6. Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant women and postpartum with HIV) for improving maternal, neonatal, and child health

Patient or population: pregnant and postpartum women with HIV
Setting: community and healthcare settings
Intervention: digital targeted client communication via mobile devices
Comparison: digital non-targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with digital non-targeted client communication	Risk with digital targeted client communication				
Health status and well-being –maternal morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on maternal morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – neonatal morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on neonatal morbidity mortality is unknown as there was no direct evidence.
Service utilisation – antenatal care	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on attendance for antenatal care is unknown as there was no direct evidence.
Service utilisation – intrapartum care	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on intrapartum care is unknown as there was no direct evidence.
Health status and well-being – neonatal health: infant HIV status	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on the proportion of infants testing positive for HIV is unknown as there was no direct evidence.

Health behaviour change: adherence to ARV therapy	720 per 1000	907 per 1000 (770 to 1000)	RR 1.26 (1.07 to 1.48)	150 (1 RCT)	⊕⊕○○ Low ^a	The intervention may increase infant ARV adherence.
Follow-up: 6 weeks postpartum						
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

ARV: antiretroviral; **CI:** confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice for risk of bias: trial at high or unclear risk of bias across all applicable domains.

Summary of findings 7. Digital targeted client communication via mobile devices compared to standard care or no intervention (parents of children aged under five years) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to standard care or no intervention (parents of children aged < 5 years) for improving maternal, neonatal, and child health

Patient or population: parents of children aged < 5 years

Setting: community and healthcare settings

Intervention: digital targeted client communication via mobile devices

Comparison: standard care or no intervention

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with standard care	Risk with digital targeted client communication				

Health status and well-being – child morbidity and mortality combined	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on child morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – child nutritional status	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on child nutritional status is unknown as there was no direct evidence.
Health behaviour change – breastfeeding	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on breastfeeding is unknown as there was no direct evidence.
Service utilisation – attendance for necessary healthcare (attendance for vaccinations at 6–12 months, attendance at HIV medical appointments)	642 per 1000 777 per 1000 (693 to 860)	RR 1.21 (1.08 to 1.34)	5660 (10 RCTs)	⊕⊕⊕⊕ Low ^{a,b}	The intervention may increase attendance for necessary healthcare. However, the result varied considerably according to whether the healthcare attendance was for vaccinations at 6 months, vaccinations at 12 months, or an HIV medical appointment, and between studies within each of these outcome categories.
Follow-up: up to 12 months					
Health behaviour change – hygiene practices	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on hygiene practices is unknown as there was no direct evidence.
Health behaviour change – correct treatment taken	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on taking correct treatment is unknown as there was no direct evidence.
Unintended consequences	No studies reported this outcome.	—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for risk of bias: most studies at unclear risk of bias for allocation concealment.

^bDowngraded one level for inconsistency: high statistical heterogeneity ($I^2 > 90\%$).

Summary of findings 8. Digital targeted client communication via mobile devices compared to non-digital targeted client communication (parents of children aged under five years) for improving maternal, neonatal, and child health
Digital targeted client communication via mobile devices compared to non-digital targeted client communication (parents of children aged < 5 years) for improving maternal, neonatal, and child health

Patient or population: parents of children aged < 5 years
Setting: community and healthcare settings
Intervention: digital targeted client communication via mobile devices
Comparison: non-digital targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with non-digital targeted client communication	Risk with digital targeted client communication				
Health status and well-being – child morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on child morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – child nutritional status	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on child nutritional status is unknown as there was no direct evidence.
Health behaviour change – breastfeeding	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on breastfeeding is unknown as there was no direct evidence.
Service utilisation – attendance for necessary healthcare (attendance for vaccinations at 14 weeks)	839 per 1000	948 per 1000 (839 to 1000)	RR 1.13 (1.00 to 1.28)	744 (1 RCT)	⊕⊕○○ Low ^a	The intervention may slightly increase attendance for vaccinations. However, the CI includes both no increase and a large increase in attendance.
Health behaviour change – hygiene practices (oral health in children at 4 weeks, Visible Plaque Index, 0–100%, lower score is better)	Mean score: 35.6	Mean score: 33.5 (28.06 to 38.94)	MD –2.10 (–7.54 to 3.34)	143 (1 RCT)	⊕⊕○○ Low ^{b,c}	The intervention may make little or no difference to oral hygiene practices. However, the CI includes both benefit and harm.
Health behaviour change – correct treatment taken	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on taking correct treatment is unknown as there was no direct evidence.
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **MD:** mean difference; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded two levels for risk of bias: study at unclear or high risk of bias across all but one domain.

^bDowngraded one level for imprecision: confidence interval encompasses both benefit and harm.

^cDowngraded one level for risk of bias: study at unclear risk of bias for sequence generation and allocation concealment.

Summary of findings 9. Digital targeted client communication via mobile devices compared to digital non-targeted client communication (parents of children aged under five years) for improving maternal, neonatal, and child health

Digital targeted client communication via mobile devices compared to digital non-targeted client communication (parents of children aged < 5 years) for improving maternal, neonatal, and child health

Patient or population: parents of children aged < 5 years

Setting: community and healthcare settings

Intervention: digital targeted client communication via mobile devices

Comparison: digital non-targeted client communication

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with digital non-targeted client communication	Risk with digital targeted client communication				
Health status and well-being – child morbidity and mortality combined	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on child morbidity and mortality is unknown as there was no direct evidence.
Health status and well-being – child nutritional status	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on child nutritional status is unknown as there was no direct evidence.

Health behaviour – breastfeeding	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on breastfeeding is unknown as there was no direct evidence.
Service utilisation – attendance for necessary healthcare – attendance for vaccinations at 6 months	652 per 1000	411 per 1000 (215 to 782)	RR 0.63 (0.33 to 1.20)	40 (1 RCT)	⊕⊕⊕⊕ Low a,b	The intervention may reduce attendance for vaccinations, but the CI includes both an increase and a decrease in attendance.
Follow-up: 6 months						
Health behaviour change – hygiene practices	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on hygiene practices is unknown as there was no direct evidence.
Health behaviour change – correct treatment taken	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on taking correct treatment is unknown as there was no direct evidence.
Unintended consequences	No studies reported this outcome.		—	(0 studies)	—	The effect of the intervention on unintended consequences is unknown as there was no direct evidence.

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for risk of bias: unclear risk of bias for allocation concealment, high risk of bias for incomplete outcome reporting and other bias.

^bDowngraded one level for imprecision: small number of events and confidence interval encompassing potential harmful effect and potential beneficial effect of the intervention.

BACKGROUND

Description of the condition

The enormous burden of disease due to poor sexual, reproductive, maternal, newborn, child, and adolescent health (SRMNCAH) renders them urgent global health priorities. In 2016, poor maternal and neonatal health, and communicable and nutritional diseases, which particularly adversely affect children, accounted for more than a quarter of healthy years of life lost worldwide (Hay 2016). Neonatal preterm birth complications, HIV/AIDS, neonatal encephalopathy due to birth asphyxia and trauma, lower respiratory infections, and diarrhoeal diseases were among the 10 leading causes of total years of life lost, with the burden heavily concentrated in children under five years of age (Naghavi 2017). Those living in low- and middle-income countries are disproportionately affected by poor SRMNCAH (Black 2016).

In 2015, an estimated 303,000 women died during and following pregnancy and childbirth, and maternal mortality remains a leading cause of death for adolescent women. The vast majority of these deaths occurred in low-resource settings, and most could have been prevented (Alkema 2016). Almost three-quarters of maternal deaths are due to direct obstetric causes, such as haemorrhage, hypertensive disorders, sepsis, abortion, embolism, and complications of labour (Say 2014). The majority of indirect causes of maternal deaths are due to the exacerbation of pre-existing conditions; with HIV accounting for an estimated 5.5% of global maternal deaths (Say 2014). Intrinsically linked to maternal health, 1.7 million stillbirths occurred in 2016 worldwide, with key causes including pregnancy and childbirth complications, lifestyle factors, diabetes and hypertension, maternal infections, preterm birth, and birth defects (Naghavi 2017; WHO 2017). Furthermore, five million children under the age of five years died in 2016, with almost half of these deaths occurring among newborns. The three leading global causes of death in children aged under five years were lower respiratory infections, neonatal preterm birth complications, and neonatal encephalopathy due to birth asphyxia and trauma (Naghavi 2017). Of those who survive, one-third of children fail to reach their full physical, cognitive, psychological, or socioemotional potential (or a combination of these) as a result of poverty, poor health and nutrition, insufficient care and stimulation, and other risk factors of importance to early childhood development (Every Women Every Child 2015).

Indicative of the continued global commitment to the survival and well-being of women and children, the UN Secretary General's Global Strategy for Women's and Children's Health was launched in 2010. In 2015, this was recast as the Global Strategy for Women's, Children's and Adolescents' Health aligning its priorities with the ambitious targets relating to the improvement of sexual, reproductive, maternal, newborn, and child health which feature in the Sustainable Development Goals (UN 2015). In 2011, the World Health Organization (WHO) published its global review of RMNCAH interventions, *Essential Interventions, Commodities and Guidelines for Reproductive, Maternal, New-born and Child Health*, with the aim of developing consensus on the content of packages of interventions to address the main causes of maternal, newborn, and child deaths (PMNCH 2011). The health issues targeted by the recommended interventions span adolescence and prepregnancy health (e.g. prevention and management of sexually transmitted infections (STIs) including HIV, family planning, and preconception care); pregnancy (e.g. provision of safe abortion

and postabortion care; appropriate antenatal care (ANC) including screening for maternal illness, preventive dietary supplements, and immunisations; prevention of pre-eclampsia); childbirth (e.g. medical interventions such as caesarean section where indicated); maternal postnatal care (e.g. detection and management of sepsis; family planning); newborn postnatal care (e.g. early initiation of exclusive breastfeeding; kangaroo care (prolonged skin-to-skin contact between baby and carer); detection and management of infections); and infancy and childhood (e.g. adequate nutrition; prevention and management of illness). However, despite some progress, the burden of poor sexual, reproductive, maternal, newborn, and child health remains substantial. New interventions are urgently needed to support further improvement in SRMNCAH, especially in low- and middle-income countries.

Description of the intervention

Targeted client communication (TCC), also referred to as health promotion messaging or behaviour change communication, is the transmission of targeted health content to a specified population or people within a predefined health or demographic group (WHO 2018). TCC can fall along a continuum of tailored communication, such as individualised or personalised notifications, as well as untailored content which draws on predetermined content developed for the identified population group (Hawkins 2008). In order to define the populations for the TCC, eligible people need to be identified and subscribed into a system that allows the transmission of the health content information. Additionally, the health system initiates the first transmission of information, rather than a client seeking information, as in telemedicine and on-demand information services. Following this initial communication from the health system to the client, clients may subsequently respond or continue engagement with the health system, also referred to as bidirectional communication. In contrast, non-targeted client communication (non-TCC) is the transmission of health promotion content delivered to the general population or an undefined population.

TCC has the potential to improve RMNCAH through targeting knowledge, motivation, and behaviour change in order to increase client demand and utilisation of the essential interventions detailed in the WHO Guidelines for Reproductive, Maternal, New-born and Child Health described above (PMNCH 2011). For example, for the successful promotion of breastfeeding, TCC may enhance the provision of health system services through the provision of education relating to breastfeeding, providing links to local services, and providing social support.

Mobile devices may be a particularly effective way of delivering TCC. Mobile phone ownership is almost universal in high-income countries and estimated to have reached over 90% in low- and middle-income countries (ICT 2016). Phones are generally carried wherever people go and can be accessed 24 hours a day. Given their broad reach, mobile devices may provide a cost-effective mechanism for engaging with target populations and delivering health information relating to SRMNCAH.

How the intervention might work

TCC via mobile devices can be used to target the individual-level knowledge, attitudes, and behaviours of importance for the prevention and management of health issues, including those relating to the WHO essential interventions for reproductive,

maternal, new-born, and child health (PMNCH 2011). For example, mobile device-based interventions can (Kaufman 2017):

- provide information and education relevant to the health issue being targeted (e.g. information relating to breastfeeding, adequate child nutrition, vaccinations, and the recognition of symptoms of severe childhood infections);
- facilitate timely access to health advice and services when required (e.g. by providing details of local healthcare services);
- provide reminders (e.g. for HIV medication adherence; for antenatal appointment attendance; for childhood vaccination appointment attendance);
- provide social and psychological support for the behaviour change targeted (e.g. through the provision of encouragement and positive reinforcement; and specifically targeting of psychological factors such as lack of motivation and low self-efficacy).

Why it is important to do this review

Mobile device-based interventions are of particular interest given their low-cost and potential for widespread delivery, however, the current evidence base supporting their implementation for the improvement of maternal, newborn, and child health is limited. The most recent reviews concerned with the effectiveness of mobile device-based interventions for maternal, newborn, or child health have been limited to studies conducted in low/middle-income countries and have included non-randomised controlled trials, which are prone to bias (Lee 2016). Broader reviews of randomised controlled trials (RCTs) of digital health interventions for healthcare consumers (e.g. Free 2013a) are in need of updating to consider the more recent emerging evidence in this field. Other reviews of relevance, concerned with preventive healthcare, reminders for appointment attendance, and self-management of long-term illnesses have focused specifically on SMS (short message service) and MMS (multimedia message service) mobile phone messaging (de Jongh 2012; Gurol-Urganci 2013; Vodopivec-Jamsek 2012), thereby excluding other phone-based delivery mechanisms, such as voice calls, Interactive Voice Response (IVR), and mobile application delivered instant messages.

This review is one of two linked systematic reviews which were carried out to directly inform WHO guidelines on digital interventions for health system strengthening (WHO 2019). This review focuses on the effectiveness of TCC via mobile devices for maternal, newborn, and child health, and the other review examines the effectiveness of TCC via mobile devices for sexual and reproductive health (Palmer in preparation). Although the potential for mobile and digital technologies is acknowledged, there remains considerable demand from ministries of health, donors, and decision-makers for evidence-based guidance on the value of digital tools for improving health. In response to this global need for government decision-makers, the WHO has developed a guideline on digital interventions for health system strengthening to inform government-led investments. In combination, the current review, and the linked review focusing on sexual and reproductive health (Palmer in preparation) complement a qualitative evidence synthesis on the use of TCC for RMNCAH (Ames 2019); combined, these reviews aim to provide a comprehensive overview of the impact, acceptability, and implementation considerations for formulating guideline recommendations.

OBJECTIVES

The overall aim of this review was to assess the effects of TCC via mobile devices on health behaviour, service use, and health and well-being for maternal, new-born and child health. This review focuses on RMNCH priorities in LMIC relating to the WHO essential interventions for reproductive, maternal, new-born and child health (PMNCH 2011).

Our specific objectives relate to three distinct populations and outcomes relevant to these populations. For each population group outlined below, we sought to determine whether targeted client communication via mobile devices can address challenges related to health and well-being, health behaviour, and service utilization. Interventions and comparisons are the same throughout.

- To assess the effects of TCC via mobile devices on behavioural, health/well-being, and service utilisation outcomes relevant to maternal and new-born health among pregnant and postpartum women (up to 6 weeks) and their partners or others who support them.
- To assess the effects of TCC via mobile devices on behavioural, health/well-being, and service utilisation outcomes relevant to maternal and new-born health among pregnant and postpartum women (up to 6 weeks) living with HIV and their partners or others who support them.
- To assess the effects of TCC via mobile devices, delivered to parent and caregivers (i.e. legal guardians, not healthcare professionals) of children under the age of 5 years, on behavioural, health/well-being, and service utilisation outcomes relevant to child health

Secondary objectives

Had there been sufficient studies we planned to assess whether the effects of targeted client communication accessible via mobile devices differ according to:

- Purpose of the intervention (e.g. to remind/recall versus to inform/educate or to support);
- Income region (by World Bank income group) (World Bank 2017);
- Delivery mechanism (e.g. voice, SMS, interactive voice response).

METHODS

Criteria for considering studies for this review

Types of studies

We included randomised controlled trials (RCTs). We included full-text studies, conference abstracts, and unpublished data irrespective of their publication status and language of publication.

We excluded small-scale studies which had randomised fewer than 20 participants.

Types of participants

We included trials with the following participants:

- pregnant and postpartum women up to six weeks after birth, and their partners or others who supported them (where pregnancy status had not been disaggregated, studies in which

at least 70% of women were pregnant or up to six weeks postpartum);

- pregnant and postpartum women up to six weeks after birth living with HIV, and their partners or others who supported them (where HIV status had not been disaggregated, studies in which at least 70% of pregnant and postpartum women were living with HIV);
- parents and carers of children aged under five years (where age had not been disaggregated, studies in which at least 70% of children were under five years of age).

Types of interventions

We included trials that assessed TCC delivered via mobile devices, where the content of the communication was intended to improve maternal, new-born, or child health, or a combination of these.

Targeted client communication

By TCC, we mean the transmission of targeted health content to a specified population or people within a predefined health or demographic group. Unless otherwise stated, we use the terms 'clients', 'patients', and 'consumers' to refer to the individuals whose behaviour, health service use, or health and well-being were being targeted.

We included all of the following:

- studies in which the healthcare consumers were the recipients of the transmitted information;
- studies in which health content was transmitted from the health system to the client, also referred to as unidirectional communication;
- studies in which health content was transmitted from the client to the health system or a health worker, provided that the first communication was initiated by the health system to the client's mobile device. This can occur as bidirectional communication in which clients may respond or exchange with the health system following an initial communication from the health system to the client.

We excluded:

- studies in which the communication between the client and health system was first initiated by the client. Studies in which clients initiated contact with providers were included in a separate review on client-to-provider telemedicine ([Gonçalves-Bradley 2018](#));
- studies in which health content was transmitted to the general population or an undefined population group;
- studies in which the client used fully automated services, including websites, to self-care or access clinical information;
- studies in which TCC was combined with a health worker tool for tracking client's health status, as this combination was included in a separate review ([Agarwal 2018](#)).

Mobile device/multimedia delivery of targeted health communication

By mobile devices, we mean mobile phones of any type (but not analogue landline telephones), as well as tablets and personal digital assistants, which facilitate communication via different multimedia channels including SMS, voice calls, IVR, MMS, and

smartphone applications (apps) when used for instant messaging purposes.

We included studies that used the following communication channels:

- mobile text messaging (including SMS, and unstructured supplementary service data (USSD));
- MMS, including video and audiovisual messages;
- IVR;
- voice calls and call-backs;
- WhatsApp and other instant messaging services (such as Facebook messenger);
- apps, only when they provided an instant messaging function to provide TCC.

We excluded studies that use the following communication channels:

- web portals, applications, and websites that did not have a targeted communication component to notify clients (i.e. which did not provide an instant messaging function, and thereby provided passive information which relied on clients to actively access);
- emails alone;
- social media websites such as Facebook, Baidu, Twitter (unless there was explicit mention of the provision of instant messaging services to individuals to provide target client communication).

Mixed modes of delivery

We included studies in which the intervention delivered to mobile devices was the primary intervention component under evaluation.

When considering interventions delivered by multiple modes, we included interventions involving additional components which could have conceivably been delivered by a mobile device (e.g. an intervention delivered by SMS in combination with email, websites, social media) as all of these delivery mechanisms would allow the entire intervention to be received via a mobile device. Interventions including additional components that could not conceivably be delivered by a mobile device were excluded (e.g. an intervention delivered by SMS in combination with face-to-face counselling).

Content and purpose of the targeted client communication

We included TCC dealing with the health issues listed below. We derived the list of health issues from two key resources by the WHO on Essential Interventions for RMNCH:

- Partnership for Maternal, Newborn & Child Health. 2011. *A Global Review of the Key Interventions Related to Reproductive, Maternal, Newborn and Child Health (RMNCH)* ([PMNCH 2011](#)); and
- WHO, *Packages of Interventions for Family Planning, Safe Abortion care, Maternal, Newborn and Child Health* ([WHO 2010](#)).

For pregnant and postpartum women (up to six weeks), interventions could target:

- ANC;
- birth preparedness;
- skilled attendant at birth;

- emergency obstetric care;
- postpartum care;
- kangaroo mother care;
- tetanus immunisation;
- anaemia prevention and control;
- STI testing and treatment in pregnancy;
- sexual violence;
- malaria prevention and treatment;
- smoking cessation during pregnancy;
- antiretroviral (ARV) adherence (for pregnant and postpartum women living with HIV);
- early infant diagnosis (for pregnant and postpartum women living with HIV);
- retention of mother and infant pairs in elimination of mother-to-child transmission (eMTCT) care (for pregnant and postpartum women living with HIV).

Parents and other carers of children under five years of age, interventions could target:

- postnatal care;
- Immunisation;
- breastfeeding;
- integrated management of newborn and childhood illnesses (IMNCI);
- water, sanitation, and hygiene (WASH);
- Management of diarrhoeal illnesses, oral rehydration salts (ORS), zinc;
- growth monitoring and nutrition;
- early infant diagnosis in HIV-exposed children; ARV therapy for HIV-exposed and HIV-infected children;
- early childhood development.

Interventions could serve at least one of the following purposes (Kaufman 2017):

- to inform and educate identified clients;
- to remind and recall identified clients;
- to teach skills to identified clients;
- to provide support (i.e. for the behaviour change targeted, disease prevention, or health improvement);
- facilitate decision-making;
- enable communication.

We have only included health issues that could potentially be addressed through targeted communication to the *client*, as opposed to those that relate to the provision of clinical care which would be targeted through communication to the healthcare provider. Further details of outcomes that were included can be found below.

Types of comparisons

We included trials with the following comparisons:

- targeted communication delivered to the client via mobile device compared with standard care or no intervention;

- targeted communication accessible to the client via mobile device compared with targeted, non-digital communication (e.g. letters, face-to-face communication with clients);
- targeted communication accessible to the client via mobile devices compared with non-targeted, digital communication via mobile devices (e.g. digital communications which did not target issues relating to maternal, newborn or child health)

We excluded comparisons of:

- one type of targeted communication accessible to the client via mobile devices compared with another type of targeted communication accessible via mobile devices held by the client (e.g. mobile messaging compared with mobile voice);
- studies that compare different technical specifications of telecommunication technologies (e.g. different communication channels, software, etc.);
- studies comparing TCC via mobile device in addition to another intervention that could not conceivably be delivered by mobile device (e.g. face-to-face counselling), compared with TCC via mobile device alone;
- studies comparing TCC via mobile device in addition to another intervention that could not conceivably be delivered by mobile device (e.g. face-to-face counselling), compared with standard care/no intervention.

Types of outcome measures

The outcome measures extracted were according to the population targeted. Below we present the outcomes relevant for pregnant and postpartum women (up to six weeks) living without HIV and living with HIV and the outcomes extracted for children under the age of five years. We extracted both objectively measured and self-reported outcomes for all lengths of follow-up reported.

Where a study reported the same outcome measure for multiple time points, we extracted data for the outcome at the longest follow-up time point. Where we identified studies that reported multiple outcome measures falling under the same outcome category, we extracted all outcome measures. For example, under the outcome category of 'partner violence', we would have extracted measures of sexual, physical, and emotional violence, to ensure that we were able to reflect different aspects within a single outcome category.

Where we identified studies that reported multiple outcome measures of the same *outcome*, we applied a set of rules to decide which outcome measure(s) to report in our review in order to avoid over-representing single trials that reported on multiple measures relating to a single outcome. Where a study reported both dichotomous and continuous measures relating to a single outcome, we applied the following rules to identify one dichotomous outcome measure and one continuous outcome measure to present. The rationale for presenting both a dichotomous and a continuous outcome measure, where available, was because trials may have been underpowered to detect a difference in a clinically important dichotomous outcome (e.g. proportion adherent to medication), but may have power to detect a mean difference (MD) in the equivalent continuous outcome (e.g. MD in number of days covered by medication).

Where objective measurement was possible, we prioritised reporting an objectively measured outcome over a self-reported

outcome measure. For example, had a study included the outcome of STI status, and recorded both biochemically confirmed STI status and self-reported STI status, we would have reported the biochemically confirmed STI status outcome. For outcomes that had not been directly objectively measured, likely for some health behaviour outcomes, we listed the outcomes of the trial (without considering either effect size or its statistical significance) and two review authors made a decision about which was most 'clinically' important or which was the most appropriate measure of the outcome under focus (or both). For example, in terms in clinical importance, if a study reported the outcome measure of attendance to at least one antenatal appointment and the outcome measure of attendance to all antenatal appointments, we presented the outcome relating to attendance to *all* antenatal appointments as this likely to have greater clinical impact.

Primary outcomes

Pregnant and postpartum women (living without HIV and living with HIV)

The following outcomes were identified based on the list of health issues that could be targeted by included interventions. As described above, these are based on two key resources by the WHO on Essential Interventions for RMNCH:

- the Partnership for Maternal, Newborn & Child Health. 2011. *A Global Review of the Key Interventions Related to Reproductive, Maternal, Newborn and Child Health (RMNCH) (PMNCH 2011)*; and
- WHO, *Packages of Interventions for Family Planning, Safe Abortion care, Maternal, Newborn and Child Health (WHO 2010)*.

Health behaviour change

- Smoking cessation; b. alcohol consumption; c. adherence to preventive regimens for pre-eclampsia (calcium, magnesium, low-dose aspirin); d. adherence to antenatal regimens (e.g. folic acid); e. adherence to preventive/treatment regimens for anaemia (ante- and postnatal iron supplements); f. adherence to malaria prevention strategies (insecticide-treated nets (ITNs), intermittent-preventive treatment (IPTp) with sulphadoxine-pyrimethamine (SP)); g. adherence to treatment for treatable infections (e.g. chlamydia, syphilis, mastitis); h. adherence to management strategies for pre-existing conditions, e.g. diabetes; i. adherence treatment for mental health conditions; j. initiation of kangaroo care; k. initiation of breastfeeding; l. postpartum contraceptive uptake; m. adherence to deworming regimen; n. other lifestyle changes, e.g. exercise, health diet; o. adherence to ARVs (tablet count, prescription data).

Service utilisation

- ANC:** a. attendance for ANC appointment (e.g. more than one appointment with skilled personnel, more than four appointments with skilled personnel, attendance for vaccinations, attendance for screening, e.g. HIV, syphilis, anaemia, hypertensive disorders, attendance for malaria prevention services e.g. ITNs, IPTp with SP); b. attendance for high-risk pregnancies (preterm, multiple pregnancies, previous maternal haemorrhage); c. attendance to eMTCT care.
- Intrapartum care:** a. place of birth (home, hospital); b. skilled attendant at birth.

- Postnatal care:** a. attendance for postpartum care appointment; b. attendance for information and counselling on nutrition, safe sex, family planning and provision of contraceptive methods; c. attendance to eMTCT care.
- Postnatal care (newborn):** a. attendance for postpartum care appointment.

Health status and well-being

- Maternal morbidity/mortality (physical):** a. STI (any) status; b. HIV status/HIV management (CD4 count, viral load); c. tetanus; d. syphilis; e. anaemia (e.g. haemoglobin, haematocrit, ferritin); d. malaria; e. pre-eclampsia (blood pressure); f. birth-related complications (e.g. severe bleeding, receipt of blood transfusion, postdelivery haemoglobin, infection); g. postabortion complications (e.g. incomplete abortion, severe bleeding, infection); h. mastitis; i. maternal mortality (objective and self-reported measures).
- Maternal morbidity (mental):** a. depression (validated measure); b. anxiety (validated measure); c. puerperal psychosis.
- Neonatal Health:** a. gestational age at birth; b. birth weight; c. Apgar score; d. perinatal death; e. neonatal death; d. HIV status.
- Partner violence:** a. sexual violence; b. physical violence; c. emotional violence (objective, e.g. hospital admissions and self-report measures).
- Well-being:** a. validated measures of health-related quality of life.

Parents and other carers of children under five years of age

Health behaviour change of parent/carers

- Breastfeeding:** a. duration of breastfeeding; b. exclusive breastfeeding (among babies under six months of age).
- Nutrition:** a. introduction of complementary foods (among babies aged six to eight months); b. dietary diversity/adequacy (children aged six to 23 months who received foods from four or more food groups); c. minimum meal frequency; d. consumption of iron-rich or iron-fortified foods; e. adherence to vitamin A supplements.
- Hygiene practices:** a. handwashing; b. general hygiene (e.g. washing kitchen utensils, sterilisation of household goods); c. cord care; d. safe disposal of faeces.
- Treatment adherence:** a. adherence to ARV in HIV-exposed and HIV-positive children; b. adherence to prescribed regimen for other illnesses (e.g. local infections); c. adherence to treatment for malnutrition; d. correct treatment for diarrhoeal disease (e.g. use of ORS, rice water); e. adherence to prescribed regimen for prevention of malaria (ITNs, IPTp with SP).

Service utilisation

- a. attendance for vaccinations (including measles/diphtheria/pertussis/tetanus vaccine); b. attendance to healthcare services when child showed symptoms of severe illness (e.g. sepsis); c. antibiotic treatment for pneumonia; d. time to diagnosis of HIV in HIV-exposed infants.

Health status and well-being of child

- Normal growth:** a. weight; b. height; c. body mass index; d. growth trajectories (e.g. stunting).
- Nutritional status:** a. anaemia; b. vitamin deficiencies.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

- **Disease:** a. malaria; b. measles; c. HIV; d. pneumonia; e. diarrhoeal disease; f. tetanus; g. pertussis; h. diphtheria; i. other infections; j. CD4 count, viral load (among children with HIV); k. child mortality.
- **Cognitive development:** a. achievement of developmental milestones.

Secondary outcomes

We extracted the following secondary outcomes for all populations.

- Patient/client acceptability and satisfaction with the intervention (among those who received the intervention).
- Resource use, including cost to the system (e.g. human resources/time, supplies, and equipment). This measure had to be prespecified and available directly from the main trial report (i.e. we did not search for separate reports on cost-effectiveness analyses).
- Unintended consequences (these could have included: misreading or misinterpretation of data; transmission of inaccurate data; loss of verbal and non-verbal communication cues, including between provider and client; issues of privacy and disclosure; affecting interpersonal relationships; negative impact on equity; issues with implementation fidelity resulting in an undesirable effect on health outcomes, such as failure or delay in the message delivery that results in missed appointments).

Reporting one or more of the outcome measures listed here in the trial was not an inclusion criterion for the review.

Search methods for identification of studies

Electronic searches

We searched the following electronic databases:

- Cochrane Central Register of Controlled Trials (CENTRAL, the Cochrane Library) (August 2017);
- MEDLINE (OvidSP) (July 2017);
- Embase Classic + Embase (OvidSP) (August 2017);
- POPLINE (August 2017);
- WHO Global Health Library (August 2017).

Search strategies were tailored according to database requirements. The search strategies for each database are reported in [Appendix 1](#); [Appendix 2](#); [Appendix 3](#); [Appendix 4](#); and [Appendix 5](#). The search strategies were designed to retrieve studies relevant to the two linked reviews: 1. Targeted client communication via mobile devices for improving maternal, neonatal, and child health, and 2. Targeted client communication via mobile devices for improving reproductive and sexual health ([Palmer in preparation](#)).

We searched for studies published since 2010. A review entitled "The effectiveness of mobile-health technology-based health behaviour change or disease management interventions for health care consumers: a systematic review" by [Free 2013a](#) carried out searches for studies published up to 2010. With the exception of our focus on a narrower range of health issues (sexual, reproductive, maternal, newborn, child health), our inclusion and exclusion criteria were consistent with [Free 2013a](#). Therefore, we included all relevant studies from [Free 2013a](#) covering the period up to 2010.

Searches were initially carried out in July/August 2017 and all relevant studies identified up to this date have been reported in this review. Prior to publication, a search update was carried out in July 2019. Relevant studies from the update search were not included in the review but reported in the [Characteristics of studies awaiting classification](#) section. The PRISMA diagram in [Figure 1](#) represents the flow of studies up to July 2019.

Figure 1. Study flow diagram. MNCH: maternal, neonatal, and child health; SRH: sexual and reproductive health.




Figure 1. (Continued)

(meta-analysis)

Searching other resources

We searched for ongoing trials in the following trial registries:

- WHO ICTRP (International Clinical Trials Registry Platform; www.who.int/ictrp) (July 2019; [Appendix 6](#));
- US National Institutes of Health Ongoing Trials Register ClinicalTrials.gov (www.clinicaltrials.gov) (July 2019; [Appendix 7](#)).

We also searched Epistemonikos (www.epistemonikos.org/) (July 2019) to identify all relevant systematic reviews and screened them for relevant primary studies. Additionally, WHO issued a call for papers through popular digital health communities of practice such as the Global Digital Health Network and Implementing Best Practices, to identify additional primary studies as well as grey (unpublished) literature.

On completion of screening, we ran a search for all related citations of the included studies, and these citations were screened.

Data collection and analysis

The review was carried out as described in the protocol ([Palmer 2018a](#)), with exceptions noted in the [Differences between protocol and review](#) section.

Selection of studies

Two review authors independently screened all titles and abstracts identified from searches to determine those which met the inclusion criteria. We retrieved in full text any papers identified as potentially relevant. Two review authors independently screened full-text articles for inclusion or exclusion, with discrepancies resolved by discussion and by consulting a third review author where necessary. Selected potentially relevant papers excluded from the review at the full-text stage are listed as excluded studies, with reasons provided in the [Characteristics of excluded studies](#) table. We recorded citation details and any available information about ongoing studies (see [Characteristics of ongoing studies](#) table). We collated and reported details of duplicate publications, so that each study (rather than each report) was the unit of interest in the review. The screening and selection process is presented in an adapted PRISMA flow chart ([Figure 1](#)) ([Liberati 2009](#)).

Data extraction and management

One review author extracted data from included studies, and this was cross-checked by a second review author. Any discrepancies were resolved by discussion until consensus was reached, or through consultation with a third review author where necessary. We developed and piloted a data extraction form using the Cochrane Consumers and Communication Review Group Data Extraction Template (available at: ccrg.cochrane.org/author-resources).

Data extracted included the following items:

- methods: study design; total duration of study; study setting; and date of study;

- participants: number randomised; number lost to follow-up/withdrawn; number analysed; mean age; age range; gender; and inclusion criteria and exclusion criteria;
- interventions: details of intervention and comparison group conditions (including detail of what 'standard care' included). This included intervention delivery mechanism (e.g. text messages/MMS/apps/combined); how the intervention was developed; if the intervention was personalised; and frequency and duration of intervention receipt;
- outcomes: primary and secondary outcomes specified and collected; unintended consequences; and time points reported;
- notes: funding for trial and notable conflicts of interest of trial authors.

All extracted data was entered into Review Manager 5 ([Review Manager 2014](#)) by one review author, and checked for accuracy against the data extraction sheets by a second review author working independently.

Assessment of risk of bias in included studies

We assessed and reported the methodological risk of bias of included studies in accordance with the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)), and the guidelines of the Cochrane Consumers and Communication Review Group ([Ryan 2013](#)), which recommends the explicit reporting of the following individual elements for RCTs: random sequence generation; allocation sequence concealment; blinding (participants, personnel); blinding (outcome assessment); completeness of outcome data; selective outcome reporting; and other bias. We considered blinding separately for different outcomes where appropriate (e.g. blinding may have the potential to differently affect subjective versus objective outcome measures). Under 'Other bias', we considered other potential sources of bias such as the presence of baseline imbalances related to the outcome under study, and evidence of contamination. For cluster-RCTs we also assessed and reported the risk of bias associated with an additional domain: selective recruitment of cluster participants. We judged each item as being at high, low, or unclear risk of bias as set out in the criteria provided by [Higgins 2011](#), and provided a quote from the study report and a justification for our judgement for each item in the 'Risk of bias' table. One review author independently assessed the risk of bias of included studies and a second review author cross-checked all assessments. Any disagreements were resolved by discussion to reach consensus.

Measures of treatment effect

For individually randomised controlled trials reporting dichotomous outcomes, we analysed data based on the number of events and the number of people assessed in the intervention and comparison groups. We used these to calculate the risk ratio (RR) or odds ratio (OR) and 95% confidence interval (CI). For continuous measures, we analysed data based on the mean, standard deviation (SD) and number of people assessed for both the intervention and comparison groups to calculate mean difference (MD) and 95% CI. If the MD was reported without individual group data, we used this to report the study results. If

more than one study measured the same outcome using different tools, we calculated the standardised mean difference (SMD) and 95% CI using the inverse variance method in Review Manager 5 (Review Manager 2014). For cluster RCTs, where the study reported cluster-adjusted effect estimates (e.g. RR, OR) or counts, these were extracted.

Unit of analysis issues

We included both individually and cluster RCTs. Where cluster-RCTs were included, in the first instance, we extracted effect estimates that were adjusted for within-cluster correlation by the study authors. If no adjusted estimates were available in the study report, we derived intracluster correlation coefficients (ICCs) for the outcomes of interest from other included studies or from a paper by Pagel 2011 and calculated adjusted effect estimates prior to meta-analysis. We calculated a design effect using these ICCs and mean cluster size, which was used to calculate the effective number of events per control/intervention and effective number of participants per control/intervention (for methods see Higgins 2011).

Where we identified multiarm trials with more than one relevant intervention arm but only one control arm, we pooled the intervention arms for a single pair-wise comparison as recommended by the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We excluded intervention arms from multi-arm trials that were not eligible for inclusion, but listed them in the [Characteristics of included studies](#) table.

Dealing with missing data

We planned to contact investigators or study sponsors to obtain missing data where possible (e.g. when a study was identified as abstract only); however, we did not have the resources to do so. Where a study was identified as abstract only (Kamau-Mbuthia 2013; Kebaya 2014), we searched for associated full reports. We commented on the potential impact of studies that apparently measured outcomes but did not contribute usable data in the [Effects of interventions](#) section. Where missing data were considered a potential source of serious bias, we planned to conduct a sensitivity analysis to explore the impact of including such studies in the overall assessment of results.

For participant data, we planned, where possible, to conduct analysis on an intention-to-treat basis; otherwise data were analysed as reported. We reported loss to follow-up and assessed this as a source of potential bias.

Assessment of heterogeneity

We carried out meta-analyses when we considered it meaningful to do so. In order to be pooled, studies had to have been conducted among the same population (i.e. we did not pool studies across the three distinct populations as defined under [Types of participants](#)), targeting and measuring the same outcome, and comparing the intervention with similar control groups conditions (i.e. pooling was carried out separately for the following three comparisons: TCC delivered by mobile device versus 1. standard care or no intervention, 2. non-digital TCC, or 3. digital non-targeted communication. We assessed the degree of heterogeneity by visual inspection of forest plots and by examining the Chi² test for heterogeneity. Heterogeneity was quantified using the I² statistic and interpreted by considering the size and direction of effects

and the strength of the evidence for heterogeneity, based on the P value from the Chi² test (Higgins 2011). Where heterogeneity was present in pooled effect estimates, we intended to explore possible reasons for variability by conducting our prespecified subgroup analysis; however, there were an insufficient number of studies in the pooled analyses to conduct meaningful subgroup analyses. Where we noted other potential explanations for high heterogeneity (e.g. differing baseline level of risk), and there was a sufficient number of studies, we conducted subgroup analyses to examine these.

Assessment of reporting biases

We assessed reporting bias qualitatively based on the characteristics of the included studies. For example, if only small studies that indicated positive findings were identified for inclusion, or information that we obtained from contacting experts and authors of studies suggested that there were relevant unpublished studies, we would have considered this as potential evidence of publishing bias and would have reported it as so. If we had identified sufficient studies (at least 10) for inclusion in a meta-analysis, we planned to construct a funnel plot to investigate small-study effects, which may indicate the presence of publication bias. We also searched for trial registry entries and published protocols of all included studies, and used this to assess the risk of bias due to selective reporting.

Data synthesis

We decided whether to meta-analyse data based on whether the included trials were similar enough in terms of participants, settings, intervention, comparison, and outcome measures to ensure meaningful conclusions from a statistically pooled result.

For studies which measured the same outcome at different time points, we extracted the outcome measured at the longest follow-up time point. We had planned to categorise lengths of follow-up as follows: short-term follow-up: less than three months; moderate-term follow-up: three to 12 months; long-term follow-up: more than 12 months. However, given the limited number of studies with the same aim, comparison, and outcome measure that could be pooled, we decided to pool outcomes across different lengths of follow-up.

Due to the anticipated variability in the interventions of included studies, we used random-effects models for meta-analysis. The primary meta-analyses included all studies regardless of their score on the risk of bias assessment.

For continuous outcomes, we calculated MDs with 95% CIs. If more than one study measured the same continuous outcome using different tools, we calculated the SMD and 95% CI using the inverse variance method in Review Manager 5 (Review Manager 2014). We calculated RR or OR with 95% CI for dichotomous outcomes. For cluster-RCTs that presented adjusted effect estimates (e.g. OR, RR), we combined these estimates with using generic inverse variance random-effects (Higgins 2011).

Subgroup analysis and investigation of heterogeneity

We planned to carry out the following subgroup analyses for the objectively measured outcomes of health status:

- income region (by World Bank income group: lower income, lower-middle income, upper-middle income, high income) ([World Bank 2017](#));
- delivery mechanisms (i.e. mobile phone messaging only, mobile apps only, combined mobile phone messaging and apps, combined apps and other).

However, there was an insufficient number of studies reporting the same health status outcomes to conduct these subgroup analyses.

Sensitivity analysis

We planned to carry out the following sensitivity analyses:

- only including studies with low risk of bias on the sequence generation, allocation concealment, and incomplete outcome data domains;
- only including studies with objectively measured outcomes.

However, there was only a sufficient number of studies to conduct the first specified sensitivity analysis.

'Summary of findings' table

We prepared 'Summary of findings' tables to present the results of meta-analyses or narrative synthesis (or both) for the major comparisons of the review, for the following prespecified outcomes.

Pregnant and up to six weeks postpartum women

- Maternal morbidity and mortality combined.
- Neonatal morbidity and mortality combined.
- Breastfeeding.
- Attendance for ANC.
- Birth in a health facility/occurring with a skilled attendant present.
- Unintended consequences.

Pregnant and up to six weeks postpartum women living with HIV

- Maternal morbidity and mortality combined.
- Neonatal morbidity and mortality combined.
- Attendance for ANC (including eMTCT care).
- Birth in a health facility.
- Unintended consequences.
- Neonate HIV status.
- Adherence to ARV therapy.

Parents and other carers of children under five years of age

- Child morbidity and mortality combined.
- Child nutritional status (including anthropometric measures).
- Breastfeeding.
- Clinic attendance for necessary healthcare (e.g. for vaccinations, care for severe illness, HIV tests)
- Hygiene practices (e.g. handwashing).
- Correct treatment taken (e.g. correct treatment for diarrhoeal disease, adherence to ARV therapy).
- Unintended consequences.

Two review authors independently used the GRADE criteria to assess the certainty of the evidence based on the methods described in Chapter 11 of the *Cochrane Handbook for Systematic Reviews of Interventions*, using the GRADEprofiler (GRADEpro) software ([Schünemann 2011](#)). Where meta-analyses were not possible, we presented results in a narrative format.

Ensuring relevance to healthcare decisions

The protocol and review received input and feedback from members of the WHO throughout the review process to ensure the relevance of the review for health policy and practice decisions. The review was also refereed by content experts and including a consumer referee as part of the Cochrane Consumers and Communication Group standard editorial processes.

RESULTS

Description of studies

Results of the search

The results of the search are shown in [Figure 1](#). The search of the databases retrieved 11,259 records, with a further 1726 records identified from clinical trial registries, and one additional record from other sources. After deduplication, 10,332 records remained for title and abstract screening, of which 9206 were clearly irrelevant and excluded. We assessed 1126 full-text articles for eligibility and 168 studies (including 64 ongoing studies) met the inclusion criteria for either this review, or the linked review of TCC via mobile devices for sexual and reproductive health ([Palmer in preparation](#)).

For this review, the search identified 48 eligible studies, of which 21 were added to the [Characteristics of studies awaiting classification](#) section. A total of 27 studies were included in the synthesis reported below, 11 concerned pregnant and postpartum populations, three related to pregnant and postpartum populations living with HIV, and 13 related to parents of children under the age of five years.

Included studies

The [Characteristics of included studies](#) table presents details of the design, methods, participants, intervention, comparison, and outcome measures for the studies included in this review.

Participants and settings

In the 11 studies conducted among pregnant and postpartum populations, the sample size ranged from 68 to 2637, totalling 7626 participants. The three studies concerned with pregnant and postpartum women living with HIV had sample sizes varying from 150 to 550, resulting in 1088 women. All participants were recruited from a mix of community and healthcare settings.

The 13 studies among parents of children aged under five years were conducted with 8749 participants, with sample sizes ranging from 57 to 2054. All trials recruited through healthcare settings or community settings linked with healthcare facilities.

[Table 1](#) presents details of the settings in which each trial was carried out. Of the 11 trials concerned with pregnant and postpartum women, four were conducted in high-income countries, two in upper middle-income countries, four in lower middle-income countries, and one in a low-income country. All

three trials among pregnant and postpartum women living with HIV were carried out in Kenya (a lower middle-income country). Of the 13 studies targeting parents of children under five years of age, five were conducted in high-income countries, seven in lower middle-income countries, and one in a low-income country.

Interventions

Details of the interventions are provided in the [Characteristics of included studies](#) table and [Table 2](#).

Of the 11 studies conducted among pregnant and postpartum populations, nine evaluated interventions that aimed to provide information, education, or support ([Evans 2014](#); [Jareethum 2008](#); [Joshi 2015](#); [Kamau-Mbuthia 2013](#); [Maslowsky 2016](#); [McConnell 2016](#); [Moniz 2013](#); [Naughton 2017](#); [Yudin 2017](#)), and two interventions also provided reminders alongside information/education or support ([Lund 2012](#); [Omole 2018](#)). Six trials delivered the intervention by text messages only ([Evans 2014](#); [Kamau-Mbuthia 2013](#); [Lund 2012](#); [Moniz 2013](#); [Omole 2018](#); [Yudin 2017](#)); two delivered via voice calls only ([Maslowsky 2016](#); [McConnell 2016](#)); and the remaining three studies examined interventions delivered by multiple mechanisms, such as text messages, telephone calls, and email ([Naughton 2017](#)), texts messages and telephone calls ([Jareethum 2008](#)), and multimedia messaging including video and audiovisual messages, combined with voice calls ([Joshi 2015](#)). Four studies did not report on personalisation of the intervention ([Jareethum 2008](#); [Kamau-Mbuthia 2013](#); [Moniz 2013](#); [Yudin 2017](#)); and the rest reported some extent of personalisation, such as messaging specific to the baby's gestational age, choice of language in which to receive intervention, and preferred time of receipt of messaging. In eight of the trials among pregnant and postpartum women, the control group received standard care/no intervention, and in two trials the control group received digital non-targeted communication, both in the form of text messages containing general health information ([Moniz 2013](#); [Omole 2018](#)). One trial had two control groups contributing to this review; of which one was standard care, and the other was non-digital targeted communication (which involved delivering the content of the intervention through home visits by community health workers) ([McConnell 2016](#)).

Of the trials conducted among pregnant and postpartum women living with HIV, one tested an intervention providing reminders ([Kebaya 2014](#)), and the other two provided information, education, or support alongside reminders ([Kassaye 2016](#); [Odeny 2014](#)). One trial delivered the intervention through voice calls ([Kebaya 2014](#)), and the other two via text messaging only ([Kassaye 2016](#); [Odeny 2014](#)). One trial did not report if the intervention was personalised ([Kebaya 2014](#)); one was tailored according to pregnancy stage ([Kassaye 2016](#)); and one allowed participants to select their preferred time of receipt of messaging and their preferred name ([Odeny 2014](#)). Two of the trials among pregnant and postpartum women living with HIV compared the intervention to a control of standard care/no intervention ([Kassaye 2016](#); [Odeny 2014](#)), and, in one trial, the control group received digital non-targeted communication in the form of standard healthcare text messages ([Kebaya 2014](#)).

Most trials among parents of children aged under five years tested interventions providing reminders only, as most of these studies were concerned with efforts to increase attendance for childhood vaccinations ([Ahlers-Schmidt 2012](#); [Bangure 2015](#); [Bigna 2015](#);

[Domek 2016](#); [Eze 2015](#); [Gibson 2017](#); [Haji 2016](#); [Hofstetter 2015a](#); [Niederhauser 2015](#)). Three trials evaluated interventions aiming to provide both reminders and education, information, or support ([Brown 2016](#); [Hannan 2016](#); [Stockwell 2015](#)); and one provided education, information, or support without reminders ([Sharma 2011](#)). Ten interventions were delivered via text messages ([Ahlers-Schmidt 2012](#); [Bangure 2015](#); [Domek 2016](#); [Eze 2015](#); [Gibson 2017](#); [Haji 2016](#); [Hofstetter 2015a](#); [Niederhauser 2015](#); [Sharma 2011](#); [Stockwell 2015](#)), one through voice calls ([Brown 2016](#)), and two through a combination of text messages and voice calls ([Bigna 2015](#); [Hannan 2016](#)). Four studies did not report on whether the intervention had any degree of personalisation ([Ahlers-Schmidt 2012](#); [Bigna 2015](#); [Haji 2016](#); [Hannan 2016](#)); two studies reported that the intervention was not personalised ([Bangure 2015](#); [Brown 2016](#)); and the rest reported at least some degree of personalisation such as use of the child's name or name of clinic at which the appointment was due (or both), and delivery in preferred language. In 10 of the trials among parents of children aged under five years, the control group received standard care or no intervention, in one trial the control received digital non-targeted communication in the form of text messages about general infant health, without the reminders for vaccination appointments that were received by the intervention group ([Niederhauser 2015](#)); and in one trial the control group received non-digital targeted communication in the form of pamphlets ([Sharma 2011](#)). One trial among parents of children had two control groups for the purposes of this review; one received standard care and the other received non-digital targeted communication in the form of sticker vaccination reminders ([Haji 2016](#)).

Outcomes

Health behaviours

Six trials among pregnant and postpartum women reported on at least one health behaviour change outcome. Two trials recorded smoking during pregnancy ([Evans 2014](#); [Naughton 2017](#)), one reported on alcohol consumption during pregnancy ([Evans 2014](#)), three reported on breastfeeding ([Kamau-Mbuthia 2013](#); [Maslowsky 2016](#); [McConnell 2016](#)), one reported on uptake of postpartum contraception ([Maslowsky 2016](#)), and one on adherence to iron/folic acid tablets during the antenatal period ([Joshi 2015](#)).

Two of the three trials among pregnant and postpartum women living with HIV reported on behaviour change – one in the form of maternal utilisation and adherence to ARV therapy, infant ARV/PMTCT, and infant HIV testing ([Kassaye 2016](#)); and one on recorded adherence to newborn PMTCT treatment (nevirapine prophylaxis) ([Kebaya 2014](#)).

Only one of the trials carried out with parents of children aged under five years reported a behaviour change outcome, and this was an indicator of hygiene practices for oral health (the Visible Plaque Index (VPI) in children) ([Sharma 2011](#)).

Service utilisation

Seven of the trials among pregnant and postpartum women reported on health service utilisation outcomes: one measured the proportion of women attending more than four ANC appointments ([Lund 2012](#)); two measured attendance for antenatal influenza vaccine ([Moniz 2013](#); [Yudin 2017](#)); three reported whether birth occurred in a health facility or whether it was attended to by a skilled birth assistant ([Joshi 2015](#); [Lund 2012](#); [Omole 2018](#));

and two measured attendance for postpartum care appointments (Maslowsky 2016; McConnell 2016).

All three of the trials among pregnant and postpartum women living with HIV recorded service utilisation outcomes. One recorded the proportion of births occurring in a health facility, ANC, and mean number of face-to-face or mobile communications with healthcare workers (Kassaye 2016); two reported attendance for postpartum care visits (Kebaya 2014; Odeny 2014).

Eleven of the trials targeting parents of children aged under five years reported on health service utilisation outcomes, most which related to attendance for childhood vaccinations (Ahlers-Schmidt 2012; Bangure 2015; Brown 2016; Domek 2016; Eze 2015; Gibson 2017; Haji 2016; Hannan 2016; Hofstetter 2015a; Niederhauser 2015; Stockwell 2015). Two studies reported on other types of health service utilisation – one on attendance to HIV medical appointments (among HIV-positive and HIV-exposed infants) (Bigna 2015); and one reported on urgent care seeking (Hannan 2016).

Health and well-being

Three trials among pregnant and postpartum women measured health and well-being outcomes. One reported on neonatal diarrhoeal disease and infant growth up to three months postpartum (Kamau-Mbuthia 2013); one reported on preterm delivery, birthweight, and gestational age at birth (Jareethum 2008); and one trial measured perinatal death and maternal mortality (Lund 2012).

Two of three trials among pregnant and postpartum women living with HIV recorded HIV status of the neonate (Kassaye 2016; Odeny 2014), and one trial reported neonatal deaths/stillbirths (Odeny 2014). None of trials among parents of children aged under five years recorded health and well-being outcomes.

Unintended consequences

None of the trials carried out among the three populations of interest reported data on unintended consequences or adverse events.

Acceptability

Eight studies among pregnant and postpartum women recorded at least one measure relating to acceptability/satisfaction with the intervention (Jareethum 2008; Joshi 2015; Lund 2012; Maslowsky 2016; Moniz 2013; Naughton 2017; Omole 2018; Yudin 2017), whereas as none of the studies among pregnant and postpartum women living with HIV did so. Of the trials carried out with parents of children aged under five years, six reported on indicators of acceptability/satisfaction with the intervention (Ahlers-Schmidt 2012; Bangure 2015; Domek 2016; Gibson 2017; Hofstetter 2015a; Stockwell 2015).

Resource use

One trial conducted among pregnant and postpartum women reported on the costs associated with delivering the intervention (Naughton 2017). No trials among pregnant and postpartum women living with HIV reported on the intervention costs. Five of the trials among parents of children aged under five years provided information relating to intervention costs (Bangure 2015; Bigna 2015; Eze 2015; Haji 2016; Hannan 2016).

Funding

Of the trials conducted among pregnant and postpartum women, nine were funded by non-commercial grants (Evans 2014; Joshi 2015; Kamau-Mbuthia 2013; Maslowsky 2016; McConnell 2016; Moniz 2013; Naughton 2017; Omole 2018; Yudin 2017), and one did not state the source of funding (Jareethum 2008).

Two of the trials conducted among pregnant and postpartum women living with HIV reported non-commercial funding sources (Kassaye 2016; Odeny 2014), and one did not report the source of funding (Kebaya 2014).

Most trials conducted among parents of children aged under five years reported non-commercial funding sources (Ahlers-Schmidt 2012; Bangure 2015; Bigna 2015; Domek 2016; Gibson 2017; Haji 2016; Hannan 2016; Hofstetter 2015a; Lund 2012; Stockwell 2015), and four of the studies did not report their source of funding (Brown 2016; Eze 2015; Niederhauser 2015; Sharma 2011).

Excluded studies

Following full-text screening, we excluded 958 articles (Figure 1). The details of relevant excluded trials are provided in the [Characteristics of excluded studies](#) table. Reasons for exclusion included not having a randomised controlled design or enrolling an irrelevant population (e.g. parents of older children/adolescents). The most common reasons for exclusion were related to the intervention being evaluated. In some cases, the intervention was not considered to be TCC as per our definition, or the intervention included a digital tracking component, or was used in conjunction with other interventions (e.g. face-to-face interventions).

Studies awaiting classification

We updated the search in August 2019, and identified a further 21 studies, which are summarised in the [Characteristics of studies awaiting classification](#) table.

Ongoing studies

We identified 32 ongoing studies, which are summarised in the [Characteristics of ongoing studies](#) table.

Risk of bias in included studies

Details of the risk of bias assessments for each of the included studies are presented in the 'Risk of bias' tables in the [Characteristics of included studies](#) table, and in [Figure 2](#).

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Selective cluster recruitment	Blinding of participants and personnel (performance bias): All outcomes	Blinding of objective outcome assessment (detection bias)	Blinding of subjective outcome assessment (detection bias)	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Ahlers-Schmidt 2012	+	?		-	+	-	+	?	?
Bangure 2015	+	?		-	?	-	+	+	+
Bigna 2015	+	+		-	+	-	+	+	?
Brown 2016	?	?	?	-	?		+	?	-
Domek 2016	+	+		-	?	-	+	+	?
Evans 2014	+	?		-		-	+	+	?
Eze 2015	+	?		-	+	-	?	?	-
Gibson 2017	+	+	-	-	+	?	+	+	+
Haji 2016	?	?	-	-	-	-	?	+	-
Hannan 2016	+	?		-		-	+	?	?
Hofstetter 2015a	+	+		+	+	+	+	?	+
Jareethum 2008	+	?		-	+	-	+	+	+
Joshi 2015	?	?	?	-		-	?	?	?
Kamau-Mbuthia 2013	?	?		-		-	?	-	?
Kassaye 2016	?	?	?	-		-	+	?	-
Kebaya 2014	?	?		-	?	?	?	-	?
Lund 2012	?	?	+	-	?		+	+	+
Maslowsky 2016	+	?		-		-	+	?	+
McConnell 2016	+	?		-		-	?	+	?
Moniz 2013	?	+		?	+	?	+	+	+
Naughton 2017	+	?		-	+	-	?	+	?
Niederhauser 2015	+	?		-	?	-	+	?	?
Odeny 2014	+	+		-	+		+	?	+

Figure 2. (Continued)

Niederhauser 2015	+	?	-	?	-	-	?	?
Odeny 2014	+	+	-	+		+	?	+
Omole 2018	?	?	-	-	?	?	-	?
Sharma 2011	?	?	-	+		-	?	+
Stockwell 2015	?	+	-	+	-	+	+	+
Yudin 2017	+	+	+		+	+	+	?

Allocation

Pregnant and postpartum women

One trial demonstrated adequate random sequence generation and allocation concealment, and so was at low risk of bias for both domains (Yudin 2017). Five trials reported adequate random sequence generation and were at low risk of bias on this domain, but did not provide sufficient information on their procedures relating to allocation concealment and therefore were at unclear risk of bias for this domain (Evans 2014; Jareethum 2008; Maslowsky 2016; McConnell 2016; Naughton 2017). One trial was at unclear risk of bias for random sequence generation, and at low risk of bias for allocation concealment (Moniz 2013). Four studies were at unclear risk of bias for both random sequence generation and allocation concealment domains as they provided insufficient information (Joshi 2015; Kamau-Mbuthia 2013; Lund 2012; Omole 2018).

Pregnant and postpartum women living with HIV

One trial demonstrated adequate random sequence generation and allocation concealment, and so was at low risk of bias for both domains (Odeny 2014). Two studies were at unclear risk of bias for both random sequence generation and allocation concealment domains as they provided insufficient information (Kassaye 2016; Kebaya 2014).

Parents and carers of children aged under five years

Four trials demonstrated adequate random sequence generation and allocation concealment, and so were at low risk of bias for both domains (Bigna 2015; Domek 2016; Gibson 2017; Hofstetter 2015a). Five studies reported adequate random sequence generation and were at low risk of bias on this domain, but did not provide sufficient information on their procedures relating to allocation concealment and therefore were at unclear risk of bias for this domain (Ahlers-Schmidt 2012; Bangure 2015; Eze 2015; Hannan 2016; Niederhauser 2015). Three studies were at unclear risk of bias for both random sequence generation and allocation concealment domains as they provided insufficient information (Brown 2016; Haji 2016; Sharma 2011), and one study was at unclear risk of bias for random sequence generation, but low risk of bias for allocation concealment (Stockwell 2015).

Blinding

Pregnant and postpartum women

Nine trials were at high risk of bias for the blinding of participants and personnel domain (Evans 2014; Jareethum 2008; Joshi 2015; Kamau-Mbuthia 2013; Lund 2012; Maslowsky 2016; McConnell 2016; Naughton 2017; Omole 2018). One trial did not provide sufficient information relating to blinding of participants and

personnel and so was at unclear risk of bias for this domain (Moniz 2013). One trial was at low risk of bias for blinding of participants and personnel (Yudin 2017). Three studies demonstrated adequate blinding for objective outcomes and were at low risk of bias on this domain (Jareethum 2008; Moniz 2013; Naughton 2017), two studies did not provide adequate information and so were at unclear risk of bias on this domain (Lund 2012; Omole 2018), and six studies did not record objective outcomes (Evans 2014; Joshi 2015; Kamau-Mbuthia 2013; Maslowsky 2016; McConnell 2016; Yudin 2017). In relation to blinding for subjective outcome assessment, one trial was a low risk of bias (Yudin 2017), two were at unclear risk of bias due to lack of information provided (Moniz 2013; Omole 2018), seven were at high risk of bias (Evans 2014; Jareethum 2008; Joshi 2015; Kamau-Mbuthia 2013; Maslowsky 2016; McConnell 2016; Naughton 2017), and the remaining study did not record any subjective outcomes (Lund 2012).

Pregnant and postpartum women living with HIV

All three trials were at high risk of bias for the blinding of participants and personnel domain (Kassaye 2016; Kebaya 2014; Odeny 2014).

One study was at high risk of bias for subjective outcome assessment, and did not record any objectively measured outcomes (Kassaye 2016), one provided insufficient information and was considered to be at unclear risk of bias for blinding of both objective and subjective outcomes (Kebaya 2014), and third study was at low risk of bias for blinding of objective outcomes, and did not record any subjective outcomes (Odeny 2014).

Parents of children aged under five years

With the exception on one trial which was at low risk of bias for blinding of participants and personnel (Hofstetter 2015a), all trials among parents of children aged under five years were at high risk of bias for this domain. One study was at low risk of bias for blinding of both objective and subjective outcome assessments (Hofstetter 2015a), one study was at low risk of bias for blinding of objective outcome assessments, and did not record any subjective outcomes (Sharma 2011), one study was at low risk of bias for objective outcomes and unclear risk of bias for subjective outcomes (Gibson 2017), and four studies were at low risk of bias for blinding of objective outcome assessments and high risk of bias for blinding of subjective outcome assessments (Ahlers-Schmidt 2012; Bigna 2015; Eze 2015; Stockwell 2015). Three studies were at unclear risk of bias for blinding of objective outcome assessment, and high risk of bias for subjective outcome assessment (Bangure 2015; Domek 2016; Niederhauser 2015). One study was at unclear risk of bias for blinding of objective outcomes, and did not record subjective outcomes (Brown 2016); one study did not record objective outcomes, and was at high risk of bias for subjective

outcome blinding (Hannan 2016); and one study was at high risk of bias for blinding for both objective and subjective outcome assessment (Haji 2016).

Incomplete outcome data

Pregnant and postpartum women

Four of the trials among pregnant and postpartum women were at low risk of bias for incomplete outcome data (Jareethum 2008; Lund 2012; Moniz 2013; Yudin 2017) and seven were at high or unclear risk of bias for this domain (Evans 2014; Joshi 2015; Kamau-Mbuthia 2013; Maslowsky 2016; McConnell 2016; Naughton 2017; Omole 2018).

Pregnant and postpartum women living with HIV

For the incomplete outcome data domain, one trial among pregnant and postpartum women living with HIV was at low risk of bias (Odeny 2014), one was at unclear risk of bias (Kebaya 2014), and one was at high risk of bias (Kassaye 2016).

Parents of children aged under five years

Eight of the trials among parents of children aged under five years were at low risk of bias for incomplete outcome data (Ahlers-Schmidt 2012; Bangure 2015; Bigna 2015; Brown 2016; Domek 2016; Gibson 2017; Hofstetter 2015a; Stockwell 2015), two were at unclear risk of bias (Eze 2015; Haji 2016), and three were at high risk of bias for this domain (Hannan 2016; Niederhauser 2015; Sharma 2011).

Selective reporting

Pregnant and postpartum women

Seven studies were at low risk of bias for the selective outcome reporting domain as their protocols or trial registry entries (or both) could be identified and all expected outcomes were reported (Evans 2014; Jareethum 2008; Lund 2012; McConnell 2016; Moniz 2013; Naughton 2017; Yudin 2017). Two studies were at unclear risk of bias as their protocols could not be identified (Joshi 2015; Maslowsky 2016). Two studies were at high risk of bias on the selective outcome reporting domain due to inconsistencies between the prespecified and actual outcome reporting (Kamau-Mbuthia 2013; Omole 2018).

Pregnant and postpartum women living with HIV

Two of the trials among pregnant and postpartum women living with HIV were at unclear risk of bias for selective outcome reporting as their protocols could not be identified (Kassaye 2016; Odeny 2014), and one was at high risk of bias due to inconsistency between prespecified and actual outcome reporting (Kebaya 2014).

Parents of children aged under five years

Six studies among parents of children aged under five years were at low risk of bias for the selective outcome reporting domain as their protocols or trial registry entries (or both) could be identified and all expected outcomes were reported (Bangure 2015; Bigna 2015; Domek 2016; Gibson 2017; Haji 2016; Stockwell 2015). Seven studies were at unclear risk of bias as their protocols could not be identified (Ahlers-Schmidt 2012; Brown 2016; Eze 2015; Hannan 2016; Hofstetter 2015a; Niederhauser 2015; Sharma 2011).

Other potential sources of bias

Pregnant and postpartum women

Four studies were at low risk of bias for the 'other bias' domain (Jareethum 2008; Lund 2012; Maslowsky 2016; Moniz 2013), and seven were at unclear risk of bias (Evans 2014; Joshi 2015; Kamau-Mbuthia 2013; McConnell 2016; Naughton 2017; Omole 2018; Yudin 2017).

Pregnant and postpartum women living with HIV

For the 'other bias' domain, one trial among pregnant and postpartum women living with HIV was at low risk of bias (Odeny 2014), one was at unclear risk of bias (Kebaya 2014), and one was at high risk of bias as due to important baseline differences between groups (Kassaye 2016).

Parents of children aged under five years

Five of the trials conducted among parents of children aged under five years were at low risk of other bias (Bangure 2015; Gibson 2017; Hofstetter 2015a; Sharma 2011; Stockwell 2015). Five trials were at unclear risk of bias for this domain (Ahlers-Schmidt 2012; Bigna 2015; Domek 2016; Hannan 2016; Niederhauser 2015). Three trials were at high risk of bias for this domain due to: significant demographic differences between clusters in a clustered RCT (Brown 2016); participants being swapped between the intervention and control group when people who did not own a mobile phone had been allocated to the intervention (Eze 2015); and an issue whereby attending a different facility may result in misclassification of the outcome of vaccination (Haji 2016).

Selective cluster recruitment

Pregnant and postpartum women

Three of the 11 trials conducted among pregnant and postpartum women were cluster RCTs and one was at low risk of bias for the domain of selective cluster recruitment (Lund 2012), one was at unclear risk of bias (Joshi 2015), and one was at high risk of bias due to participant enrolment after cluster allocation had been made (Omole 2018). The remaining trials were individually randomised trials and so were not assessed regarding cluster recruitment.

Pregnant and postpartum women living with HIV

One of the three trials conducted among pregnant and postpartum women living with HIV was a cluster RCT and at unclear risk of bias for the domain of selective cluster recruitment (Kassaye 2016). The other two trials were individually randomised trials and so were not assessed regarding cluster recruitment.

Parents of children aged under five years

Three of the 13 trials conducted among parents of children aged under five years were cluster RCTs. One was at unclear risk of bias for the domain of selective cluster recruitment (Brown 2016), and two were at high risk of bias due to convenient enrolment of participants likely to have taken place after cluster allocation to intervention arms (Haji 2016), and participant enrolment after cluster allocation had been made (Gibson 2017). The remaining trials were individually randomised trials and so were not assessed regarding cluster recruitment.

Effects of interventions

See: **Summary of findings 1** Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women) for improving maternal, neonatal, and child health; **Summary of findings 2** Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health; **Summary of findings 3** Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant and postpartum women) for improving maternal, neonatal, and child health; **Summary of findings 4** Digital targeted client communication via mobile devices compared to standard care or no intervention (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health; **Summary of findings 5** Digital targeted client communication via mobile devices compared to non-digital targeted client communication (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health; **Summary of findings 6** Digital targeted client communication via mobile devices compared to digital non-targeted client communication (pregnant and postpartum women with HIV) for improving maternal, neonatal, and child health; **Summary of findings 7** Digital targeted client communication via mobile devices compared to standard care or no intervention (parents of children aged under five years) for improving maternal, neonatal, and child health; **Summary of findings 8** Digital targeted client communication via mobile devices compared to non-digital targeted client communication (parents of children aged under five years) for improving maternal, neonatal, and child health; **Summary of findings 9** Digital targeted client communication via mobile devices compared to digital non-targeted client communication (parents of children aged under five years) for improving maternal, neonatal, and child health

Pregnant and postpartum women

Targeted client communication via mobile devices versus standard care

Summary of findings 1 presents the evidence relating to the effect of TCC via mobile devices compared to standard care or no intervention among pregnant and postpartum women for the outcomes of maternal mortality and morbidity, neonatal mortality and morbidity, breastfeeding, attendance for ANC, receipt of intrapartum care, and unintended consequences.

Health behaviour change

There was at best moderate-certainty evidence relating to the effect of TCC via mobile devices on health behaviours when compared to standard care.

One study carried out in Kenya provided low-certainty evidence relating to the effect of TCC via mobile devices for the promotion of exclusive breastfeeding in a setting where all women in the control group exclusively breastfed (exclusive breastfeeding at 9 weeks postpartum: RR 0.92, 95% CI 0.79 to 1.08; n = 40; low-certainty evidence – downgraded due to risk of bias and imprecision; **Analysis 1.1**) (McConnell 2016). Another study, conducted in Ecuador, which had comparatively lower breastfeeding rates in the standard care arm, suggests that the intervention may increase exclusive breastfeeding at three months postpartum (RR 1.30, 95% CI 1.06 to 1.59; n = 135; low-certainty evidence – downgraded for risk of

bias (unclear risk of bias on the allocation concealment domain and high rates of loss to follow-up) and imprecision; **Analysis 1.1**) (Maslowsky 2016). A second RCT in Kenya found that mobile phone-based support may increase the proportion of postpartum women receiving help with breastfeeding (RR 2.15, 95% CI 1.78 to 2.58; n = 332; low-certainty evidence – downgraded twice for risk of bias; **Analysis 1.2**) (Kamau-Mbuthia 2013).

One study conducted among women living in rural villages in India demonstrated that voice messages delivered to mobile devices probably increased the proportion of participants who took iron and folate tablets for 100 days during pregnancy (RR 1.71, 95% CI 1.42 to 2.07; n = 908; moderate-certainty evidence – downgraded once due to unclear risk of bias for random sequence generation and allocation concealment; **Analysis 1.3**) (Joshi 2015).

We are uncertain about the effects of the intervention on postpartum contraceptive use at three months (study in Ecuador) and at nine weeks postpartum (study in Kenya) because the certainty of the evidence was very low (RR 1.35, 95% CI 0.75 to 2.46; n = 175; downgraded twice for risk of bias and once for imprecision; **Analysis 1.4**) (Maslowsky 2016; McConnell 2016).

Two studies reported on smoking during pregnancy: one UK trial among pregnant women who smoked suggested that the intervention may increase objectively verified continuous abstinence (RR 2.76, 95% CI 0.89 to 8.54; n = 407; low-certainty evidence – downgraded due to risk of bias and imprecision; **Analysis 1.6**) (Naughton 2017). One trial in the USA among pregnant women (who were not recruited based on smoking status) suggested that the intervention may reduce the proportion of pregnant women reporting having smoked in the last 30 days (RR 0.43, 95% CI 0.17 to 1.10; n = 459; low-certainty evidence – downgraded due to risk of bias and imprecision; **Analysis 1.5**) (Evans 2014). The USA study also found that the intervention may have little or no impact on preventing alcohol consumption during pregnancy (RR 1.00, 95% CI 0.97 to 1.03; n = 459; low-certainty evidence – due to risk of bias and indirectness based evidence from a single high-income country; **Analysis 1.7**) (Evans 2014). However, over 97% of participants across the intervention and control group reported no alcohol consumption at baseline leaving little room for any change to be observed postintervention. The effect of the intervention may be different in populations with different levels of baseline risk.

Service utilisation

Evidence for whether TCC via mobile devices increased service utilisation was mixed. TCC via mobile devices may have a modest effect on increasing the proportion of women attending four or more antenatal appointments (cluster-adjusted OR 1.54, 95% CI 0.80 to 2.96; n = 2550; low-certainty evidence – downgraded due to risk of bias and imprecision, CI encompasses both benefit and harm; **Analysis 1.8**) (Lund 2012). Based on pooled analyses of this trial and another conducted in Canada, TCC via mobile devices may increase attendance for antenatal vaccinations (influenza and tetanus) (OR 1.36, 95% CI 0.90 to 2.06; n = 714; low-certainty evidence – downgraded due to risk of bias and imprecision (**Analysis 1.9**) (Lund 2012; Yudin 2017). The study by Lund 2012 also suggests that TCC via mobile devices may increase antenatal attendance for preventive treatment for malaria (cluster adjusted OR 1.69, 95% CI 0.82 to 3.48; low-certainty evidence – downgraded for risk of bias and imprecision; **Analysis 1.10**).

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Two studies examined the effect of TCC via mobile devices on the proportion of births with a skilled attendant present. The intervention may make little or no difference in one low-risk setting where 99% of both the control and intervention group had a skilled attendant at birth (OR 1.00, 95% CI 0.34 to 2.94; $n = 1743$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 1.11](#)) ([Joshi 2015](#)). The second trial conducted in a high-risk setting (Tanzanian urban and rural settings where about 50% of all deliveries take place at home with unskilled attendance) found that there may be benefit in urban populations (cluster-adjusted OR 4.45, 95% CI 1.36 to 14.51; $n = 1077$), but not rural populations (cluster adjusted OR 0.83, 95% CI 0.36 to 1.92; $n = 1473$) (low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 1.11](#)) ([Lund 2012](#)) (this study only reported effect estimates accounting for clustering separately by urban and rural for this outcome).

Two studies examined the effect of TCC via mobile devices on utilisation of care for newborns. One trial conducted in Kenya found that the intervention may make little or no difference to care utilisation (RR 0.98, 95% CI 0.87 to 1.11; $n = 56$; low-certainty evidence – downgraded for risk of bias and imprecision; [Analysis 1.12](#)). However, this was a low-risk setting where 96% of the control group attended care for their newborn leaving little room for demonstration of an improvement ([McConnell 2016](#)). Another trial, conducted in Ecuador – a high risk setting where utilisation of care for newborns was lower (53.3% attendance in the control group) – reported that the intervention may increase the proportion of women attending services for infant care within 10 days of delivery (RR 1.35, 95% CI 1.02 to 1.78; $n = 135$; low-certainty evidence – downgraded for risk of bias and imprecision; [Analysis 1.12](#)) ([Maslowsky 2016](#)). [Maslowsky 2016](#) also reported that "no effect of the intervention on attendance at maternal postpartum visits was observed," without providing data for extraction.

One study in Kenya showed that TCC via mobile device may increase attendance for a postpartum care appointment at 10 days postpartum (RR 1.50, 95% CI 0.30 to 7.52; $n = 56$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 1.13](#)), and may have little or no effect on newborn vaccination (pentavalent and polio at up to nine weeks) (RR 1.08, 95% CI 0.89 to 1.32; $n = 40$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 1.14](#)) ([McConnell 2016](#)).

Health status and well-being

Based on very low-certainty evidence from one study in Tanzania, we are uncertain as to whether TCC via mobile devices has an effect on maternal mortality (RR 2.86, 95% CI 0.30 to 27.40; $n = 2637$; downgraded due to risk of bias and twice for imprecision; [Analysis 1.15](#)) ([Lund 2012](#)). While this effect estimate is large, it is based on very few events (four maternal deaths in the intervention group and one in the control group), and the CI is very wide, consistent with a decrease in mortality as well as a large increase; therefore, we cannot be certain what the effect might be. This study also provided low-certainty evidence that the intervention may slightly reduce severe obstetric complications up to six weeks postpartum (RR 0.86, 95% CI 0.70 to 1.07; $n = 2550$; downgraded due to risk of bias and imprecision; [Analysis 1.15](#)) ([Lund 2012](#)). The trial conducted by [McConnell 2016](#) provided low-certainty evidence that TCC via mobile devices may reduce the likelihood of reporting having experienced 'any maternal health problem' up to 10 days postpartum (RR 0.50, 95% CI 0.90 to 2.76; $n = 56$; downgraded

due to risk of bias and imprecision; [Analysis 1.15](#)). There was low-certainty evidence that TCC via mobile devices may reduce self-reported breast pain and breast engorgement up to three months postpartum (pain: RR 0.28, 95% CI 0.09 to 0.80; $n = 332$; engorgement: RR 0.58, 95% CI 0.31 to 1.10; $n = 332$; [Analysis 1.15](#)) ([Kamau-Mbuthia 2013](#)). There was low-certainty evidence that the intervention may result in little or no difference to the number of acute maternal episodes requiring a clinic visit up to three months postpartum (MD 0.06, 95% CI –0.19 to 0.31; $n = 135$; downgraded due to risk of bias and imprecision; [Analysis 1.16](#)) ([Maslowsky 2016](#)).

There was very low-certainty evidence relating to the effect of the intervention on neonatal mortality and morbidity based on three trials (in Tanzania and Kenya) (OR 1.00, 95% CI 0.61 to 1.64; $n = 2870$; downgraded once due to risk of bias and twice for imprecision; [Analysis 1.17](#)) ([Kamau-Mbuthia 2013](#); [Lund 2012](#); [McConnell 2016](#)). There was moderate-certainty evidence that the intervention may reduce the number of acute neonatal episodes requiring a clinic visit up to three months postpartum (MD –0.53, 95% CI –0.92 to –0.14; $n = 135$; downgraded due to risk of bias; [Analysis 1.18](#)) ([Maslowsky 2016](#)). One study reported on the effect of TCC via mobile devices on gestational age at birth and birth weight (gestational age: MD 0.10 weeks, 95% CI –0.45 to 0.65; $n = 61$; birth weight: MD –173 g, 95% CI –448.87 to 102.87; $n = 61$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 1.19](#)) ([Jareethum 2008](#)). Two studies conducted in Thailand and Zanzibar reported preterm birth (RR 0.85, 95% CI 0.31 to 2.33; $n = 2557$; low-certainty evidence – downgraded due to risk of bias and an imprecise effect estimate, which encompassed both large potential benefit and harm of the intervention; [Analysis 1.20](#)) ([Jareethum 2008](#); [Lund 2012](#)). A sensitivity analysis was also conducted to examine the effect of excluding the cluster RCT by [Lund 2012](#) from the analysis with preterm birth as the outcome. This resulted in no substantive change in the findings; the point estimate remained in the direction of intervention benefit (RR 0.18, 95% CI 0.01 to 3.64; [Analysis 1.21](#)), but the CI was considerably wider, encompassing both benefit and harm.

Targeted client communication via mobile devices versus non-digital targeted communication

[Summary of findings 2](#) presents the evidence relating to the effect of TCC via mobile devices compared to non-digital targeted communication among pregnant and postpartum women for the outcomes of maternal mortality and morbidity, neonatal mortality and morbidity, breastfeeding, attendance for ANC, receipt of intrapartum care, and unintended consequences.

Health behaviour change

One study conducted in Kenya compared TCC via mobile devices to non-digital targeted communication (in the form of home visits by community health workers). This provided low-certainty evidence relating to the effect on exclusive breastfeeding at nine weeks postpartum (RR 0.92, 95% CI 0.79 to 1.07; $n = 42$; [Analysis 2.1](#); downgraded due to risk of bias and imprecision) (this study reported 100% of participants were breastfeeding in the control arm, so there was no room for benefit to be observed and the effect of the intervention may be different in populations with different levels of baseline risk) ([McConnell 2016](#)). The same study also provided low-certainty evidence for the effect on contraceptive use at nine weeks postpartum (RR 1.45, 95% CI 0.78 to 2.69; $n =$

42; downgraded due to risk of bias and imprecision; [Analysis 2.2](#) ([McConnell 2016](#)).

Service utilisation

The trial by [McConnell 2016](#) provided moderate-certainty evidence that the intervention made little or no difference to attendance for newborn care at 10 days postpartum (RR 0.97, 95% CI 0.87 to 1.09; n = 59; downgraded due to risk of bias; [Analysis 2.3](#)) or attendance for newborn vaccination at nine weeks (pentavalent and polio vaccine) (RR 1.01, 95% CI 0.88 to 1.16; n = 42; downgraded due to risk of bias; [Analysis 2.4](#)). This trial also provided low-certainty evidence relating to the intervention effect on attendance for a maternal care postpartum appointment (RR 0.56, 95% CI 0.18 to 1.79; n = 59; downgraded due to risk of bias and imprecision; [Analysis 2.5](#)) ([McConnell 2016](#)).

Health status and well-being

Based on the [McConnell 2016](#) trial, there was low-certainty evidence that the intervention may reduce 'any maternal health problem' reported up to 10 days postpartum (RR 0.19, 95% CI 0.04 to 0.79; n = 59; [Analysis 2.6](#)) and 'any newborn health problem' reported up to 10 days postpartum (RR 0.52, 95% CI 0.25 to 1.06; n = 59; [Analysis 2.7](#)). For both outcomes, the certainty of the evidence was downgraded due to risk of bias and imprecision due to the small number of events.

Targeted client communication via mobile devices versus digital non-targeted communication

[Summary of findings 3](#) presents the evidence relating to the effect of TCC via mobile devices compared to digital non-targeted communication among pregnant and postpartum women for the outcomes of maternal mortality and morbidity, neonatal mortality and morbidity, breastfeeding, attendance for ANC, receipt of intrapartum care, and unintended consequences.

Health behaviour change

No studies reported health behaviour outcomes for the comparison of TCC via mobile devices compared with digital non-targeted communication.

Service utilisation

Based on one study in the USA comparing TCC via mobile devices with digital non-targeted communication, there was low-certainty evidence that the intervention may have little or no effect on attendance for antenatal influenza vaccination (RR 1.05, 95% CI 0.71 to 1.58; n = 204; downgraded due to risk of bias and imprecision; [Analysis 3.1](#)) ([Moniz 2013](#)). Based on a single study conducted in Nigeria we are uncertain as to whether TCC via mobile devices has an effect on the proportion of births occurring with skilled attendant when compared with digital non-targeted communication (RR 1.00, 95% CI 0.69 to 1.45; n = 16; very low-certainty evidence – downgraded twice due to risk of bias and once due to imprecision; [Analysis 3.2](#)) ([Omole 2018](#)).

Health status and well-being

No studies reported health status and well-being outcomes for the comparison of TCC via mobile devices compared with digital non-targeted communication.

Patient/client acceptability and satisfaction with the intervention

There was generally high rates of satisfaction with the interventions received. One UK trial targeting smoking cessation in pregnancy reported that 61.7% of intervention participants thought the messages were helpful, and 80.8% would recommend the support, while 14.2% reported that the messages were 'annoying' and 13.3% discontinued the support by texting 'STOP' ([Naughton 2017](#)). One study in Canada evaluating the effect of text messages aiming to increase influenza vaccine uptake during pregnancy found that 88% of intervention recipients were satisfied with the messages; 81% found the information in the messages useful; 65% thought the messages were appropriately timed; and 60% thought that the number of messages was right (received two messages per week for four weeks) ([Yudin 2017](#)). About 99% of intervention recipients in a trial conducted in Nigeria stated that they thought the SMS reminders improved ANC attendance; over 95% stated the messages influenced their decision to deliver in a health facility and to attend a postnatal clinic; 77.8% stated that the messages helped them to complete immunisations and preventive therapy for malaria in pregnancy; and 96.6% agreed that they supported the use of a SMS platform for informing ANC clients about obstetric danger signs ([Omole 2018](#)). One trial in India reported that 89.3% of the intervention group were satisfied with the content of the voice messages and 97.4% with animations received ([Joshi 2015](#)). One trial among pregnant women in Thailand who received text messages about pregnancy and the early postpartum period reported high levels of satisfaction (a mean score of 9.25, based on a visual analogue scale of 1 to 10 points) ([Jareethum 2008](#)). The majority of participants receiving the SMS intervention in one trial conducted Tanzania rated the messages positively. About 59% stated that receiving text messages influenced the number of times they attended ANC and 71% reported that the messages helped them in various areas, including learning about danger signs in pregnancy and feeling that the health system cared for them ([Lund 2012](#)). The trial of an intervention delivered through phone calls providing postpartum health education and support in Ecuador reported almost universal satisfaction – 98% of intervention participants agreed that they would like to have access to the service at their next birth, and 100% that they would recommend the service to a friend ([Maslowsky 2016](#)). Most recipients of messages communicating the importance of influenza vaccinations during pregnancy along with general health advice (USA) reported that they liked the text messages (90%), found them to be helpful (86%), wanted to receive text messages in the future (82%), and thought they increased their satisfaction with their ANC (80%) ([Moniz 2013](#)).

Resource use

[Naughton 2017](#) reported that the total cost of delivering the 'MiQuit' text messaging intervention was GBP 4.62 per participant, and, based on the relevant incremental quit rate estimate of 3.46%, this resulted in an incremental cost per additional quitter of GBP 133.53 (95% CI –395.78 to 843.62).

Unintended consequences

None of the trials among pregnant and postpartum women collected or reported data on unintended consequences.

Equity considerations

The three trials concerned with breastfeeding were conducted in upper-middle and lower-middle income countries, and two

of which specifically highlighted their inclusion of low-income participants (Kamau-Mbuthia 2013; McConnell 2016). Both trials of interventions aiming to reduce harmful behaviour during pregnancy (smoking and alcohol consumption) were conducted in high-income countries, and it is not possible to assess whether the results are applicable to low-income settings (Evans 2014; Naughton 2017). The two trials of TCC via mobile devices for promoting attendance for antenatal influenza vaccination were also from high-income countries: one primarily included unmarried participants with low levels of education (Moniz 2013), while the other study population was made up of mostly married or partnered women with higher levels of education (Yudin 2017); despite the apparent socioeconomic differences in these populations, both studies demonstrated no benefit.

Of note for consideration of issues relating to equity, is that more than half of the trials applied a language-based criteria in their inclusion/exclusion criteria. Given the nature of the intervention, it is likely that those studies not explicitly stating the need to speak the native language to partake in the study, will have also excluded those lacking such fluency in a particular language. This raises the issue of exclusion of illiterate populations and recent migrants, who are known to be a particularly vulnerable population, but are unable to provide consent to take part in studies that rely on phone-based communications in a specific language.

Pregnant and postpartum women living with HIV

Targeted client communication via mobile devices versus standard care

Summary of findings 4 presents the evidence relating to the effect of TCC via mobile devices compared to standard care or no intervention among pregnant and postpartum women living with HIV for the outcomes of maternal mortality and morbidity, neonatal mortality and morbidity, attendance for ANC (including eMTCT care), receipt of intrapartum care, unintended consequences, neonate HIV status, and adherence to ARV therapy.

Health behaviours

One study reported the effect of TCC via mobile devices on whether mothers took any type of ARVs antenatally and postnatally compared to standard care in Kenya (Kassaye 2016). There was low-certainty evidence suggesting the intervention may make little or no difference to maternal antenatal ARV usage (RR 1.04, 95% CI 0.91 to 1.19; n = 503; downgraded twice due to risk of bias; Analysis 4.1). The same study also provided very low-certainty evidence relating to the effect of the intervention on maternal postnatal ARV usage (RR 0.87, 95% CI 0.61 to 1.24; n = 471; downgraded twice for risk of bias and once for imprecision; Analysis 4.2). This study also reported infant ARV/PMTCT treatment adherence as an outcome, providing low-certainty evidence that the intervention may have little or no effect (RR 1.01, 95% CI 0.98 to 1.04; n = 223; downgraded twice due to risk of bias; Analysis 4.3).

Pooled analyses of two studies carried out in Kenya provided low-certainty evidence that TCC via mobile devices may have little or no effect on increasing rates of infant HIV testing in comparison to standard care (RR 1.04, 95% CI 0.95 to 1.13; n = 838; downgraded twice due to risk of bias; Analysis 4.4) (Kassaye 2016; Odeny 2014). Repeating this analysis excluding the cluster RCT (Kassaye 2016), resulted in very little change from the original pooled estimate (RR 1.08, 95% CI 1.00 to 1.16; n = 378).

Service utilisation

One trial conducted in Kenya provided very low-certainty evidence relating to the effect of a text messaging intervention on the proportion of births occurring in health facilities, with CI encompassing both potential intervention benefit and harm (RR 0.85, 95% CI 0.62 to 1.15; n = 134; downgraded twice due to risk of bias and once due to imprecision; Analysis 4.6) (Kassaye 2016). This trial also provided low-certainty evidence that the intervention may result in a slight increase in the mean number of face-to-face or mobile communications with a healthcare worker during ANC (MD 1.50, 95% CI -0.36 to 3.36; n = 297; downgraded twice due to risk of bias; Analysis 4.7).

One trial carried out among pregnant women living with HIV in Kenya found that a text messaging intervention may increase attendance to appointments for postpartum care (RR 1.66, 95% CI 1.02 to 2.70; n = 381; moderate-certainty evidence – downgraded due to the low number of events recorded; Analysis 4.5) (Odeny 2014).

Health status and well-being

Based on low-certainty evidence from a single trial in Kenya, TCC via mobile devices may make little or no difference to neonatal health (RR for neonatal death or stillbirth: 1.12, 95% CI 0.39 to 3.28; n = 381; downgraded twice for imprecision; Analysis 4.8) (Odeny 2014). Given the wide CI consistent with both benefit and harm, there is uncertainty about what the true effect might be. Pooled analyses of two trials in Kenya also provided very low-certainty evidence, meaning we are uncertain as to whether TCC via mobile devices has an effect on the proportion of infants testing HIV positive (RR 0.54, 95% CI 0.11 to 2.56; n = 852; downgraded one level due to risk of bias and two levels due to imprecision; Analysis 4.9). The CI encompassed both benefit and harm, meaning we are uncertain as to what the true effect might be (Kassaye 2016; Odeny 2014). Repeating this analysis excluding the cluster RCT (Kassaye 2016), resulted in very little change from the original pooled estimate (RR 0.64, 95% CI 0.11 to 3.80; n = 378).

Targeted client communication via mobile devices versus non-digital targeted communication

Summary of findings 5 presents the evidence relating to the effect of TCC via mobile devices compared to non-digital targeted communication among pregnant and postpartum women living with HIV for the outcomes of maternal mortality and morbidity, neonatal and morbidity, attendance for ANC (including eMTCT care), receipt of intrapartum care, unintended consequences, neonate HIV status, and adherence to ARV therapy.

Health behaviours

No studies reported health behaviour outcomes for the comparison of TCC via mobile devices compared with non-digital targeted communication among pregnant and postpartum women living with HIV.

Service utilisation

No studies reported service utilisation outcomes for the comparison of TCC via mobile devices compared with non-digital targeted communication among pregnant and postpartum women living with HIV.

Health and well-being

No studies reported health and well-being outcomes for the comparison of TCC via mobile devices compared with non-digital targeted communication among pregnant and postpartum women living with HIV.

Targeted client communication via mobile devices versus digital non-targeted communication

[Summary of findings 6](#) presents the evidence relating to the effect of TCC via mobile devices compared to digital non-targeted communication among pregnant and postpartum women living with HIV for the outcomes of maternal mortality and morbidity combined, neonatal and morbidity combined, attendance for ANC (including eMTCT care), receipt of intrapartum care, unintended consequences, neonate HIV status, and adherence to ARV therapy.

Health behaviour

One trial conducted in Kenya among women living with HIV recruited within 24 hours of birth examined the effect of a two-weekly phone calls about PMTCT compared with standard healthcare messages. Based on this study, TCC via mobile devices may be beneficial for infant ARV/PMTCT treatment adherence at six weeks postpartum compared with digital non-targeted communication (RR 1.26, 95% CI 1.07 to 1.48; n = 150; low-certainty evidence – downgraded twice due to risk of bias with the trial at unclear or high risk of bias across all applicable domains; [Analysis 5.1](#)) ([Kebaya 2014](#)).

Service utilisation

One trial provided low-certainty evidence that the intervention may increase attendance to appointments for postpartum care at 10 weeks (RR 1.86, 95% CI 1.34 to 2.58; n = 150; downgraded twice for risk of bias; [Analysis 5.2](#)) ([Kebaya 2014](#)).

Health and well-being

No studies reported health status and well-being outcomes for the comparison of TCC via mobile devices compared with digital non-targeted communication among pregnant and postpartum women living with HIV.

Acceptability

No studies among pregnant and postpartum women living with HIV reported measures of acceptability.

Resource use

No studies among pregnant and postpartum women living with HIV reported on resource use.

Unintended consequences

No studies among pregnant and postpartum women living with HIV reported on unintended consequences or adverse outcomes.

Equity considerations

All three trials were conducted in Kenya (a lower-middle income country), and it is not possible to assess the applicability of these findings to other settings. One study specifically stated that it excluded women without a phone, who did not receive ANC, and who could not read or did not have someone to read, making it highly likely that particularly vulnerable women were unable to take part in this trial ([Odeny 2014](#)). The other two studies provided

little information on exclusion criteria, however, all trials recruited from healthcare facilities, meaning that those not accessing care will not have had the opportunity for inclusion.

Parents or carers of children aged less than five years

Targeted client communication via mobile devices versus standard care

[Summary of findings 7](#) presents the evidence relating to the effect of TCC via mobile devices compared to standard care or no intervention among parents of children aged under five years for the outcomes of child morbidity and mortality, child nutritional status, breastfeeding, clinic attendance for necessary healthcare, hygiene practices, correct treatment taken, and unintended consequences.

Health behaviours

No studies reported health behaviour outcomes for the comparison of TCC via mobile devices compared with standard care among parents of children under five years of age.

Service utilisation

Overall, for the outcome of attendance to services for necessary health care, we found low-certainty evidence for a modest intervention benefit based on the pooled results of 10 studies concerned with attendance for childhood vaccinations at six to 12 months and attendance for HIV medical appointments (among HIV-positive and HIV-exposed infants) (RR 1.21, 95% CI 1.08 to 1.34; n = 5660; downgraded for risk of bias and inconsistency; [Analysis 6.1](#)) ([Ahlers-Schmidt 2012](#); [Bangure 2015](#); [Bigna 2015](#); [Brown 2016](#); [Domek 2016](#); [Gibson 2017](#); [Haji 2016](#); [Hannan 2016](#); [Hofstetter 2015a](#); [Stockwell 2015](#)). Given the very high degree of inconsistency indicated by the I^2 value ($I^2 = 91%$), we also present subgroup analyses by type of service and timing of vaccination, described below.

Pooled analyses of five studies (conducted in the USA, Zimbabwe, Guatemala, and Kenya) indicated that TCC via mobile devices may result in a slight increase in attendance for vaccinations at six months (RR 1.14, 95% CI 1.01 to 1.28; n = 1586; low-certainty evidence – downgraded due to risk of bias and inconsistency; [Analysis 6.1](#)) ([Ahlers-Schmidt 2012](#); [Bangure 2015](#); [Domek 2016](#); [Haji 2016](#); [Hannan 2016](#)). The I^2 value was 63%, with one study reporting an point estimate indicative of reduced vaccination rates in the intervention arm ([Ahlers-Schmidt 2012](#)), and the other four studies reporting effect estimates indicative of benefit. Excluding the cluster RCT ([Haji 2016](#)) from this analysis resulted in a negligible change in the pooled effect estimate (RR 1.13, 95% CI 0.96 to 1.32).

Four studies (conducted in Nigeria, Kenya, USA) provided low-certainty evidence that TCC via mobile devices may increase attendance for vaccinations at 12 months (RR 1.24, 95% CI 1.02 to 1.52; n = 3832) ([Brown 2016](#); [Gibson 2017](#); [Hofstetter 2015a](#); [Stockwell 2015](#)) ([Analysis 6.1](#)). This body of evidence was downgraded due to risk of bias and inconsistency between individual effect estimates ($I^2 = 92%$), with two trials reporting effect estimates consistent with intervention benefit ([Brown 2016](#); [Stockwell 2015](#)), and two trials reporting little or no difference with CI encompassing no effect ([Gibson 2017](#); [Hofstetter 2015a](#)). Repeating this analysis without including the cluster RCTs ([Brown 2016](#); [Gibson 2017](#)), resulted in little change to the effect estimate, but a widening of the CI (RR 1.28, 95% CI 0.82 to 2.01).

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Pooled analyses of four RCTs (conducted in the USA, Nigeria, and Kenya) provided moderate-certainty evidence that TCC via mobile devices probably has a modest effect on increasing 'timeliness of vaccination' (vaccine receipt within a certain time period) compared with standard care (RR 1.18, 95% CI 1.04 to 1.34; $n = 2400$; downgraded due to risk of bias; [Analysis 6.2](#)) ([Ahlers-Schmidt 2012](#); [Eze 2015](#); [Gibson 2017](#); [Stockwell 2015](#)). Excluding the cluster RCT from this analysis ([Gibson 2017](#)) resulted in a negligible change in the pooled effect estimate (RR 1.19, 95% CI 0.96 to 1.46).

One trial indicated that TCC via mobile devices probably increases the attendance of HIV-positive and HIV-exposed children to medical appointments in Cameroon (RR 1.63, 95% CI 1.26 to 2.11; $n = 242$; moderate-certainty evidence – downgraded due to imprecision given the small number of events; [Analysis 6.1](#)) ([Bigna 2015](#)).

One study conducted in the USA provided very low-certainty evidence meaning we are uncertain as to whether the two-way communication phone and texting intervention had an effect on attendance to the emergency department in the six months after birth (no attendance: RR 1.32, 95% CI 1.03 to 1.70; $n = 129$; downgraded due to risk of bias, imprecision resulting from a small number of events, and indirectness (a single study conducted in a high-income country; [Analysis 6.3](#)) ([Hannan 2016](#)).

Health and well-being

No studies reported health and well-being outcomes for the comparison of TCC via mobile devices compared with standard care among parents of children under five years of age.

Targeted client communication via mobile devices versus non-digital targeted communication

[Summary of findings 8](#) presents the evidence relating to the effect of TCC via mobile devices compared to non-digital targeted communication among parents of children under five years of age for the outcomes of child morbidity and mortality, child nutritional status, breastfeeding, clinic attendance for necessary healthcare, hygiene practices, correct treatment taken, and unintended consequences.

Health behaviours

One trial carried out in India comparing the effect of providing oral health education via text messages with the provision of the same information via pamphlets suggests that the intervention may have little or no effect on children's oral health at four weeks, measured with the VPI (MD -2.10 , 95% CI -7.54 to 3.34 ; $n = 143$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 7.1](#)) ([Sharma 2011](#)).

Service utilisation

Based on one study, there was low-certainty evidence that TCC via mobile devices might result in a slight increase in attendance for vaccinations compared with non-digital targeted communication (in the form of stickers) (RR 1.13, 95% CI 1.00 to 1.28; $n = 744$; downgraded twice due to risk of bias; [Analysis 7.2](#)) ([Haji 2016](#)).

Health and well-being

No studies reported health and well-being outcomes for the comparison of TCC via mobile device compared with non-digital targeted communication among parents of children under five years of age.

Targeted client communication via mobile devices versus digital non-targeted communication

[Summary of findings 9](#) presents the evidence relating to the effect of TCC via mobile devices compared to digital non-targeted communication among parents of children under five years of age for the outcomes of child morbidity and mortality, child nutritional status, breastfeeding, clinic attendance for necessary healthcare, hygiene practices, correct treatment taken, and unintended consequences.

Health behaviour

No studies reported health behaviour outcomes for the comparison of TCC via mobile device compared with digital non-targeted communication among parents of children under five years of age.

Service utilisation

One trial in Hawaii compared vaccination reminders delivered via SMS with the control arm receiving general health information for their babies. This study found that SMS reminders may reduce attendance for vaccinations at seven months; however, the CI encompassed both benefit and harm (RR 0.63, 95% CI 0.33 to 1.20; $n = 40$; low-certainty evidence – downgraded due to risk of bias and imprecision; [Analysis 8.1](#)) ([Niederhauser 2015](#)).

Health status and well-being

No studies reported health status and well-being outcomes for the comparison of TCC via mobile device compared with digital non-targeted communication among parents of children under five years of age.

Acceptability and satisfaction

There was generally a high level of acceptability and satisfaction with the interventions assessed among parents of children aged under five years. One trial in the USA of text message reminders for childhood vaccinations interviewed 18/50 participants randomised to the intervention group, all of whom reported that they found the messages 'helpful' or 'somewhat helpful' and that they would be willing to receive text message reminders in the future ([Ahlers-Schmidt 2012](#)). Most (93%) intervention recipients in one trial of SMS immunisation reminders in Zimbabwe perceived the use of SMS as 'very beneficial' ([Bangure 2015](#)). One study of SMS vaccination reminders in Guatemala reported that 99.1% of intervention parents agreed that they would be interested in receiving text message reminders in the future and 67.5% said that they would be willing to pay for future SMS reminders ([Domek 2016](#)). One study in Kenya reported that 97.5% of participants thought the number of reminders was 'just right' (SMS vaccination reminders were sent three days and the day before each scheduled immunisation visits) ([Gibson 2017](#)). Another study involving reminders for the measles, mumps, and rubella (MMR) vaccination in the USA reported that 86.8% of participants liked the messages and 9.3% did not like them ([Hofstetter 2015a](#)). Finally another US study reported that nearly all intervention recipients (98.0%) were very satisfied or satisfied with the messages which sought to increase attendance for childhood influenza vaccination ([Stockwell 2015](#)).

Resource use

Five studies reported on the resource use of the interventions. One study reported a cost of USD 59.22 for all the messages ($n = 1368$).

(Bangure 2015). Another reported a cost of USD 0.27 per child for the project (Haji 2016). One study reported that the intervention was estimated to save between USD 51,030 and USD 104,277 in healthcare costs (Hannan 2016). Another study did not report the total costs but noted that text messaging was the most efficient intervention when both the direct costs of the intervention and staff working time were considered (Bigna 2015). Only one study reported on a cost-effectiveness analysis and found that projected cost of using SMS reminders was about 25% what it would cost to use Junior Community Health Extension Workers (CHEWs) for functional home visits in one year (Eze 2015).

Unintended consequences

No studies among parents of children aged under five years reported on unintended consequences.

Equity considerations

The trials targeting childhood immunisations were conducted in a range of high-, lower-middle-, and low-income countries. Two studies specifically included low-income families (Hannan 2016; Stockwell 2015), for example, the intervention examined by Hannan 2016 targeted low-income first-time mothers. A third study assessing text message reminders for immunisation recruited from two public health clinics in Guatemala City serving a publicly insured and low-income population (Domek 2016). Of the participants involved in the text messaging trial conducted in Kenya, 77% were unemployed and 49% had only primary level education (Haji 2016).

Sensitivity analyses

We planned to conduct a sensitivity analysis only including studies at low risk of bias (those scored at low risk of bias for the sequence generation, allocation concealment, and incomplete outcome data domains). The following trials had a low risk of bias: Yudin 2017 (pregnant and postpartum women); Odeny 2014 (pregnant and postpartum women living with HIV); and Bigna 2015; Domek 2016; Gibson 2017; Hofstetter 2015a (parents of children aged under five years). Therefore, only the comparison of TCC via mobile devices and standard care for outcome of uptake of childhood vaccinations at 12 months could undergo sensitivity analyses. As reported above, the pooled results of four studies was RR 1.28 (95% CI 0.82 to 2.01) (Brown 2016; Gibson 2017; Hofstetter 2015a; Stockwell 2015). However, when repeating this analysis pooling only the two studies at low risk of bias, the effect estimate was attenuated to RR 1.03 (95% CI 0.98 to 1.09; n = 2802) (Gibson 2017; Hofstetter 2015a).

Studies not contributing usable data

All included studies provided data that could be analysed using meta-analyses for examining the intervention effect on health behaviour, service utilisation, health and well-being, or a combination of these, meaning that conclusions relating to effectiveness have been made based on all available data from included studies.

DISCUSSION

Summary of main results

We included 27 studies in our review. For pregnant and postpartum women and parents of children aged under five years, trials were conducted in a range of low- to high-income countries. All trials

among pregnant women living with HIV were conducted in one lower-middle income country.

Pregnant and postpartum women

See [Summary of findings 1](#); [Summary of findings 2](#); and [Summary of findings 3](#).

Targeted client communication delivered by mobile device versus standard care

Women receiving TCC via mobile devices may breastfeed more in settings where breastfeeding is less common. They may go to more ANC appointments, and may use skilled birth attendants more in settings where this is less common. We do not know if TCC via mobile devices affects women's or infant morbidity or mortality because the evidence is of very low certainty.

Targeted client communication via mobile devices compared to non-digital targeted client communication or digital non-targeted communication

When compared to non-digital TCC, TCC via mobile devices may have little or no effect on breastfeeding, and may improve maternal and newborn health. We do not know what the effect is on health service use because the evidence is missing.

We do not know what the effect of TCC via mobile devices is on women's use of a skilled birth attendant when compared to digital non-targeted communication because the evidence is of very low certainty. TCC via mobile devices may make little or no difference to whether women attendance appointments for vaccination during pregnancy when compared to digital non-targeted communication. We do not know what the effect of TCC via mobile devices is on women's or babies' morbidity or mortality or breastfeeding when compared to digital non-targeted communication because the evidence is missing.

The effects of TCC via mobile devices on unintended consequences is unknown due to lack of studies. Only one study reported on resource use associated with the intervention. Satisfaction with the intervention was generally high when assessed, but few studies conducted in low-income settings reported on measures of satisfaction.

Pregnant and postpartum women living with HIV

See [Summary of findings 4](#); [Summary of findings 5](#); and [Summary of findings 6](#).

Targeted client communication delivered by mobile device versus standard care

Women who receive TCC via mobile devices may go to slightly more ANC appointments. We do not know what the effect of TCC via mobile devices is on whether women give birth in a health facility because the evidence was of very low certainty. We do not know what the effect of TCC via mobile devices is on babies' health or mortality because the evidence is of very low certainty. No studies reported on maternal mortality or morbidity. TCC via mobile devices may make little or no difference to maternal and infant adherence to ARV therapy.

Targeted client communication via mobile devices compared to non-digital targeted client communication or digital non-targeted communication

We do not know what the effect of TCC via mobile devices is on women's or babies' health or mortality, use of healthcare services during pregnancy, or whether women use skilled attendants at birth because the evidence is missing. Women who receive TCC via mobile devices may follow their baby's ARV treatment better than women who receive digital non-targeted communication.

For pregnant and postpartum women living with HIV, we are uncertain of the satisfaction and resource use associated with receiving TCC via mobile devices, or the effects on unintended consequences, due to lack of evidence.

Parents or carers of children aged under five years

See [Summary of findings 7](#); [Summary of findings 8](#); and [Summary of findings 9](#).

Targeted client communication delivered by mobile device versus standard care

We do not know what the effect of parental receipt of TCC via mobile devices is on breastfeeding, children's health or nutrition, whether they received the right treatments, or parents' hygiene practices because the evidence is missing. Parents who receive TCC via mobile devices may take their children to more healthcare services, such as vaccination appointments; however, the results varied widely between studies.

Targeted client communication via mobile devices compared to non-digital targeted client communication or digital non-targeted communication

We do not know what the effect of TCC via mobile devices is on breastfeeding, children's health or nutrition, or whether they received the correct treatments, because the evidence is missing. Parents who receive TCC via mobile devices may take their children to more vaccination appointments compared to parents who receive non-digital targeted communication although the effects vary. There is little or no difference observed between these groups in terms of oral hygiene practices. Parents who receive TCC via mobile devices may take their children to fewer vaccination appointments than parents who receive digital non-targeted communication, but it is possible that the intervention makes little or no difference.

The effects of TCC via mobile devices on unintended consequences is unknown due to lack of studies. Acceptability and satisfaction with the intervention was high in the six studies that reported these outcomes. Costs were low in the five studies that reported on intervention costs.

Overall completeness and applicability of evidence

This review includes evidence from 27 studies, 11 of which were conducted among pregnant and postpartum women, three among pregnant and postpartum women living with HIV, and 13 among parents of children aged under five years. In collating those trials relating to the same population, employing the same comparison groups, and measuring the same outcomes, these specific bodies of evidence were generally made up of only a small number of studies, thereby limiting the extent to which conclusions can be made.

With the exception of trials among pregnant and postpartum women living with HIV, all of which were carried out in Kenya, there was generally a varied geographical spread for where studies were conducted, with representation of low- to high-income countries. Furthermore, the trials recruited from a mix of community and healthcare settings.

No trials targeted or measured correct treatment of childhood illness or child nutritional outcomes. No trials reported measuring unintended consequences or adverse events. However, HIV and reproduction can be sensitive topics and there is the potential for harm resulting from interventions concerned with these issues. In the accompanying review focusing on reproductive and sexual health, one trial among women living with HIV reported one participant withdrew due to her undisclosed HIV status being compromised ([Palmer in preparation](#)).

Quality of the evidence

Using GRADE methodology, we assessed the certainty of the evidence for all outcomes. Among the pregnant and postpartum population, there was low or very low-certainty evidence relating to the effect of TCC via mobile devices for most outcomes. There was moderate-certainty evidence for only three outcomes among this population, generally based on the results of a single trial. For pregnant and postpartum women living with HIV, the evidence for all outcomes except one were considered of low or very low certainty. Based on trials conducted among parents of children aged under five years, for most outcomes the evidence was of low or very low certainty. Across all three populations, the most common reasons for downgrading were risk of bias of the studies and imprecision of the effect estimates.

Given the nature of the interventions, participant blinding was generally unfeasible. The trials among pregnant and postpartum women primarily depended on self-reported outcome measures. Therefore, among these studies there is risk of response bias which could result in an overestimation of intervention benefit. Most trials among parents of children aged under five years concerned with increasing uptake of childhood vaccination were able to use objective outcome measures based on clinic attendance recording systems, meaning that for these studies, having unblinded intervention recipients would be unlikely to bias the effect estimates recorded. Few other trials used objective outcomes, which would be less subject to bias than self-reported measures.

Potential biases in the review process

There were limited resources to follow-up with authors of reports that did not provide sufficient information for data extraction/risk of bias assessments. Furthermore, publication bias, whereby trials with positive findings are more likely to be published, may have biased the selection of included studies in this review. However, efforts were made to overcome this through searching clinical trial registries for prospectively registered trials.

Agreements and disagreements with other studies or reviews

As with other communication and behaviour change interventions, a challenge in synthesising the results of TCC interventions delivered by mobile device is the heterogeneity of interventions, ranging from reminders to educational and more complex behaviour change support. As evidenced by the heterogeneity

of some pooled estimates in our review and similar reviews (Amankwaa 2018; McFadden 2017), this can result in real differences in effects of interventions even when they use the same delivery mechanisms and target the same outcomes.

There was uncertain evidence relating to the effect of TCC via mobile devices for improving postpartum contraception use. The CI was wide so modest benefits similar to those reported in other Cochrane Reviews of contraception interventions delivered by mobile phone for general populations of women or women postabortion cannot be excluded (Palmer in preparation; Smith 2015).

Our finding that TCC delivered by mobile devices may improve exclusive breastfeeding at three months in settings where exclusive breastfeeding rates are modest is consistent with one Cochrane Review which found that breastfeeding support is effective in increasing exclusive breastfeeding and the duration of breastfeeding. We note, however, that the trial of TCC included in our review was delivered by mobile phone calls and the Cochrane Review found that face-to-face support was more effective than telephone support (McFadden 2017).

This review showed TCC delivered by mobile devices may increase abstinence from smoking in pregnancy, but evidence is low certainty. These findings are consistent with the effects of TCC delivered by mobile phone which doubles continuous abstinence from smoking among general populations of smokers willing to make a quit attempt (Free 2011; Palmer 2018b; Whittaker 2016). Effective smoking cessation support delivered by mobile phone includes a range of behavioural change techniques, and the behaviour change techniques found in effective psychosocial support for smoking cessation in pregnant women have also been described (Lorenatto 2012). Since our search was completed, one trial of TCC by mobile phone for smoking cessation in pregnant women compared to digital non-targeted communication reported an increase in validated no smoking in the last seven days at three months (OR 1.47, 95% CI 1.21 to 1.78) (Abroms 2017a) and one recent systematic review including the Abroms 2017a trial reported an increase in smoking cessation during pregnancy (pooled OR 1.59, 95% CI 1.07 to 2.38) (Griffiths 2018). Our review findings are also consistent with the findings of one systematic review of health education and psychosocial interventions, not specifically delivered by mobile devices, for smoking cessation during pregnancy which found evidence of a beneficial intervention effect (Chamberlain 2017).

Our finding of a lack of effect of TCC via mobile devices for alcohol reduction in pregnancy is consistent with one Cochrane Review of psychological and education interventions for reducing alcohol intake, not specifically delivered by mobile devices, which found no intervention effects for most outcomes (Stade 2009).

The effect of TCC via mobile devices for improving antenatal and postnatal adherence to ARV therapy among women living with HIV was uncertain. The effects on ARV/PMTCT adherence in infants were mixed. One previous Cochrane Review examining text messaging for adherence to ARV therapy in populations of all ages reported evidence of beneficial effects from two trials (Horvath 2012), but further trials of ARV therapy adherence interventions delivered by mobile phone have been published since 2012, with mixed results reported in more recent systematic reviews (Quintana 2018; Shah 2019). One systematic review reported an increase in adherence

for adults based on a pooled analysis of interventions delivered by SMS, but uncertain effects of interventions delivered by voice message (Amankwaa 2018).

The finding that TCC via mobile devices may improve service utilisation, such as attendance to antenatal and postnatal care appointments among pregnant and postpartum women, and attendance for infant HIV medical appointments, is in accordance with our findings relating to other populations (Palmer in preparation), and previous research findings that mobile phone-based reminders improve healthcare appointment attendance (Free 2013b; Guro-Urganci 2013).

Some feasibility studies have shown promise for TCC delivered by mobile devices altering nutrition in infants or children (Singh 2014). However, we did not identify any RCTs of nutritional interventions or trials reporting infant or child nutrition outcomes that met our inclusion criteria.

Our findings that TCC via mobile devices among parents of children aged under five years may have modest effects on increasing attendance for childhood immunisations and necessary health care is consistent with the findings of a previous review of mobile phone-based interventions (Free 2013a), and a Cochrane Review examining face-to-face interventions for parents about childhood vaccinations, which also found evidence for small intervention benefits (Kaufman 2018). However, in our review, the effects varied in different studies. This could be due to bias, as both of the trials that found a positive effect were at unclear risk of bias for random sequence generation. It is also possible that there was a ceiling effect in the Gibson 2017 trial due to the high uptake of vaccination in the control arm (82%) leaving less scope for extra improvement to be demonstrated. It may also be that for some vaccinations remembering an appointment is not the main barrier to uptake, so simple reminders would not be expected to improve attendance. The Hofstetter 2015a trial in the USA was targeting MMR, which in the context of vaccine concerns might be intractable to improvement using reminder messages only. Educational messages targeting parental concerns about the MMR vaccine could be more relevant.

AUTHORS' CONCLUSIONS

Implications for practice

We cannot make strong recommendations for implementation of targeted client communication (TCC) delivered by mobile devices on the basis of this review as the certainty of the evidence was primarily very low or low.

TCC interventions delivered by mobile devices, such as appointment reminders have already been implemented in many settings. Where healthcare providers and policy makers are considering implementation, we suggest prioritising interventions for which evidence is suggestive of an improvement in outcomes, such as TCC delivered by mobile devices for general populations of pregnant women for antenatal care attendance, skilled attendance at birth, and for childhood vaccinations. However, careful consideration should be given to the costs and resources needed in conjunction with the limited strength of evidence for gains in service use and health behaviours. Few studies in this review reported on costs, but those that did so generally indicated costs to be low. Furthermore, the interventions delivered

in trials which observed evidence for a small increase in attendance for vaccinations (parents of children aged under five years) compared to standard care, consisted of simple reminders which may be more straightforward to implement than more complex interventions. The text messaging intervention evaluated by [Odeny 2014](#) consisted of just 14 messages sent to pregnant women living with HIV from 28 weeks' gestation to six weeks postdelivery, but seemed to result in gains in relation to attendance for the postpartum appointment, and infant HIV testing. In other areas of health behaviour change, such as smoking cessation, effective TCC delivered by mobile devices has been shown to be highly cost effective ([Guerriero 2013](#)). The majority of TCC interventions in this review were delivered by text message (SMS). The World Health Organization and the International Telecommunication Union have developed guidance on implementing SMS-based smoking cessation support at scale, and much of this guidance would apply to implementing any SMS health platform ([WHO/ITU 2015](#)). Nonetheless, setting up, maintaining and monitoring SMS platforms, and delivery requires resources and incurs ongoing costs, which can represent a considerable burden and challenge in some settings. Furthermore, the findings of one complementary synthesis of qualitative research which explored clients' perceptions and experiences of targeted communication via mobile devices on topics related to reproductive, maternal, newborn, child, and adolescent health (RMNCAH) ([Ames 2019](#)), highlights continuing barriers to the widespread implementation of TCC via mobile devices. Despite the high rate of mobile phone ownership globally, use of these devices can still be limited by poor access to mobile networks and the internet, phones breaking, the inability to afford airtime, and the inability to charge the device. Given that there are contexts in which uninterrupted access to mobile devices cannot be guaranteed, it may be premature to roll out TCC via mobile devices if it is at the expense of other channels of communication.

For some outcomes, the findings suggested intervention benefits may be gained only in settings where the behaviour targeted is relatively uncommon and there is greater scope for improvement. For example, while TCC via mobile devices had little or no effect in a low-risk setting where the majority of births already occurred with a skilled attendant present, there was evidence that the intervention increased skilled birth attendance in a moderate-risk urban setting where two-thirds of births occurred with skilled attendance (but not in the rural area included in the same study). Therefore, background levels of health service use and behaviours (and existing health facility infrastructure), should be considered prior to implementation. While the majority of comparisons in the review were between TCC via mobile devices and standard care/no intervention, consideration should also be given to the apparent lack of intervention benefit when the comparison group is in receipt of something quite similar to the intervention. For example, the trial by [Niederhauser 2015](#) examined the effect of a TCC mobile device-delivered intervention involving childhood vaccination reminders at four weeks and two weeks prior to the routine vaccination appointments, compared with the control group who received well baby messages. However, the study report states that the participants in both groups were also receiving the "routine reminders that they receive from their healthcare providers," therefore, the finding of no additional benefit of the reminder messaging intervention, and perhaps even a slight negative effect, might be considered unsurprising. This highlights that an understanding of what is delivered as part of existing

medical practice, which may vary between contexts, is important to prevent misplaced interventions which result in few or no gains.

Healthcare providers and policy makers should be especially cautious regarding the implementation of interventions specifically targeting pregnant women living with HIV. While the evidence is of low or very low certainty, the confidence intervals around the effect estimates suggest that TCC delivered by mobile devices may have little benefit or no effect. Due to the sensitivity and stigma associated with HIV status, it is essential that such interventions are designed so that content and delivery ensures confidentiality of sensitive material. In contexts where women's access to phones is controlled by others, or where phones are shared, it may not be feasible for content to remain private. In such circumstances it may be better to include HIV-positive pregnant women in mobile phone interventions for general populations of pregnant women, such as for attendance for antenatal care, rather than targeting them in isolation on the basis of their HIV status. This is consistent with the findings of one synthesis of qualitative research concerned with perceptions and experiences of TCC via mobile devices on topics related to RMNCAH ([Ames 2019](#)). Practical considerations for messaging concerned with stigmatised or personal health conditions included the use of neutral language, and tailoring the content, timing, and frequency of the messages ([Ames 2019](#)).

No studies across the three populations in this review reported on unintended consequences or adverse events. However, given that reproduction can be a sensitive topic and HIV is subject to considerable stigma, there is the potential for harm resulting from interventions concerned with these issues. For example, in the accompanying review focusing on reproductive and sexual health, a trial among women living with HIV reported that one participant withdrew due to her undisclosed HIV status being compromised. Furthermore, globally one in three women has experienced intimate partner violence in their lifetime and pregnancy is a time when domestic violence can escalate ([WHO 2013](#)). There is the potential that TCC on sensitive topics could result in further escalations in abuse for women in abusive relationships. Consequently, those implementing interventions should consider how the risks of such potential unintended consequences can be minimised and monitored.

Issues relating to language and literacy have been highlighted in this review as potential barriers in achieving equity of access and benefit from communication-based interventions. This is also a finding of the qualitative synthesis of clients' perceptions experiences with such interventions ([Ames 2019](#)), in which language, literacy, techno-literacy, or a combination of these were raised as potential limiting factors for accessing these types of interventions. Based on this, some participants reported a preference for voice calls in which they could request clarifying information. Where interventions are being implemented, this should be accompanied by monitoring and evaluation of impact (including both benefits and harms), equity, and associated costs.

Implications for research

- The effect of TCC delivered by mobile devices on health service use (antenatal care, delivery in a health facility) and vaccinations at six and 12 months requires further assessment. The vaccination interventions in this review primarily provided reminders. Further TCC interventions, which

- also target attitudinal barriers to childhood vaccinations, should be evaluated.
- Given the somewhat equivocal evidence relating to the effects of TCC delivered by mobile devices for pregnant women living with HIV, future research should explore whether carefully designed interventions can improve outcomes in this population, and the contexts in which this can occur.
 - We found no trials of TCC delivered by mobile devices that targeted treatment of childhood illness, and infant and child nutrition. Therefore, research is needed to evaluate whether well-designed TCC interventions targeting these behaviours can be effective.
 - Future research should ensure potential unintended consequences and adverse events are considered in the design of the intervention and evaluation, so that risks are minimised and any such events that do occur are adequately captured.
 - Researchers should ensure that interventions are adequately described in sufficient detail for them to be replicated, for example, with reference to guidance on reporting interventions ([Agarwal 2016](#); [Hoffmann 2014](#)).
 - Both trials and implementation research should evaluate equity in access to, and the effects of, TCC delivered by mobile devices for maternal, neonatal, and child health.
 - Where interventions have been shown to be effective, cost-effectiveness analysis of TCC delivered by mobile device for maternal, neonatal, and child health is needed.
- Further high-quality trials are warranted to evaluate the effects of TCC by mobile devices on behaviours such as postpartum contraception use and breastfeeding. Given that lay support is associated with increased exclusive breastfeeding and duration of breastfeeding, trials of lay support for breastfeeding delivered by mobile devices could be considered ([McFadden 2017](#)).

ACKNOWLEDGEMENTS

We thank the editors and staff of Cochrane Consumers and Communication, particularly Bronwen Merner, Ann Jones, and Rebecca Ryan for their input into the protocol.

We are grateful to the Guideline Development Group and External Review Group of the *WHO Guideline on digital interventions for health system strengthening* for their constructive feedback in formulating the guiding questions for this systematic review. Cochrane Response, an evidence services unit operated by Cochrane provided screening and data extraction services for this review. This work was funded by the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a cosponsored program executed by the World Health Organization (WHO). We thank John Evers for designing and running the search strategies for this review.

REFERENCES

References to studies included in this review

Ahlers-Schmidt 2012 {published data only}

Ahlers-Schmidt CR, Chesser AK, Nguyen T, Brannon J, Hart TA, Williams KS, et al. Feasibility of a randomized controlled trial to evaluate Text Reminders for Immunization Compliance in Kids (TRICKS). *Vaccine* 2012;**30**(36):5305-9.

Bangure 2015 {published data only}

Bangure D, Chirundu D, Gombe N, Marufu T, Mandozana G, Tshimanga M, et al. Effectiveness of short message services reminder on childhood immunization programme in Kadoma, Zimbabwe – a randomized controlled trial, 2013. *BMC Public Health* 2015;**15**:137.

Bigna 2015 {published data only}

Bigna JJ, Noubiap JJ, Kouanfack C, Plottel CS, Koulla-Shiro S. Effect of mobile phone reminders on follow-up medical care of children exposed to or infected with HIV in Cameroon (MORE CARE): a multicentre, single-blind, factorial, randomised controlled trial. *Lancet Infectious Diseases* 2014;**14**(7):600-8.

Bigna JJ, Noubiap JJ, Plottel CS, Kouanfack C, Koulla-Shiro S. Barriers to the implementation of mobile phone reminders in pediatric HIV care: a pre-trial analysis of the Cameroonian MORE CARE study. *BMC Health Services Research* 2014;**14**:523.

Bigna JJ, Noubiap JJ, Plottel CS, Kouanfack C, Koulla-Shiro S. Factors associated with non-adherence to scheduled medical follow-up appointments among Cameroonian children requiring HIV care: a case-control analysis of the usual-care group in the MORE CARE trial. *Infectious Diseases of Poverty* 2014;**3**(1):44.

* Bigna JJ. Automated text message reminders to promote good health. *Lancet Infectious Diseases* 2015;**15**(1):19-20.

Mbuagbaw L. Mobile phone reminders for paediatric HIV follow-up care. *Lancet Infectious Diseases* 2014;**14**(7):540-1.

Brown 2016 {published data only}

Brown VB, Oluwatosin OA, Akinyemi JO, Adeyemo AA. Effects of community health nurse-led intervention on childhood routine immunization completion in primary health care centers in Ibadan, Nigeria. *Journal of Community Health* 2016;**41**(2):265-73.

Domek 2016 {published data only}

Domek GJ, Contreras-Roldan IL, O'Leary ST, Bull S, Furniss A, Kempe A, et al. SMS text message reminders to improve infant vaccination coverage in Guatemala: a pilot randomized controlled trial. *Vaccine* 2016;**34**(21):2437-43.

Evans 2014 {published data only}

Evans WD, Wallace B, Szekely D, Nielsen P, Murray E, Abrams L, et al. Initial outcomes from a 4-week follow-up study of the text4baby program in the military women's population: randomized controlled trial. *Journal of Medical Internet Research* 2014;**16**(5):e131.

Eze 2015 {published data only}

Eze GU, Adeleye OO. Enhancing routine immunization performance using innovative technology in an urban area of Nigeria. *West African Journal of Medicine* 2015;**34**(1):3-10.

Gibson 2017 {published data only}

Gibson DG, Ochieng B, Kagucia EW, Were J, Hayford K, Moulton LH, et al. Mobile phone-delivered reminders and incentives to improve childhood immunisation coverage and timeliness in Kenya (M-SIMU): a cluster randomised controlled trial. *Lancet Global Health* 2017;**5**(4):e428-e38.

Haji 2016 {published data only}

Haji A, Lowther S, Ngan'ga Z, Gura Z, Tabu C, Sandhu H, et al. Reducing routine vaccination dropout rates: evaluating two interventions in three Kenyan districts, 2014. *BMC Public Health* 2016;**16**:152.

Hannan 2016 {published data only}

Hannan J, Brooten D, Page T, Galindo A, Torres M. Low-income first-time mothers: effects of APN follow-up using mobile technology on maternal and infant outcomes. *Global Pediatric Health* 2016;**3**:2333794X16660234.

Hofstetter 2015a {published data only}

Hofstetter AM, DuRivage N, Vargas CY, Camargo S, Vawdrey DK, Fisher A, et al. Text message reminders for timely routine MMR vaccination: a randomized controlled trial. *Vaccine* 2015;**33**(43):5741-6.

Jareethum 2008 {published data only}

Jareethum R, Titapant V, Chantra T, Sommai V, Chuenwattana P, Jirawan C. Satisfaction of healthy pregnant women receiving short message service via mobile phone for prenatal support: a randomized controlled trial. *Journal of the Medical Association of Thailand* 2008;**91**(4):458-63.

Joshi 2015 {published data only}

Joshi S, Patil N, Hegde A. Impact of mHealth initiative on utilization of antenatal care services in rural Maharashtra, India. *Indian Journal of Maternal and Child Health* 2015;**17**(2):1-7.

Kamau-Mbuthia 2013 {published data only}

* Kamau-Mbuthia E, Mbugua S, Webb G, Kalungu S, Sarange C, Lou W, et al. Cell phone based peer counseling to support exclusive breastfeeding is associated with more frequent help and decreased breastfeeding problems. *Annals of Nutrition & Metabolism* 2013;**63**:196-7.

Webb G, Kamau-Mbuthia E, Mbugua S, Kalungu S, Sarange C, Lou W, et al. Infant medication, illness and growth in a randomized controlled trial of exclusive breastfeeding support in Kenya. *Annals of Nutrition & Metabolism* 2013;**63**:752.

Kassaye 2016 {published data only}

Kassaye SG, Ong'ech J, Sirengo M, Kose J, Matu L, McOdida P, et al. Cluster-randomized controlled study of SMS text messages for prevention of mother-to-child transmission of HIV in rural Kenya. *AIDS Research & Treatment* 2016;**2016**:1289328.

Kebaya 2014 {published data only}

* Kebaya L, Nduati R, Wamalwa D, Kariuki N, Bashir A. Efficacy of mobile phone use on adherence to nevirapine prophylaxis and retention in care among the HIV-exposed infants in PMTCT: a randomised controlled trial. *Archives of Disease in Childhood* 2014;**99**:A329.

Kebaya LM, Wamalwa D, Kariuki N, Admani B, Nduati RW. Efficacy of mobile phone use on adherence to nevirapine prophylaxis and retention in care among HIV-exposed infants. *Topics in Antiviral Medicine* 2015;**23**:407.

Lund 2012 {published data only}

* Lund S, Hemed M, Nielsen BB, Said A, Said K, Makungu MH, et al. Mobile phones as a health communication tool to improve skilled attendance at delivery in Zanzibar: a cluster-randomised controlled trial. *BJOG* 2012;**119**(10):1256-64.

Lund S, Nielsen BB, Hemed M, Boas IM, Said A, Said K, et al. Mobile phones improve antenatal care attendance in Zanzibar: a cluster randomized controlled trial. *BMC Pregnancy & Childbirth* 2014;**14**:29.

Lund S, Nielsen BB, Hemed M, Said A, Said K, Makungu MH, et al. Mobile phones as a health communication tool to improve maternal and perinatal health in Zanzibar: a cluster randomised controlled trial. *Tropical Medicine & International Health* 2013;**18**:22.

Lund S, Rasch V, Hemed M, Boas IM, Said A, Said K, et al. Mobile phone intervention reduces perinatal mortality in Zanzibar: secondary outcomes of a cluster randomized controlled trial. *JMIR mHealth and uHealth* 2014;**2**(1):e15.

Maslowky 2016 {published data only}

Maslowky J, Frost S, Hendrick CE, Trujillo Cruz FO, Merajver SD. Effects of postpartum mobile phone-based education on maternal and infant health in Ecuador. *International Journal of Gynaecology & Obstetrics* 2016;**134**(1):93-8.

McConnell 2016 {published data only}

McConnell M, Ettenger A, Rothschild CW, Muigai F, Cohen J. Can a community health worker administered postnatal checklist increase health-seeking behaviors and knowledge?: evidence from a randomized trial with a private maternity facility in Kiambu County, Kenya. *BMC Pregnancy Childbirth* 2016;**16**(1):136.

Moniz 2013 {published data only}

Moniz MH, Hasley S, Meyn LA, Beigi RH. Improving influenza vaccination rates in pregnancy through text messaging: a randomized controlled trial. *Obstetrics & Gynecology* 2013;**121**(4):734-40.

Naughton 2017 {published data only}

Cooper S, Foster K, Naughton F, Leonardi-Bee J, Sutton S, Ussher M, et al. Pilot study to evaluate a tailored text message intervention for pregnant smokers (MiQuit): study protocol for a randomised controlled trial. *Trials* 2015;**16**:29.

* Naughton F, Cooper S, Foster K, Emery J, Leonardi-Bee J, Sutton S, et al. Large multi-centre pilot randomized controlled

trial testing a low-cost, tailored, self-help smoking cessation text message intervention for pregnant smokers (MiQuit). *Addiction* 2017;**112**(7):1238-49.

Niederhauser 2015 {published data only}

Niederhauser V, Johnson M, Tavakoli AS. Vaccines4Kids: assessing the impact of text message reminders on immunization rates in infants. *Vaccine* 2015;**33**(26):2984-9.

Odeny 2014 {published data only}

Odeny TA, Bukusi EA, Cohen CR, Yuhas K, Camlin CS, McClelland RS. Texting improves testing: a randomized trial of two-way SMS to increase postpartum prevention of mother-to-child transmission retention and infant HIV testing. *AIDS* 2014;**28**(15):2307-12.

Omole 2018 {published data only}

Omole O, Ijadunola MY, Olotu E, Omotoso O, Bello B, Awoniran O, et al. The effect of mobile phone short message service on maternal health in south-west Nigeria. *International Journal of Health Planning and Management* 2018;**33**(1):155-70.

Sharma 2011 {published data only}

Sharma R, Hebbal M, Ankola AV, Murugabupathy V. Mobile-phone text messaging (SMS) for providing oral health education to mothers of preschool children in Belgaum City. *Journal of Telemedicine & Telecare* 2011;**17**(8):432-6.

Stockwell 2015 {published data only}

Stockwell MS, Hofstetter AM, DuRivage N, Barrett A, Fernandez N, Vargas CY, et al. Text message reminders for second dose of influenza vaccine: a randomized controlled trial. *Pediatrics* 2015;**135**(1):e83-91.

Yudin 2017 {published data only}

Yudin MH, Mistry N, De Souza LR, Besel K, Patel V, Blanco Mejia S, et al. Text messages for influenza vaccination among pregnant women: a randomized controlled trial. *Vaccine* 2017;**35**(5):842-8.

References to studies excluded from this review
Atukunda 2017 {published data only}

Atukunda EC, Musiimenta A, Musinguzi N, Wyatt MA, Ashaba J, Ware NC, et al. Understanding patterns of social support and their relationship to an ART adherence intervention among adults in rural Southwestern Uganda. *AIDS and Behavior* 2017;**21**(2):428-40.

Bracken 2014 {published data only}

Bracken H, Lohr PA, Taylor J, Morroni C, Winikoff B. RU OK? The acceptability and feasibility of remote technologies for follow-up after early medical abortion. *Contraception* 2014;**90**(1):29-35.

Broberg 2013 {published data only}

Broberg G, Jonasson JM, Ellis J, Gyrd-Hansen D, Anjemark B, Glantz A, et al. Increasing participation in cervical cancer screening: telephone contact with long-term non-attendees in Sweden. Results from RACOMIP, a randomized controlled trial. *International Journal of Cancer* 2013;**133**(1):164-71.

Carlsen 2013 {published data only}

Carlsen EM, Kyhnaeb A, Renault KM, Cortes D, Michaelsen KF, Pryds O. Telephone-based support prolongs breastfeeding duration in obese women: a randomized trial. *American Journal of Clinical Nutrition* 2013;**98**(5):1226-32.

Collier 2005 {published data only}

Collier AC, Ribaudo H, Mukherjee AL, Feinberg J, Fischl MA, Chesney M, et al. A randomized study of serial telephone call support to increase adherence and thereby improve virologic outcome in persons initiating antiretroviral therapy. *Journal of Infectious Diseases* 2005;**192**(8):1398-406.

Gallegos 2014 {published data only}

Gallegos D, Russell-Bennett R, Previte J, Parkinson J. Can a text message a week improve breastfeeding? *BMC Pregnancy and Childbirth* 2014;**14**(1):374.

Haberer 2016 {published data only}

Haberer JE, Musiimenta A, Atukunda EC, Musinguzi N, Wyatt MA, Ware NC, et al. Short message service (SMS) reminders and real-time adherence monitoring improve antiretroviral therapy adherence in rural Uganda. *AIDS* 2016;**30**(8):1295-300.

Hashemian 2015 {published data only}

Hashemian TS, Kritiz-Silverstein D, Baker R. Text2floss: the feasibility and acceptability of a text messaging intervention to improve oral health behavior and knowledge. *Journal of Public Health Dentistry* 2015;**75**(1):34-41.

Herring 2016 {published data only}

Herring SJ, Cruice JF, Bennett GG, Rose MZ, Davey A, Foster GD. Preventing excessive gestational weight gain among African American women: a randomized clinical trial. *Obesity* 2016;**24**(1):30-6.

Hofstetter 2015b {published data only}

Hofstetter AM, Vargas CY, Camargo S, Holleran S, Vawdrey DK, Kharbanda EO, et al. Impacting delayed pediatric influenza vaccination: a randomized controlled trial of text message reminders. *American Journal of Preventive Medicine* 2015;**48**(4):392-401.

Irons 2015 {published data only}

Irons M, Tomaszewski K, Buchanan CR, Trent M. Understanding adolescent nonresponsiveness to text messages: lessons from the Depotext trial. *Journal of Urban Health* 2015;**92**(3):502-12.

Jimenez 2017 {published data only}

Jimenez ME, DuRivage NE, Bezpalko O, Suh A, Wade R, Blum NJ, et al. A pilot randomized trial of a video patient decision aid to facilitate early intervention referrals from primary care. *Clinical Pediatrics* 2017;**56**(3):268-77.

Kofinas 2014 {published data only}

Kofinas JD, Varrey A, Sapra KJ, Kanj RV, Chervenak FA, Asfaw T. Adjunctive social media for more effective contraceptive counseling: a randomized controlled trial. *Obstetrics & Gynecology* 2014;**123**(4):763-70.

Lau 2013 {published data only}

Lau JT, Tsui HY, Lau MM. A pilot clustered randomized control trial evaluating the efficacy of a network-based HIV peer-education intervention targeting men who have sex with men in Hong Kong, China. *AIDS Care* 2013;**25**(7):812-9.

Lau 2014 {published data only}

Lau YK, Cassidy T, Hacking D, Brittain K, Haricharan HJ, Heap M. Antenatal health promotion via short message service at a midwife obstetrics unit in South Africa: a mixed methods study. *BMC Pregnancy and Childbirth* 2014;**14**(1):284.

Lewis 2012 {published data only}

Lewis BA, Gjerdingen DK, Avery MD, Guo H, Sirard JR, Bonikowske AR, et al. Examination of a telephone-based exercise intervention for the prevention of postpartum depression: design, methodology, and baseline data from the healthy mom study. *Contemporary Clinical Trials* 2012;**33**(6):1150-8.

Maduka 2013 {published data only}

Maduka O, Tobin-West CI. Adherence counseling and reminder text messages improve uptake of antiretroviral therapy in a tertiary hospital in Nigeria. *Nigerian Journal of Clinical Practice* 2013;**16**(3):302-8.

Mauriello 2016 {published data only}

Mauriello LM, Van Marter DF, Umanson CD, Castle PH, de Aguiar EL. Using mHealth to deliver behavior change interventions within prenatal care at community health centers. *American Journal of Health Promotion* 2016;**30**(7):554-62.

Milani 2015 {published data only}

Milani HS, Azargashb E, Beyraghi N, Defaie S, Asbaghi T. Effect of telephone-based support on postpartum depression: a randomized controlled trial. *International Journal of Fertility & Sterility* 2015;**9**(2):247-53.

Moore 2013 {published data only}

Moore DJ, Montoya JL, Blackstone K, Rooney A, Gouaux B, Georges S, et al. Preliminary evidence for feasibility, use, and acceptability of individualized texting for adherence building for antiretroviral adherence and substance use assessment among HIV-infected methamphetamine users. *AIDS Research and Treatment* 2013;**2013**:585143.

Moore 2015 {published data only}

Moore DJ, Poquette A, Casaletto KB, Gouaux B, Montoya JL, Posada C, et al. Individualized texting for adherence building (iTAB): improving antiretroviral dose timing among HIV-infected persons with co-occurring bipolar disorder. *AIDS and Behavior* 2015;**19**(3):459-71.

Murthy 2019 {published data only}

Murthy N, Chandrasekharan S, Prakash MP, Kaonga NN, Peter J, Ganju A, et al. The impact of an mHealth voice message service (mMitra) on infant care knowledge, and practices among low-income women in India: findings from a pseudo-randomized controlled trial. *Maternal Child Health Journal* 2019;**23**(12):1658-69.

Mwapasa 2017 {published data only}

Mwapasa V, Joseph J, Tchereni T, Jousset A, Gunda A. Impact of mother–infant pair clinics and short-text messaging service (SMS) reminders on retention of HIV-infected women and HIV-exposed infants in eMTCT care in Malawi: a cluster randomized trial. *Journal of Acquired Immune Deficiency Syndromes* 2017;**75**:S123-31.

Oakley-Girvan 2016 {published data only}

Oakley-Girvan I, Londono C, Canchola A, Watkins Davis S. Text messaging may improve abnormal mammogram follow-up in Latinas. *Oncology Nursing Forum* 2016;**43**(1):36-43.

Patel 2014 {published data only}

Patel A, Stern L, Unger Z, Debevec E, Roston A, Hanover R, et al. Staying on track: a cluster randomized controlled trial of automated reminders aimed at increasing human papillomavirus vaccine completion. *Vaccine* 2014;**32**(21):2428-33.

Peitzmeier 2016 {published data only}

Peitzmeier SM, Khullar K, Potter J. Effectiveness of four outreach modalities to patients overdue for cervical cancer screening in the primary care setting: a randomized trial. *Cancer Causes & Control* 2016;**27**(9):1081-91.

Pérez-Ferre 2010 {published data only}

Pérez-Ferre N, Galindo M, Fernández MD, Velasco V, Runkle I, de la Cruz MJ, et al. The outcomes of gestational diabetes mellitus after a telecare approach are not inferior to traditional outpatient clinic visits. *International Journal of Endocrinology* 2010;**2010**:386941.

Pollak 2014 {published data only}

Pollak KI, Alexander SC, Bennett G, Lyna P, Coffman CJ, Bilheimer A, et al. Weight-related SMS texts promoting appropriate pregnancy weight gain: a pilot study. *Patient Education and Counseling* 2014;**97**(2):256-60.

Prieto 2016 {published data only}

Prieto JT, Zuleta C, Rodríguez JT. Modeling and testing maternal and newborn care mHealth interventions: a pilot impact evaluation and follow-up qualitative study in Guatemala. *Journal of the American Medical Informatics Association* 2016;**24**(2):352-60.

Rampersaud 2016 {published data only}

Rampersaud GC, Sokolow A, Gruspe A, Colee JC, Kauwell GP. Folate/folic acid knowledge, intake, and self-efficacy of college-aged women: impact of text messaging and availability of a folic acid-containing supplement. *Journal of American College Health* 2016;**64**(6):460-8.

Rand 2015 {published data only}

Rand CM, Brill H, Albertin C, Humiston SG, Schaffer S, Shone LP, et al. Effectiveness of centralized text message reminders on human papillomavirus immunization coverage for publicly insured adolescents. *Journal of Adolescent Health* 2015;**56**(5):S17-20.

Rand 2017 {published data only}

Rand CM, Vincelli P, Goldstein NP, Blumkin A, Szilagyi PG. Effects of phone and text message reminders on completion of the human papillomavirus vaccine series. *Journal of Adolescent Health* 2017;**60**(1):113-9.

Reeder 2014 {published data only}

Reeder JA, Joyce T, Sibley K, Arnold D, Altindag O. Telephone peer counseling of breastfeeding among WIC participants: a randomized controlled trial. *Pediatrics* 2014;**134**(3):e700.

Reid 2014 {published data only}

Reid MJ, Dhar SI, Cary M, Liang P, Thompson J, Gabaitiri L, et al. Opinions and attitudes of participants in a randomized controlled trial examining the efficacy of SMS reminders to enhance antiretroviral adherence: a cross-sectional survey. *Journal of Acquired Immune Deficiency Syndromes* 2014;**65**(2):e86-8.

Richman 2016 {published data only}

Richman AR, Maddy L, Torres E, Goldberg EJ. A randomized intervention study to evaluate whether electronic messaging can increase human papillomavirus vaccine completion and knowledge among college students. *Journal of American College Health* 2016;**64**(4):269-78.

Robbins 2013 {published data only}

Robbins GK, Testa MA, Su M, Safren SA, Morse G, Lammert S, et al. Site nurse-initiated adherence and symptom support telephone calls for HIV-positive individuals starting antiretroviral therapy, ACTG 5031: substudy of ACTG 384. *HIV Clinical Trials* 2013;**14**(5):235-53.

Sridhar 2013 {published data only}

Sridhar A, Chen A, Glik D. Plan a birth control: randomized controlled trial of a mobile health application. *Contraception* 2013;**88**(3):463.

Sridhar 2014 {published data only}

Sridhar A, Chen AY. Mobile health application for long-acting reversible contraceptive information: a secondary analysis. *Obstetrics & Gynecology* 2014;**123**:111S.

Stern 2013 {published data only}

Stern L, Unger Z, Debevec E, Ginde S, Morfesis J, Patel A. Staying on track: a cluster randomized controlled trial of automated reminders for HPV vaccine series completion. *Contraception* 2013;**88**(3):438-9.

Stockwell 2012 {published data only}

Stockwell MS, Kharbanda EO, Martinez RA, Vargas CY, Vawdrey DK, Camargo S. Effect of a text messaging intervention on influenza vaccination in an urban, low-income pediatric and adolescent population: a randomized controlled trial. *Journal of the American Medical Association* 2012;**307**(16):1702-8.

Szilagyi 2013 {published data only}

Szilagyi PG, Albertin C, Humiston SG, Rand CM, Schaffer S, Brill H, et al. A randomized trial of the effect of centralized reminder/recall on immunizations and preventive care visits for adolescents. *Academic Pediatrics* 2013;**13**(3):204-13.

Takeuchi 2016 {published data only}

Takeuchi S, Horiuchi S. Randomised controlled trial using smartphone website vs leaflet to support antenatal perineal massage practice for pregnant women. *Women and Birth* 2016;**29**(5):430-5.

Tarrant 2014 {published data only}

Tarrant AM, Fong DY, Heys M, Lee IL, Sham A, Hui Choi WH. Professional breastfeeding support to increase the exclusivity and duration of breastfeeding: a randomised controlled trial. *Hong Kong Medical Journal* 2014;**20**:34.

Trent 2013 {published data only}

Trent M, Tomaszewski K. Family planning appointment attendance among urban youth: results from the Depotext trial. *Journal of Adolescent Health* 2013;**52**(2):S88.

Trent 2015 {published data only}

Trent M, Thompson C, Tomaszewski K. Text messaging support for urban adolescents and young adults using injectable contraception: outcomes of the DepoText pilot trial. *Journal of Adolescent Health* 2015;**57**(1):100-6.

Van Ryswyk 2015 {published data only}

Van Ryswyk EM, Middleton PF, Hague WM, Crowther CA. Postpartum SMS reminders to women who have experienced gestational diabetes to test for type 2 diabetes: the DIAMIND randomized trial. *Diabetic Medicine* 2015;**32**(10):1368-76.

Wright 2012 {published data only}

Wright A, Poon EG, Wald J, Feblowitz J, Pang JE, Schnipper JL, et al. Randomized controlled trial of health maintenance reminders provided directly to patients through an electronic PHR. *Journal of General Internal Medicine* 2012;**27**(1):85-92.

Young 2013 {published data only}

Young SD, Cumberland WG, Lee SJ, Jaganath D, Szekeres G, Coates T. Social networking technologies as an emerging tool for HIV prevention: a cluster randomized trial. *Annals of Internal Medicine* 2013;**159**(5):318-24.

Young 2014 {published data only}

Young SD, Holloway I, Jaganath D, Rice E, Westmoreland D, Coates T. Project HOPE: online social network changes in an HIV prevention randomized controlled trial for African American and Latino men who have sex with men. *American Journal of Public Health* 2014;**104**(9):1707-12.

References to studies awaiting assessment
Abroms 2017a {published data only}

Abroms LC, Johnson PR, Leavitt LE, Cleary SD, Bushar J, Brandon TH, et al. A randomized trial of text messaging for smoking cessation in pregnant women. *American Journal of Preventive Medicine* 2017;**53**(6):781-90. [DOI: doi.org/10.1016/j.amepre.2017.08.002]

Abroms 2017b {published data only}

Abroms LC, Chiang S, Macherelli L, Leavitt L, Montgomery M. Assessing the National Cancer Institute's smokefreeMOM text-

messaging program for pregnant smokers: pilot randomized trial. *Journal of Medical Internet Research* 2017;**19**(10):e333.

Altazan 2019 {published data only}

Altazan AD, Redman LM, Burton JH, Beyl RA, Cain LE, Sutton EF, et al. Mood and quality of life changes in pregnancy and postpartum and the effect of a behavioral intervention targeting excess gestational weight gain in women with overweight and obesity: a parallel-arm randomized controlled pilot trial. *BMC Pregnancy and Childbirth* 2019;**19**(1):50.

Bangal 2018 {published data only}

Bangal V, Somasundaram KV, Thitame S. Influence of mobile communication on utilization and outcome of maternal health services in rural area. *Indian Journal of Public Health Research & Development* 2018;**9**(5):504-8.

Domek 2019 {published data only}

Domek GJ, Contreras-Roldan IL, Bull S, O'Leary ST, Bolaños Ventura GA, Bronsert M, et al. Text message reminders to improve infant immunization in Guatemala: a randomized clinical trial. *Vaccine* 2019;**37**(42):6192-200.

Forinash 2018 {published data only}

Forinash AB, Yancey A, Chamness D, Koerner J, Inteso C, Miller C, et al. Smoking cessation following text message intervention in pregnant women. *Annals of Pharmacotherapy* 2018;**52**(11):1109-16.

Gibson 2019 {published data only}

Gibson DG, Kagucia EW, Were J, Obor D, Hayford K, Ochieng B. Text message reminders and unconditional monetary incentives to improve measles vaccination in western Kenya: study protocol for the mobile and scalable innovations for measles immunization randomized controlled trial. *JMIR Research Protocols* 2019;**8**(7):e13221.

Harari 2018 {published data only}

* Harari N, Rosenthal MS, Bozzi V, Goeschel L, Jayewickreme T, Onyebekwe C, et al. Feasibility and acceptability of a text message intervention used as an adjunct tool by WIC breastfeeding peer counsellors: the LATCH pilot. *Maternal & Child Nutrition* 2018;**14**(1):e12488.

Martinez-Brockman JL, Harari N, Segura-Pérez S, Goeschel L, Bozzi V, Pérez-Escamilla R. Impact of the lactation advice through texting can help (LATCH) trial on time to first contact and exclusive breastfeeding among WIC participants. *Journal of Nutrition Education and Behavior* 2018;**50**(1):33-42.

Martinez-Brockman JL, Harari N, Segura-Perez S, Goeschel L, Perez-Escamilla R. Impact of the lactation advice through texting can help (LATCH) randomized controlled trial. *FASEB Journal* 2017;**31**(Suppl):457.7.

Harrington 2019 {published data only}

* Harrington EK, Drake AL, Matemo D, Ronen K, Osoti AO, John-Stewart G, et al. An mhealth SMS intervention on postpartum contraceptive use among women and couples in Kenya: a randomized controlled trial. *American Journal of Public Health* 2019;**109**(6):934-41.

Harrington EK. Evaluation of an mHealth SMS Dialogue Strategy to Meet Women's and Couples' Postpartum Contraceptive Needs in Kenya (Mobile WACH XY): a Randomized Controlled Trial [Masters' thesis]. Vol. **10688609**. Washington (DC): University of Washington, 2017.

Hmone 2017 {published data only}

* Hmone MP, Li M, Agho K, Dibley M. Impact of SMS text messages to improve exclusive breastfeeding and reduce other adverse infant feeding practices in Yangon, Myanmar: a randomized controlled trial. *Annals of Nutrition and Metabolism* 2017;**71**:610-11.

Hmone MP, Li M, Agho K, Dibley MJ. Impact of SMS text messages to improve exclusive breastfeeding: a randomized controlled trial in Myanmar. *FASEB Journal* 2017;**31**(1):313.3.

Hmone MP, Li M, Alam A, Dibley MJ. Mobile phone short messages to improve exclusive breastfeeding and reduce adverse infant feeding practices: protocol for a randomized controlled trial in Yangon, Myanmar. *JMIR Research Protocols* 2017;**6**(6):e126.

Jasemzadeh 2018 {published data only} **IRCT2016102810804N8**

Jasemzadeh M, Khafaie MA, Jaafarzadeh N, Araban M. Effectiveness of a theory-based mobile phone text message intervention for improving protective behaviors of pregnant women against air pollution: a randomized controlled trial. *Environmental Science and Pollution Research* 2018;**25**(7):6648-55.

Oliveira-Ciabati 2017 {published data only}

Oliveira-Ciabati L, Vieira CS, Franzon AC, Alves D, Zaratini FS, Braga GC, et al. PRENACEL- a mHealth messaging system to complement antenatal care: a cluster randomized trial. *Reproductive Health* 2017;**14**(1):146.

Ortiz 2018 {published data only}

Ortiz J. Post-abortion follow-up through SMS: texting alternatives to unnecessary follow-up visits. *International Journal of Gynaecology and Obstetrics* 2018;**143**(Suppl 3):54.

Palacios 2018 {published data only}

Banna J, Campos M, Gibby C, Graulau RE, Melendez M, Reyes A, et al. Multi-site trial using short mobile messages (SMS) to improve infant weight in low-income minorities: development, implementation, lessons learned and future applications. *Contemporary Clinical Trials* 2017;**62**:56-60.

Palacios C, Campos M, Gibby C, Banna J. Multi-site trial using short mobile messages (SMS) to improve infant feeding practices among participants in the WIC program. *FASEB Journal* 2017;**31**(Suppl):959.8-959.8.

* Palacios C, Campos M, Gibby C, Meléndez M, Lee JE, Banna J. Effect of a multi-site trial using short message service (SMS) on infant feeding practices and weight gain in low-income minorities. *Journal of the American College of Nutrition* 2018;**37**(7):605-13.

Patel 2018 {published data only}

Patel A, Kuhite P, Puranik A, Khan SS, Borkar J, Dhande L. Effectiveness of weekly cell phone counselling calls and daily text messages to improve breastfeeding indicators. *BMC Pediatrics* 2018;**18**(1):337.

Seth 2018 {published data only}

Seth R, Akinboyo I, Chhabra A, Qaiyum Y, Shet A, Gupte N, et al. Mobile phone incentives for childhood immunizations in rural India. *Pediatrics* 2018;**141**(4):e20173455.

Shinde 2018 {published data only}

Shinde K, Rani U, Kumar PN. Assessing the effectiveness of immunization reminder system among nursing mothers of South India. *Research Journal of Pharmacy and Technology* 2018;**11**(5):1761-7.

Shorey 2017 {published data only} **ISRCTN99092313**

* Shorey S, Lau Y, Dennis CL, Chan YS, Tam WW, Chan YH. A randomized-controlled trial to examine the effectiveness of the 'Home-but not Alone' mobile-health application educational programme on parental outcomes. *Journal of Advanced Nursing* 2017;**73**(9):2103-17.

Shorey S, Ng YP, Danbjorg DB, Dennis CL, Morelius E. Effectiveness of the 'Home-but not Alone' mobile health application educational programme on parental outcomes: a randomized controlled trial, study protocol. *Journal of Advanced Nursing* 2017;**73**:253-64.

Unger 2018 {published data only}

Unger JA, Ronen K, Perrier T, DeRenzi B, Slyker J, Drake AL, et al. Short message service communication improves exclusive breastfeeding and early postpartum contraception in a low-to middle-income country setting: a randomised trial. *BJOG* 2018;**125**(12):1620-9.

Wang 2018 {published data only} **ISRCTN44149146**

Wang X, Luo R, Liu C, Zhang L, Yue A, Medina A, et al. Using daily text messages to improve adherence to infant micronutrient powder (MNP) packets in rural western China: a cluster-randomized controlled trial. *PLoS One* 2018;**13**(1):e0191549.

Xie 2018 {published data only}

Xie RH, Tan H, Taljaard M, Liao Y, Krewski D, Du Q, et al. The impact of a maternal education program through text messaging in rural China: cluster randomized controlled trial. *JMIR mHealth and uHealth* 2018;**6**(12):e11213.

References to ongoing studies

ACTRN12618000636257 {published data only}

ACTRN12618000636257. Investigating the impact of short automated text message reminder system in improving influenza vaccine uptake in children with chronic lung conditions: a randomised controlled trial. anzctr.org.au/Trial/Registration/TrialReview.aspx?id=374892 (first received 12 April 2018).

Adam 2019 {published data only}

Adam M, Tomlinson M, Le Roux I, LeFevre AE, McMahon SA, Johnston J, et al. The Philani MOVIE study: a cluster-randomized controlled trial of a mobile video entertainment-education intervention to promote exclusive breastfeeding in South Africa. *BMC Health Services Research* 2019;**19**:211.

CTRI/2018/04/013510 {published data only}

CTRI/2018/04/013510. Effectiveness of mHealth for improving fetal outcome: a community based intervention trial. ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=20357 (first received 27 April 2018).

Cyan 2016 {published data only}

Cyan R, Hussain H, Shehki I, Ishaq Z, Khan T, Rothenberg R. Affordable technology for saving maternal and infant lives: moving on with solutions. *Annals of Global Health* 2016;**82**:382.

Gelano 2018 {published data only}

Gelano TF, Assefa N, Bacha YD, Mahamed AA, Roba KT, Hambisa MT. Effect of mobile-health on maternal health care service utilization in Eastern Ethiopia: study protocol for a randomized controlled trial. *Trials* 2018;**19**(1):102.

Gul 2019 {published data only}

Gul X, Hameed W, Hussain S, Sheikh I, Siddiqui JU. A study protocol for an mhealth, multi-centre randomized control trial to promote use of postpartum contraception amongst rural women in Punjab, Pakistan. *BMC Pregnancy Childbirth* 2019;**19**(1):283.

IRCT20180520039728N1 {published data only}

IRCT20180520039728N1

IRCT20180520039728N1. Effectiveness of the distance education program on the mothers' empowerment in breastfeeding. en.irct.ir/trial/31317 (first received 3 July 2018).

ISRCTN13224744 {published data only}

ISRCTN13224744

ISRCTN13224744. The effectiveness of theory based intervention using social media to reduce urinary incontinence among postpartum women in Hebron city hospitals. isrctn.com/ISRCTN13224744 (first received 26 July 2018).

ISRCTN15017499 {published data only}

ISRCTN15017499

ISRCTN15017499. A trial exploring the feasibility of using telephone support (SMS and call) as a means of supporting young mothers (12–19 years) in Western Kenya soon after giving birth. isrctn.com/ISRCTN15017499 (first received 21 June 2017).

Kazi 2019 {published data only}

Kazi AM, Ahsan N, Khan A, Jamal S, Kalimuddin H, Ghulamhussain N, et al. Personalized text messages and automated calls for improving vaccine coverage among children in Pakistan: protocol for a community-based cluster randomized clinical trial. *JMIR Research Protocols* 2019;**8**(5):e12851.

Lefevre 2019 {published data only}

Lefevre A, Agarwal S, Chamberlain S, Scott K, Godfrey A, Chandra R, et al. Are stage-based health information messages effective and good value for money in improving maternal

newborn and child health outcomes in India? Protocol for an individually randomized controlled trial. *Trials* 2019;**20**(1):272.

Maycock 2015 {published data only}

Maycock BR, Scott JA, Hauck YL, Burns SK, Robinson S, Giglia R, et al. A study to prolong breastfeeding duration: design and rationale of the Parent Infant Feeding Initiative (PIFI) randomised controlled trial. *BMC Pregnancy and Childbirth* 2015;**15**(1):159.

NCT02371213 {published data only}

NCT02371213. Social networking on mobile phone to improve maternal and neonatal outcomes (HISONET). clinicaltrials.gov/show/NCT02371213 (first received 25 February 2015).

NCT03023033 {published data only}

NCT03023033. Supporting attendance for facility delivery and infant health (SAFI). clinicaltrials.gov/show/NCT03023033 (first received 18 January 2017).

NCT03135288 {published data only}

NCT03135288. Cell-phone assisted postpartum counseling on the use of long-acting reversible contraceptives. clinicaltrials.gov/ct2/show/NCT03135288 (first received 1 May 2017).

NCT03332108 {published data only}

NCT03332108. Novel approach to improving lactation support using mobile health technology. clinicaltrials.gov/ct2/show/NCT03332108 (first received 6 November 2017).

NCT03355989 {published data only}

NCT03355989. Evaluation of conditional cash transfers (CCTs) for immunization. clinicaltrials.gov/ct2/show/NCT03355989 (first received 29 November 2017).

NCT03379467 {published data only}

NCT03379467. Use of SMS and interactive reminders to improve timely immunization coverage: a randomized controlled trial. clinicaltrials.gov/ct2/show/NCT03379467 (first received 20 November 2017).

NCT03399058 {published data only}

NCT03399058. Innovative nutrition and mhealth evidence building project. clinicaltrials.gov/ct2/show/NCT03399058 (first received 16 January 2018).

NCT03402646 {published data only}

NCT03402646. Mobile phone reminders (and photovoice) for routine immunization in Nigeria – the MOPING study. clinicaltrials.gov/ct2/show/NCT03402646 (first received 18 January 2018).

NCT03516682 {published data only}

NCT03516682. The REDIVAC study-reducing delay in vaccination of children study. clinicaltrials.gov/ct2/show/NCT03516682 (first received 4 May 2018).

NCT03705455 {published data only}

NCT03705455. Immunization schedule alert platform: determining ISAP SMS efficacy in improving childhood

immunization timeliness and completeness in Nigeria. clinicaltrials.gov/ct2/show/NCT03705455 (first received 15 October 2018).

NCT03797950 {published data only}

NCT03797950. Mobile nudges to increase early vaccination coverage in rural areas: a pilot Investigation in Ghana's northern region. clinicaltrials.gov/ct2/show/NCT03797950 (first received 9 January 2019).

NCT03890978 {published data only}

NCT03890978. Improving exclusive breastfeeding via mobile phone text messages: a randomized controlled trial in southern Jordan. clinicaltrials.gov/ct2/show/NCT03890978 (first received 26 March 2019).

Odeny 2018 {published data only}

Odeny TA, Onono M, Owuor K, Helova A, Wanga I, Bukusi EA, et al. Maximizing adherence and retention for women living with HIV and their infants in Kenya (MOTIVATE! study): study protocol for a randomized controlled trial. *Trials* 2018;**19**(1):77.

PACTR201703002093382 {published data only}

PACTR201703002093382. Effect of enhanced reminders on postnatal clinic attendance in Addis Ababa: a cluster randomized control trial. pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=2093 (first received 11 March 2017).

PACTR201711002737120 {published data only}

PACTR201711002737120. Zinc adherence: a follow-up study of under-fives with acute watery diarrhoea using mobile phones: a randomised controlled trial. pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=2737 (first received 31 October 2017).

PACTR201801002231314 {published data only}

PACTR201801002231314. Utilization of short message service to enhance uptake of focused ante-natal care among women in Tharaka Nithi County, Kenya. pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=2231 (first received 23 April 2017).

PACTR201806003369302 {published data only}

PACTR201806003369302. Efficacy of text messages, voice calls and community visits on developmental milestones of children from teenage pregnancies: a randomized intervention trial. pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=3369 (first received 4 May 2018).

Rossing 2016 {published data only}

Rossing E, Ravn H, Batista CS, Rodrigues A. MHealth to improve measles immunization in Guinea-Bissau: study protocol for a randomized controlled trial. *JMIR Research Protocols* 2016;**5**(3):e158.

Salam 2018 {published data only}

Salam SS, Ali NB, Rahman AE, Tahsina T, Islam MI, Iqbal A, et al. Study protocol of a 4-parallel arm, superiority, community based cluster randomized controlled trial comparing paper and e-platform based interventions to improve accuracy of recall of last menstrual period (LMP) dates in rural Bangladesh. *BMC Public Health* 2018;**18**(1):1359.

Tobe 2018 {published data only}

Tobe RG, Haque SE, Ikegami K, Mori R. Mobile-health tool to improve maternal and neonatal health care in Bangladesh: a cluster randomized controlled trial. *BMC Pregnancy and Childbirth* 2018;**18**(1):102.

Additional references

Agarwal 2016

Agarwal S, LeFevre AE, Lee J, L'Engle K, Mehl G, Sinha C, et al. Guidelines for reporting of health interventions using mobile phones: mobile health (mHealth) evidence reporting and assessment (mERA) checklist. *BMJ* 2016;**352**:i1174. [DOI: doi.org/10.1136/bmj.i1174]

Agarwal 2018

Agarwal S, Vasudevan L, Tamrat T, Glenton C, Lewin S, Bergman H, et al. Digital tracking, provider decision support systems, and targeted client communication via mobile devices to improve primary health care. *Cochrane Database of Systematic Reviews* 2018, Issue 1. Art. No: CD012925. [DOI: [10.1002/14651858.CD012925](https://doi.org/10.1002/14651858.CD012925)]

Alkema 2016

Alkema L, Chou D, Hogan D, Zhang S, Moller AB, Gemmill A, et al. Global, regional, and national levels and trends in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet* 2016;**387**(10017):462-74.

Amankwaa 2018

Amankwaa I, Boateng D, Quansah DY, Akuoko CP, Evans C. Effectiveness of short message services and voice call interventions for antiretroviral therapy adherence and other outcomes: a systematic review and meta-analysis. *PLoS One* 2018;**13**(9):e0204091. [DOI: [10.1371/journal.pone.0204091](https://doi.org/10.1371/journal.pone.0204091)]

Ames 2019

Ames HM, Glenton C, Lewin S, Tamrat T, Akama E, Leon N. Clients' perceptions and experiences of targeted digital communication accessible via mobile devices for reproductive, maternal, newborn, child, and adolescent health: a qualitative evidence synthesis. *Cochrane Database of Systematic Reviews* 2019, Issue 10. Art. No: CD013447. [DOI: [10.1002/14651858.CD013447](https://doi.org/10.1002/14651858.CD013447)]

Black 2016

Black RE, Levin C, Walker N, Chou D, Liu L, Temmerman M, et al. Reproductive, maternal, newborn, and child health: key messages from Disease Control Priorities 3rd Edition. *Lancet* 2016;**388**(10061):2811-24.

Chamberlain 2017

Chamberlain C, O'Mara-Eves A, Porter J, Coleman T, Perlen SM, Thomas J, et al. Psychosocial interventions for supporting women to stop smoking in pregnancy. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No: CD001055. [DOI: [10.1002/14651858.CD001055.pub5](https://doi.org/10.1002/14651858.CD001055.pub5)]

de Jongh 2012

de Jongh T, Guroi-Urganci I, Vodopivec-Jamsek V, Car J, Atun R. Mobile phone messaging for facilitating self-management of long-term illnesses. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No: CD007459. [DOI: [10.1002/14651858.CD007459.pub2](https://doi.org/10.1002/14651858.CD007459.pub2)]

Every Women Every Child 2015

Every Woman Every Child. The global strategy for women's and children's and adolescents' health (2016-2030): survive, thrive, transform. www.who.int/life-course/partners/global-strategy/globalstrategyreport2016-2030-lowres.pdf (accessed prior to 3 May 2020).

Free 2011

Free C, Knight R, Robertson S, Whittaker R, Edwards P, Zhou W, et al. Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial. *Lancet* 2011;**378**(9785):49-55. [DOI: [doi.org/10.1016/S0140-6736\(11\)60701-0](https://doi.org/10.1016/S0140-6736(11)60701-0)]

Free 2013a

Free C, Phillips G, Galli L, Watson L, Felix L, Edwards P, et al. The effectiveness of mobile-health technology-based health behaviour change or disease management interventions for health care consumers: a systematic review. *PLoS Medicine* 2013;**10**(1):e1001362.

Free 2013b

Free C, Phillips G, Watson L, Galli L, Felix L, Edwards P, et al. The effectiveness of mobile-health technologies to improve health care service delivery processes: a systematic review and meta-analysis. *PLoS Medicine* 2013;**10**(1):e1001363. [DOI: doi.org/10.1371/journal.pmed.1001363]

Gonçalves-Bradley 2018

Gonçalves-Bradley DC, Buckley BS, Fønhus MS, Glenton C, Henschke N, Lewin S, et al. Mobile-based technologies to support client to healthcare provider communication and management of care. *Cochrane Database of Systematic Reviews* 2018, Issue 1. Art. No: CD012928. [DOI: [10.1002/14651858.CD012928](https://doi.org/10.1002/14651858.CD012928)]

Griffiths 2018

Griffiths SE, Parsons J, Naughton F, Fulton EA, Tombor I, Brown KE. Are digital interventions for smoking cessation in pregnancy effective? A systematic review and meta-analysis. *Health Psychology Review* 2018;**12**(4):333-56. [DOI: doi.org/10.1080/17437199.2018.1488602]

Guerriero 2013

Guerriero C, Cairns J, Roberts I, Rodgers A, Whittaker R, Free C. The cost-effectiveness of smoking cessation support delivered by mobile phone text messaging: Txt2stop. *European Journal of Health Economics* 2013;**14**(5):789-97. [DOI: doi.org/10.1007/s10198-012-0424-5]

Guroi-Urganci 2013

Guroi-Urganci I, de Jongh T, Vodopivec-Jamsek V, Atun R, Car J. Mobile phone messaging reminders for attendance at healthcare appointments. *Cochrane Database of*

Systematic Reviews 2013, Issue 12. Art. No: CD007458. [DOI: [10.1002/14651858.CD007458.pub3](https://doi.org/10.1002/14651858.CD007458.pub3)]

Hawkins 2008

Hawkins RP, Kreuter M, Resnicow K, Fishbein M, Dijkstra A. Understanding tailoring in communicating about health. *Health Education Research* 2008;**23**(3):454-66.

Hay 2016

Hay SI, Abajobit AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2016;**390**:1260-344.

Higgins 2011

Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hoffmann 2014

Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;**348**:g1687. [DOI: doi.org/10.1136/bmj.g1687]

Horvath 2012

Horvath T, Azman H, Kennedy GE, Rutherford GW. Mobile phone text messaging for promoting adherence to antiretroviral therapy in patients with HIV infection. *Cochrane Database of Systematic Reviews* 2012, Issue 3. Art. No: CD009756. [DOI: [10.1002/14651858.CD009756](https://doi.org/10.1002/14651858.CD009756)]

ICT 2016

ICT. Key ICT indicators for developed and developing countries and the world. idp.nz/Global-Rankings/ITU-Key-ICT-Indicators/6mef-ytg6 (accessed 1 December 2016).

Kaufman 2017

Kaufman J, Ames H, Bosch-Capblanch X, Cartier Y, Cliff J, Glenton C, et al. The comprehensive 'Communicate to Vaccinate' taxonomy of communication interventions for childhood vaccination in routine and campaign contexts. *BMC Public Health* 2017;**17**(1):423.

Kaufman 2018

Kaufman J, Ryan R, Walsh L, Horey D, Leask J, Robinson P, et al. Face-to-face interventions for informing or educating parents about early childhood vaccination. *Cochrane Database of Systematic Reviews* 2018, Issue 5. Art. No: CD010038. [DOI: [10.1002/14651858.CD010038.pub3](https://doi.org/10.1002/14651858.CD010038.pub3)]

Lee 2016

Lee SH, Nurmatov UB, Nwaru BI, Mukherjee M, Grant L, Pagliari C. Effectiveness of mHealth interventions for maternal, newborn and child health in low- and middle-income countries: systematic review and meta-analysis. *Journal of Global Health* 2016;**6**(1):010401.

Liberati 2009

Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Medicine* 2009;**6**:e1000100.

Lorencatto 2012

Lorencatto F, West R, Michie S. Specifying evidence-based behavior change techniques to aid smoking cessation in pregnancy. *Nicotine and Tobacco Research* 2012;**14**(9):1019-26. [DOI: doi.org/10.1093/ntr/ntr324]

McFadden 2017

McFadden A, Gavine A, Renfrew MJ, Wade A, Buchanan P, Taylor JL, et al. Support for healthy breastfeeding mothers with healthy term babies. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No: CD001141. [DOI: [10.1002/14651858.CD001141.pub5](https://doi.org/10.1002/14651858.CD001141.pub5)]

Naghavi 2017

Naghavi M, Abajobir AA, Abbafati C, Abbas KM, Abd-Allah F, Abera SF, et al. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;**390**:1151-210.

Pagel 2011

Pagel C, Prost A, Lewycka S, Das S, Colbourn T, Mahapatra R, et al. Intracluster correlation coefficients and coefficients of variation for perinatal outcomes from five cluster-randomised controlled trials in low and middle-income countries: results and methodological implications. *Trials* 2011;**12**(1):151. [DOI: [10.1186/1745-6215-12-151](https://doi.org/10.1186/1745-6215-12-151)]

Palmer 2018b

Palmer M, Sutherland J, Barnard S, Wynne A, Rezel E, Doel A, et al. The effectiveness of smoking cessation, physical activity/diet and alcohol reduction interventions delivered by mobile phones for the prevention of non-communicable diseases: a systematic review of randomised controlled trials. *PLoS One* 2018;**13**(1):e0189801. [DOI: doi.org/10.1371/journal.pone.0189801]

Palmer in preparation

Palmer MJ, Henschke N, Villanueva G, Maayan N, Bergman H, Glenton C, et al. Targeted client communication via mobile devices for improving reproductive and sexual health. Cochrane Database of Systematic Reviews In preparation.

PMNCH 2011

The Partnership for Maternal, Newborn and Child Health. A global review of the key interventions related to reproductive, maternal, newborn and child health (RMNCH). Geneva: PMNCH 2011.

Quintana 2018

Quintana Y, Martorell EA, Fahy D, Safran C. A systematic review on promoting adherence to antiretroviral therapy in HIV-infected patients using mobile phone technology.

Applied Clinical Informatics 2018;**9**(2):450-66. [DOI: [10.1055/s-0038-1660516](https://doi.org/10.1055/s-0038-1660516)]

Review Manager 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Ryan 2013

Ryan R, Hill S, Prictor M, McKenzie J, Cochrane Consumers and Communication Group. Study quality guide, May 2013. cccg.cochrane.org/authorresources (accessed 8 January 2018).

Ryan 2016

Ryan R, Horey D, Oliver S, McKenzie J, Prictor M, Santesso N, et al. Cochrane Consumers and Communication Group standard protocol text and additional guidance for review authors. CCCG cccg.cochrane.org/author-resources (accessed October 2017).

Say 2014

Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Global Health* 2014;**2**(6):e323-33.

Schünemann 2011

Schünemann HJ, Oxman AD, Higgins JP, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results and 'Summary of findings' tables. In: Higgins JP, Green S, editor(s), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Shah 2019

Shah R, Watson J, Free C. A systematic review and meta-analysis in the effectiveness of mobile phone interventions used to improve adherence to antiretroviral therapy in HIV infection. *BMC Public Health* 2019;**19**(1):915. [DOI: [10.1186/s12889-019-6899-6](https://doi.org/10.1186/s12889-019-6899-6)]

Singh 2014

Singh N. Community-based and innovative technological approaches to improve child nutrition in India [PhD thesis]. London: LSHTM University of London, 2014.

Smith 2015

Smith C, Ngo TD, Gold J, Edwards P, Vannak U, Sokhey L, et al. Effect of a mobile phone-based intervention on post-abortion contraception: a randomized controlled trial in Cambodia. *Bulletin of the World Health Organization* 2015;**93**(12):842-50a.

Stade 2009

Stade BC, Bailey C, Dzenoletas D, Sgro M, Dowswell T, Bennett D. Psychological and/or educational interventions for reducing alcohol consumption in pregnant women and women planning pregnancy. *Cochrane Database of Systematic Reviews* 2009, Issue 2. Art. No: CD004228. [DOI: [10.1002/14651858.CD004228.pub2](https://doi.org/10.1002/14651858.CD004228.pub2)]

UN 2015

United Nations. Transforming our world: the 2030 agenda for sustainable development. Geneva: United Nations (A/RES/70/1) 2015.

Vodopivec-Jamsek 2012

Vodopivec-Jamsek V, de Jongh T, Gurol-Urganci I, Atun R, Car J. Mobile phone messaging for preventive health care. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No: CD007457. [DOI: [10.1002/14651858.CD007457.pub2](https://doi.org/10.1002/14651858.CD007457.pub2)]

Whittaker 2016

Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y. Mobile phone-based interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2016, Issue 4. Art. No: CD006611. [DOI: [10.1002/14651858.CD006611.pub4](https://doi.org/10.1002/14651858.CD006611.pub4)]

WHO 2010

World Health Organization. Packages of interventions for family planning, safe abortion care, maternal, newborn and child health. apps.who.int/iris/handle/10665/70428 (accessed prior to 3 May 2020).

WHO 2013

World Health Organization, London School of Hygiene and Tropical Medicine, South Africa Medical Research Council. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. apps.who.int/iris/handle/10665/85239 (accessed prior to 3 May 2020).

WHO 2017

World Health Organization, United Nations Children's Fund. Reaching the every newborn national 2020 milestones: country progress, plans and moving forward. apps.who.int/iris/handle/10665/255719 (accessed prior to 3 May 2020).

WHO 2018

World Health Organization. Classification of digital health interventions v1.0A shared language to describe the uses of digital technology for health. apps.who.int/iris/handle/10665/260480 (accessed prior to 3 May 2020).

WHO 2019

World Health Organization. WHO guideline: recommendations on digital interventions for health system strengthening. apps.who.int/iris/handle/10665/311980 (accessed prior to 3 May 2020).

WHO/ITU 2015

World Health Organization and International Telecommunication Union. A handbook on how to implement mTobaccoCessation. apps.who.int/iris/handle/10665/251719 (accessed prior to 3 May 2020).

World Bank 2017

World Bank. World Development Indicators 2017. IGO.openknowledge.worldbank.org/handle/10986/26447 (accessed 30 July 2017).

References to other published versions of this review
Palmer 2018a

Palmer MJ, Free C, Tamrat T, Henschke N, Glenton C, Lewin L, Fønhus M, Mehl G, Villanueva G, Maayan N, Bergman H. Targeted client communication via mobile devices for maternal, neonatal, and child health: systematic review protocol. CRD42018087234. PROSPERO 2018.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Ahlers-Schmidt 2012
Study characteristics

Methods	<p>Aim: to evaluate feasibility, stability of mobile service, and effect size.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: parents of newborns being discharged from a local hospital who intended to seek child health care at the university resident and faculty clinic. Parents enrolled in the study by texting "TRICKs" to the research phone number.</p> <p>Study duration: up to 1 year</p> <p>Study dates: January 2011 to February 2011</p>
Participants	<p>Inclusion criteria: English- and Spanish-speaking parents of newborns were eligible if they owned a mobile phone with text messaging capability.</p> <p>Sample size (n): 90 (intervention: 50; standard care: 40)</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Ahlers-Schmidt 2012 (Continued)

Age (mean): intervention: 26 (SD 5) years; control: 26 (SD 6) years

Sex (female): intervention: 84%; control: 83%

County: USA (high-income)

Setting: local hospital, university resident and faculty clinic in a Midwestern metropolitan area

Interventions	<p>Intervention: reminder text messages 7 days before their child's immunisations which were due at 2, 4, and 6 months of age.</p> <p>Content: reminder and request to respond to query whether their child was immunised.</p> <p>Frequency and intensity: 7 days before immunisations at 2, 4, and 6 months, assumed to be single text messages.</p> <p>Control: standard care/no intervention: parents in the control arm received standard notification (an appointment card) of immunisations due only.</p> <p>Co-interventions: none reported</p>	
Outcomes	<p>1. Attendance for vaccinations; 2. Timely attendance for vaccinations; 3. Satisfaction with the service (time point not reported)</p> <p>Outcome assessment time point: 2, 4, 6 months – we included the longest time point (6 months).</p>	
Funding/declarations of interest	<p>Funding: funded by a Wichita Center for Graduate Medical Education and Kansas Bioscience Authority Level III grant.</p> <p>Conflicts of interest: none</p>	
Notes	<p>Trial ID: NCT01396902</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation scheme.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants or personnel.
Blinding of objective outcome assessment (detection bias)	Low risk	Vaccination results were abstracted from immunisation registry records and were considered low risk of detection bias.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	2 participants excluded from analysis, reason provided.

Ahlers-Schmidt 2012 (Continued)

Selective reporting (reporting bias)	Unclear risk	All relevant outcomes appear to be reported but there was no protocol or on-line trial registration.
Other bias	Unclear risk	Intervention and control arm participants differ significantly on income, with 28% (11/40) of control parents having an income level \geq USD 25,000 annually compared with 12% (6/50) of intervention parents.

Bangure 2015
Study characteristics

Methods	<p>Aim: to assess whether immunisation coverage among those receiving short message reminders and routine immunisation health education differ from those receiving routine immunisation health education only.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: mother or carer soon after delivery or at third or seventh day visit after delivery</p> <p>Study duration: 14 weeks</p> <p>Study dates: 1 January 2013 to 31 August 2013</p>
Participants	<p>Inclusion criteria: mother or carer soon after delivery or at third or seventh day visit after delivery, must have mobile phone and be a resident of Kadoma City.</p> <p>Sample size (n): 304 (intervention: 152; control: 152)</p> <p>Age (median): intervention: 26 (Q1 21, Q3 30); control: 27 (Q1 23, Q3 32)</p> <p>Sex: not reported</p> <p>Country: Zimbabwe (low-income)</p> <p>Setting: city clinics and general hospital</p>
Interventions	<p>Intervention: 3 automatic text message reminders. First message sent 7 days before due date for immunisation, second message sent 3 days before, last message sent 1 day before immunisation appointment.</p> <p>Content: appointment reminder</p> <p>Frequency and intensity: 3 text message reminders before the 6-, 10-, and 14-week appointments</p> <p>Control: standard care/no intervention: informed about next scheduled visit</p> <p>Co-interventions: all participants received routine health education.</p>
Outcomes	<p>1. Attendance for vaccinations (6, 10, 14 weeks – we included the longest time point, 14 weeks); 3. opinion of SMS; 4. Cost of SMS service</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>2. Age of child when immunised (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 6, 10, 14 weeks</p>
Funding/declarations of interest	<p>Funding: Center for Disease Control and Prevention, Zimbabwe</p> <p>Conflicts of interest: none declared</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Bangure 2015 (Continued)

 Notes **Trial ID:** not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Study participants assigned by computer-generated random numbers.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants or personnel.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Not reported.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	0 lost.
Selective reporting (reporting bias)	Low risk	Only outcome reported in registration is immunisation coverage at 6, 10, and 14 weeks.
Other bias	Low risk	No apparent baseline differences.

Bigna 2015
Study characteristics

Methods	<p>Aim: to determine 1. if reminders sent by text message, phone call, or concomitant text and phone calls most increase the presence at medical appointments of HIV-infected or HIV-exposed children, and 2. which is the most efficient related to working time and financial cost.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: adult-child (carer-patient) pairs were recruited for HIV-infected or exposed children aged < 15 years attending HIV care.</p> <p>Study duration: not reported</p> <p>Study dates: January 2013 to May 2013</p>
Participants	<p>Inclusion criteria: people aged \geq 18 years accompanying an HIV-infected or HIV-exposed child aged < 15 years for HIV care</p> <p>Sample size (n): 242 (text message: 60; text message + call: 61; call: 60; control: 61)</p>

Bigna 2015 (Continued)

Age: children: age range up to 15 years, although we could not confirm that $\geq 70\%$ of included children were ≤ 5 years we decided to include this study since the weighted mean age of children was 2.8 years. Carers (mean): text message + call group: 36.5 (SD 12.4) years; text message group: 41.4 (SD 12.8) years; call group: 50.5 (SD 13.2) years; control group: 42.6 (SD 12.6)

Sex: carers – men: text message + call group: 20%; text message group: 13%; call group: 15%; control group: 13%; children – boys: text message + call group: 52%; text message group: 47%; call group: 47%; control group: 49%

Country: Cameroon (lower middle-income)

Setting: 3 hospitals serving urban (Essos), semi-urban (Kousséri), and rural (Goulfey) settings

Interventions	<p>Intervention: appointment reminders via text message; voice call; or text message + voice call</p> <p>Content: appointment reminder</p> <p>Frequency and intensity: 1 call, or 1 text message then 1 call 2–3 days before appointment</p> <p>Control: standard care/no intervention. No reminder (usual practice), all participants attended standard HIV care appointments</p> <p>Co-interventions: HIV care (to all participants)</p>
Outcomes	<p>1. Attendance at HIV medical appointments; 2. Direct costs; 3. Staff working time</p> <p>Outcome assessment time point: 2 days</p>
Funding/declarations of interest	<p>Funding: no external funding, done as part of an academic degree financed from the personal funds of the corresponding author Jean Joel R Bigna.</p> <p>Conflicts of interest: authors declare no competing interests</p>
Notes	<p>Trial ID: PACTR201304000528276</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated sequence. Quote: "Randomisation and allocation were done centrally with WinPepi version 11.25. Eligible participants (adult–child pairs) were randomly assigned in blocks of four and allocated (1:1:1:1) sequentially in the order of receipt of a randomisation code."
Allocation concealment (selection bias)	Low risk	Central allocation. Quote: "Randomisation and allocation were done centrally with WinPepi version 11.25. Eligible participants (adult–child pairs) were randomly assigned in blocks of four and allocated (1:1:1:1) sequentially in the order of receipt of a randomisation code."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Personnel were blinded. Not possible to blind participants. Quote: "The treating physician, the medical administrative assistant responsible for contacting participants in the intervention groups 2, 3, and 4 via mobile phone ... were all masked to group assignments."
Blinding of objective outcome assessment (detection bias)	Low risk	Outcome assessors and analysts were blinded.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Bigna 2015 (Continued)

Quote: "the nurse (outcome assessor) responsible for recording the patient's presence or absence at the appointment, and the data analysts were all masked to group assignments."

Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures would have been at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomised participants, whether or not they received the intervention, were included in the ITT analysis.
Selective reporting (reporting bias)	Low risk	All outcomes listed in the protocol were fully reported.
Other bias	Unclear risk	Some baseline characteristics (ages of children and carers, carers' education, and time to appointment) were unevenly distributed between groups.

Brown 2016
Study characteristics

Methods	<p>Aim: to assess the effect of reminder/recall system and primary healthcare immunisation providers' training intervention on routine immunisation.</p> <p>Study design: cluster RCT</p> <p>Recruitment: children aged 0–12 weeks at their first immunisation appointment</p> <p>Study duration: until child was 12 months old</p> <p>Study dates: August 2012 to September 2013</p>
Participants	<p>Inclusion criteria: children aged 0–12 weeks at their first immunisation visits, parents living in study communities</p> <p>Sample size: 595 (reminder: 153 participants, 1 cluster; usual care: 152 participants, 1 cluster; reminder + HCP training: 149 participants, 1 cluster – not eligible or included in review; HCP training: 141 participants, 1 cluster – not eligible or included in review)</p> <p>Age: children: 0–12 months; parents: not reported</p> <p>Sex (female): children: 51.4%</p> <p>Country: Nigeria (lower middle-income)</p> <p>Setting: community/health facility</p>
Interventions	<p>Intervention: mobile phone calls for the reminder/recall interventions sessions, reminder of immunisation appointment, recall for missed appointment</p> <p>Content: appointment reminder and recall for missed appointments</p> <p>Frequency and intensity: 2 telephone reminders to the contact person whose mobile phone number was provided by the mother – 2 days before and 1 day before the appointment.</p> <p>Control: standard care/no intervention: usual care</p> <p>Co-interventions: 2-day refresher training conducted for primary healthcare immunisation providers</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Brown 2016 (Continued)

Outcomes 1. "immunization completed" = receipt of all scheduled routine childhood immunisation = 1 dose of BCG vaccine, at ≥ 4 doses of OPV, 3 doses of DPT vaccine, 3 doses of hepatitis B vaccines, and 1 dose each of measles and yellow fever vaccine by age of 12 months

Outcome assessment time point: 12 months

Funding/declarations of interest **Funding:** not reported

Conflicts of interest: not reported

Notes 2 arms that included primary healthcare training were not extracted.

Trial ID: not reported

Cluster features: 4 local government authorities were randomised. Mean cluster size: 152.5 (for the 2 study arms included in the review). The study did not adjust for clustering effect; we used ICC 0.0487 (derived from $k = 0.089$ reported in [Gibson 2017](#)) to adjust for cluster effect in the analysis.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Using ballot system, the four randomly selected LGAs [local government association] were then allocated into three intervention groups and one control group." Comment: no further details reported.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Selective cluster recruitment	Unclear risk	Quote: "A total of 605 eligible children aged 0–12 weeks at their first immunization visits having their parents living in the study communities were consecutively recruited into four different study groups from August to November 2012." Comment: unclear whether participants were recruited before or after clusters were randomised.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants or personnel since interventions were active and different.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Outcome data collected using a standard instrument to extract data from comprehensive immunisation records, but unclear blinding.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Analyses were not ITT but there were few dropouts, due to either relocation or death.
Selective reporting (reporting bias)	Unclear risk	Reports the outcome described in methods, but no protocol or online trial record.
Other bias	High risk	Significant religious and social differences between clusters at baseline.

Domek 2016
Study characteristics

Methods	<p>Aim: 1. to test the feasibility and acceptance of SMS vaccination reminders sent to parents of children presenting for their infant primary immunisation series, and 2. to evaluate the potential of a large-scale programme to optimise immunisation delivery in a LMIC.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: parents of infants aged 8–14 weeks recruited when presenting for the first dose of the 3-dose infant primary immunisation series</p> <p>Study duration: 6 months</p> <p>Study dates: March 2013 to October 2013 (6 months after last enrolment April 2013)</p>
Participants	<p>Inclusion criteria: parents of infants aged 8–14 weeks, presenting for the first dose of the 3-dose infant primary immunisation series, owned a mobile phone with SMS text messaging capability, ≥ 1 parent had to be literate and able to use SMS technology.</p> <p>Sample size (n): 321 (intervention: 160; control: 161)</p> <p>Age (mean): parents: not reported; infants: intervention: 9.7 (SD 9.1) months; control: 9.4 (SD 9.0 months)</p> <p>Sex (male): intervention: 52.5%; control: 44.4%</p> <p>Country: Guatemala (lower middle-income)</p> <p>Setting: community, served by 2 public health clinics in Guatemala City</p>
Interventions	<p>Intervention: SMS reminders for infant immunisation</p> <p>Content: appointment reminders</p> <p>Frequency and intensity: SMS text messages at 6, 4, and 2 days before scheduled vaccination visits 2 and 3 (at 4 and 6 months of age)</p> <p>Control: standard care/no intervention</p> <p>Co-interventions: all participants received written reminders in the child's immunisation card for the next dose of vaccines at the time of each vaccination based on the usual standard of care at both clinics.</p>
Outcomes	<p>1. Attendance for vaccinations: completing all vaccinations (6 months); 2. Client satisfaction</p> <p><u>Outcomes reported but not included in review:</u></p> <p>1. Completing any vaccination (6 months) – instead, we included completion of all vaccinations (see above); 2. Attendance for vaccinations at second (4 months) and third visit (6 months) – instead, we included completion of all vaccinations (see above); 3. Completing pentavalent (3 doses)/pneumococcal (2 doses)/poliomyelitis (3 doses)/rotavirus (2 doses) vaccination; 4. Age at visits 2 and 3 (outcome not eligible for inclusion); 5. Proportion of parents sent SMS messages (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 4, 6 months</p>
Funding/declarations of interest	<p>Funding: Bill & Melinda Gates Foundation Grand Challenges Explorations grant and a small operational grant from the Pan American Health Organization</p> <p>Conflicts of interest: none declared</p>
Notes	<p>Trial ID: NCT01663636</p>

Domek 2016 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Participants were allocated to either an intervention or usual care group using a computer-generated randomization scheme."
Allocation concealment (selection bias)	Low risk	Quote: "Participants were allocated to either an intervention or usual care group ... with the investigators being blind to the allocation."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants. Personnel blind to allocation, but unclear whether this blinding was sustained.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Not reported whether research nurses that collected data were blinded. Immunisation records were able to be confirmed only for children who returned to either of the 2 study sites.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	17.4% were lost to follow-up but analyses were ITT.
Selective reporting (reporting bias)	Low risk	Outcomes listed in online trial record were reported.
Other bias	Unclear risk	Parents in the usual care group had significantly higher income with more fathers working; otherwise, there were no significant differences between the baseline demographics of intervention and usual care children and their parents.

Evans 2014
Study characteristics

Methods	<p>Aim: to evaluate initial outcomes of the Text4baby intervention at 4 weeks postbaseline.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: women recruited at the end of their initial ANC visit to the Madigan Obstetrics and Gynecology clinic.</p> <p>Study duration: 4 weeks (intervention planned to continue for whole pregnancy)</p> <p>Study dates: December 2011 (recruitment) to January 2013 (recruitment)</p>
Participants	<p>Inclusion criteria: women military healthcare beneficiaries (both active duty and family members), aged 18–45 years, presented for ANC prior to 14 weeks' gestation, working mobile phone, speaking, and reading fluent English</p> <p>Sample size (n): 943 (completers (number randomised per arm not reported): intervention: 229; control: 230)</p> <p>Age (mean): 26.53 years</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Evans 2014 (Continued)

Sex: 100% women

Country: USA (high-income)

Setting: community (military) served by a large tertiary-care army medical centre

Interventions	<p>Intervention: SMS messages</p> <p>Content: 135 distinct antenatal text messages aimed at promoting ways to live a satisfying lifestyle while reducing risk and promoting maternal and child health</p> <p>Frequency and intensity: 3 messages per week for 4 weeks in this report (intervention intended for whole pregnancy)</p> <p>Control: standard care/no intervention. No SMS messages</p> <p>Co-interventions: not reported</p>
Outcomes	<p>1. Self-reported consumption of alcohol since becoming aware of pregnancy; 2. Smoking in last 30 days (self-report)</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Attitudes and knowledge on pregnancy risks, self-report (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 4 weeks</p>
Funding/declarations of interest	<p>Funding: Telemedicine and Advanced Technology Research Center (TATRC), US Army Medical and Materiel Command</p> <p>Conflicts of interest: none declared</p>
Notes	<p>Trial ID: not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Using an algorithm that generated a randomized list of individual assignments to treatment or control condition ..."
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants. Clinicians who met with participants were blinded – the randomisation occurred outside the actual clinical visit and the trial data were not accessed by the clinicians during the study.
Blinding of subjective outcome assessment (detection bias)	High risk	Subjective outcomes self-reported by unblinded participants.
Incomplete outcome data (attrition bias) All outcomes	High risk	High level of attrition – 459/943 (48.7%) completed a 4-week follow-up survey.
Selective reporting (reporting bias)	Low risk	Expected outcomes reported, as per methods in the protocol/early results paper.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Evans 2014 (Continued)

Other bias	Unclear risk	Control group reported a larger, statistically significant percent smoking in the last 30 days and consuming ≥ 3 vegetables per day at baseline compared with the intervention group.
------------	--------------	--

Eze 2015
Study characteristics

Methods	<p>Aim: 1. To provide evidence validating the need for development and deployment of automated client reminder-recall systems for the Nigerian National Routine Immunisation Program and 2. to compare its projected cost with the cost of a health personnel-based defaulter tracking system.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: carers were selected using multistage sampling method from 8 health facilities in Egor local government area of Edo State.</p> <p>Study duration: each child recruited into the study at its first immunisation session (BCG) was followed up for 18 weeks while those who were recruited at their second session (DPT1) were followed up for 12 weeks to allow for 4 extra weeks after the recommended dates of receipt of DPT3.</p> <p>Study dates: June 2010 to June 2011</p>
Participants	<p>Inclusion criteria: bringing child for routine immunisation for the first or second schedules</p> <p>Sample size (n): 1001 (intervention: 500; control: 501)</p> <p>Age (mean): carers: 29.35 (SD 5.3) years</p> <p>Sex: 895 (89%) mothers, and 10 classed as 'other' but sex not stated</p> <p>Country: Nigeria (lower middle-income)</p> <p>Setting: facilities included 1 tertiary hospital, 2 primary health centres, and 5 privately owned health facilities all of which provided routine immunisation services.</p>
Interventions	<p>Intervention: reminder and recall text messages</p> <p>Authors also report that (quote), "All text messages were by internet-based web-to-SMS (Bulk SMS) service and were tagged the name of client's health facility for easy recognition." Messages sent the day before an immunisation appointment. There were 4 immunisation time points for BCG, DPT1, DPT2, and DPT3.</p> <p>Content: short reminder text message, and recall messages to those who missed appointments</p> <p>Frequency and intensity: reminder messages sent 1 day before the appointment, the number of reminder messages received depended the time point at which the participant was recruited. Recall messages were sent to defaulters. The total number of these messages was not reported.</p> <p>Control: standard care/no intervention, details not reported</p> <p>Co-interventions: none reported</p>
Outcomes	<p>1. Timeliness of receipt of DPT3; 2. Cost and cost-effectiveness (compared to estimated cost of home visits)</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Barriers to receiving text message reminders (outcome not eligible for inclusion)</p>

Eze 2015 (Continued)

Outcome assessment time point: postintervention, exact follow-up time not reported

Funding/declarations of interest **Funding:** not reported

Conflicts of interest: not reported

Notes **Trial ID:** not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Participants recruited per facility were randomized into 2 equal groups: intervention and Control groups using the RANDOM. EXE function of the Programme for Epidemiologists (PEPI) version 4.0."
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants or personnel.
Blinding of objective outcome assessment (detection bias)	Low risk	Quote: "Data collectors could not tell if a client was in the intervention or control group."
Blinding of subjective outcome assessment (detection bias)	High risk	Unblinded carers interviewed using follow-up questionnaires.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Available-case analysis with 9.6% (96 carers) attrition. Reasons for dropping out were provided and dropouts were balanced between groups.
Selective reporting (reporting bias)	Unclear risk	All expected outcomes and outcomes listed in the methods section were reported in the results, but no protocol or online trial record was identified to check against.
Other bias	High risk	Some participants switched groups after randomisation. Authors stated that, "Some respondents, who did not have mobile phones but were randomized into the Interventional group initially, were eventually matched for age and sex and swapped with persons who own mobile phones and were randomized into the Control group." Comment: in addition, carers provided answers to questions by recall; this could have led to recall bias in the information supplied.

Gibson 2017
Study characteristics

Methods **Aim:** to assess whether SMS reminders, with or without mobile money incentives, improve the proportion of children fully immunised by their first birthday.

Study design: cluster RCT

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Gibson 2017 (Continued)

Recruitment: the Health and Demographic Surveillance System village reporters identified eligible carers and their infants. Village reporters used mobile phones to send birth notification text messages. Birth notifications were relayed to field-based community interviewers who then screened carers of newborns for eligibility into the study.

Study duration: 12 months

Study dates: October 2013 (start date of recruitment) to October 2014 (end date of recruitment)

Participants	<p>Inclusion criteria: mothers/carers of infants aged 0–34 days during the study period, current resident of 1 of the participating study villages, willing to sign informed consent for the study</p> <p>Sample size: 2018 (intervention: 476 participants, 38 clusters; control: 489 participants, 38 clusters)*</p> <p>Age: mothers aged ≤ 25 years: 52% (n = 825); aged > 25 years 48% (n = 769); infants: mean 14 (SD 8) days</p> <p>Sex: 100% female (mothers)</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: community-based (villages)</p>
Interventions	<p>Intervention: SMS reminders composed of a core text and a motivational saying</p> <p>Frequency and intensity: 3 days and the day before scheduled immunisation visits at ages 6 weeks, 10 weeks, and 14 weeks for the 3 doses of pentavalent vaccine and age 9 months for measles vaccine</p> <p>Control: standard care/no intervention, did not receive SMS reminders or incentives</p> <p>Co-interventions: all carers received a single text message at enrolment welcoming them to the study</p>
Outcomes	<p>1. Fully immunised child by 12 months of age, defined as receiving BCG, 3 doses of OPV, 3 doses of pentavalent vaccine, and measles vaccine; 2. Timely fully immunised child defined as being fully immunised within 2 weeks of the measles EPI due date; 3. Carer's opinion on number of SMS reminders per vaccine and about the influence of SMS reminder on decision to vaccinate the child</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Timely vaccinations for individual vaccines defined as receiving pentavalent 1, 2, and 3 vaccine; OPV; or measles vaccines within 2 weeks (we included overall measure instead of individual measures per vaccine); 2. Receiving all timely vaccines (children who received pentavalent 1, 2, and 3, and measles vaccines within 2 weeks of their respective EPI due date) (we included fully immunised child by measles due date instead (see above))</p> <p>Outcome assessment time point: 12 months (1 time point for all outcomes)</p>
Funding/declarations of interest	<p>Funding: Bill & Melinda Gates Foundation</p> <p>Conflicts of interest: none declared</p>
Notes	<p>*SMS plus incentives (KES 75 or KES 200) groups were not considered relevant for this systematic review and were not extracted.</p> <p>Trial ID: NCT01878435</p> <p>Cluster features: 152 villages were randomised. The study adjusted outcomes for cluster effect using coefficient of variation (k) = 0.089 in the control group; k = 0.069 in the SMS only group.</p>
Risk of bias	
Bias	Authors' judgement Support for judgement

Gibson 2017 (Continued)

Random sequence generation (selection bias)	Low risk	Computer generated. Quote: "A constrained randomisation was done with GAUSS Mathematical and Statistical System by one of the study investigators, which randomly generated 1000 allocations."
Allocation concealment (selection bias)	Low risk	The 1000 sequences were labelled with 3-digit numbers, 000 to 999, each 1 assigning 38 villages to each of the 4 groupings (A–D). At a public randomisation ceremony on 12 September 2013, village chiefs determined the final randomisation outcome by picking numbered balls from a cloth sack to select 1 of these 1000 sequences, then picking labelled (study group) balls to assign the interventions to the chosen allocation.
Selective cluster recruitment	High risk	Enrolment of participants took place after cluster allocation had been made. Quote: "Birth notifications were relayed to field-based community interviewers who then screened caregivers of newborns for eligibility into the study." Quote: "Participants provided written informed consent and were enrolled into the study by community interviewers after villages were randomly assigned."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Due to nature of intervention and study design, study participants were not blinded to their study group allocation. Field staff were not informed of a village's allocation, but this could be inferred from some enrolment and follow-up survey questions.
Blinding of objective outcome assessment (detection bias)	Low risk	Data cleaning was performed by a statistician blinded to the allocation.
Blinding of subjective outcome assessment (detection bias)	Unclear risk	Field staff were not informed of a village's allocation, but this could be inferred from some enrolment and follow-up survey questions.
Incomplete outcome data (attrition bias) All outcomes	Low risk	14% in the SMS only arm and 19% in the control arm were lost to follow-up (out-migration and death), 4.4% in the SMS only arm and 7.4% in the control arm reported immunisation verbally (no vaccination booklet) and were excluded. Primary analyses were performed with modified ITT analyses at the participant level so that participants' outcomes were analysed regardless of the degree of exposure to study interventions. The term modified refers to the requirement of being able to determine the 12-month immunisation outcomes.
Selective reporting (reporting bias)	Low risk	Table and figures reporting all outcomes stated in the paper and online trial registration (NCT0187843).
Other bias	Low risk	Sociodemographic characteristics of the analytic sample were similar across the 4 groups.

Haji 2016
Study characteristics

Methods **Aim:** to assess the impact of SMS and sticker reminders to reduce dropout rates for routine childhood vaccinations, and determined factors associated with missed vaccination in selected districts in Kenya.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Haji 2016 (Continued)

	<p>Study design: cluster RCT</p> <p>Recruitment: children aged < 12 months who were brought to the selected vaccinating health facilities in the 3 districts for their first dose of pentavalent vaccine were recruited on a first-come basis until the strategy-level target sample sizes was reached.</p> <p>Study duration: 8 weeks</p> <p>Study dates: February 2014 to December 2014 (enrolment ceased October 2014)</p>
Participants	<p>Inclusion criteria: children aged < 12 months attending for first dose of pentavalent vaccine</p> <p>Sample size: 1116 (SMS: 372 participants, 3 clusters; targeted non-digital communication (stickers): 372 participants, 3 clusters; usual care: 372 participants, 3 clusters)</p> <p>Age (mean): carers: 26 (range 14–45) years</p> <p>Sex (male): infants = 51%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: community-based, linked to 9 vaccinating health facilities in 3 districts (Machakos, Langata, and Njoro), 3 facilities in each district</p>
Interventions	<p>Intervention: SMS reminders sent from an automated web-based system in Kiswahili and English, sent 2 days before (reminding the date of next vaccine and which health facility to attend) and on the day of the scheduled vaccination for the second and third doses of pentavalent vaccine.</p> <p>Content: SMS reminders</p> <p>Frequency and intensity: 4 reminders (2 days before and on day of both second and third dose), reminders were at 10 weeks (–2 days) until 14 weeks</p> <p>Control: 1. standard care/no intervention. 2. targeted non-digital communication: reminder sticker</p> <p>Co-interventions: scheduled vaccination due date was indicated on the child's health booklet as per routine procedures. All groups received routine health education and advice on vaccination.</p>
Outcomes	<p>1. Receipt of pentavalent vaccine (at 10 and 14 weeks of age); 2. Costs</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Mean days delay in receiving vaccine doses (at 10 and 14 weeks of age) (outcome not eligible for inclusion); 2. Factors associated with missed vaccination (not included in review, results were not reported per group); 3. Did not return for vaccinations (not included in review, those receiving vaccinations were extracted)</p> <p>Outcome assessment time point: 10 and 14 weeks of age corresponding to the second and third doses of the pentavalent vaccine</p>
Funding/declarations of interest	<p>Funding: US Center for Disease Control and Prevention</p> <p>Conflicts of interest: authors declared no competing interests.</p>
Notes	<p>Districts were selected that had > 10% dropout rate for third dose of pentavalent vaccine.</p> <p>Trial ID: not reported</p> <p>Cluster features: 9 health facilities were randomised. Mean cluster size: 124. The study did not adjust for clustering effect for any of the relevant outcomes; we used ICC 0.0487 (derived from $k = 0.089$ reported in Gibson 2017) to adjust for cluster effect in the analyses.</p>

Risk of bias
Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Haji 2016 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "We selected three health facilities in each district, and randomly allocated each facility ..." Comment: no further details reported.
Allocation concealment (selection bias)	Unclear risk	No information on allocation concealment.
Selective cluster recruitment	High risk	Convenient enrolment of participants likely to have taken place after cluster allocation to intervention arms. Quote: "We selected three health facilities in each district, and randomly allocated each facility to one of the two interventions ... or to serve as the control group ... Participants were conveniently enrolled in the selected health facilities until the strategy-level target sample sizes were reached."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind; participants and personnel were aware of cluster allocations for data collection purposes.
Blinding of objective outcome assessment (detection bias)	High risk	Personnel were aware of cluster allocations for data collection purposes.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Flow of participants and clusters through the study was not clearly reported.
Selective reporting (reporting bias)	Low risk	Outcomes are reported exhaustively, including for acknowledged limitations.
Other bias	High risk	Contamination: acknowledged limitation was that (quote) "If a care giver took the child to another facility for second or third pentavalent dose, the system considered the child unvaccinated." Follow-up phone-calls determined that 35 infants had been taken to other centres. Baseline differences: there were no statistical differences in demographic characteristics among carers and children enrolled in each of the 3 groups.

Hannan 2016
Study characteristics

Methods	Aim: to test the effects of a NP 2-way communication mobile phone and texting follow-up intervention for the first 6 months postbirth in low-income first-time mothers and their healthy full-term infants. Study design: parallel individual RCT
---------	--

Hannan 2016 (Continued)

Recruitment: first-time low-income mothers were enrolled from the Mother Baby Unit at Jackson Memorial Hospital in Miami, FL.

Study duration: 6 months

Study dates: not reported

Participants

Inclusion criteria: first-time mothers, aged 18 years, any racial/ethnic group, low-income, understood spoken English or Spanish, access to a mobile phone, delivered a singleton healthy full-term infant, only infants with limited access to health care were included.

Sample size (n): 141 (intervention: 63; control: 66)

Age (mean): mothers: 25.3 (SD 5.6) years

Sex (male): infants: 55%

Country: USA (high-income)

Setting: community, linked to hospital mother and baby unit

Interventions

Intervention: 2-way NP mobile phone intervention that included calls and SMS messages

Content: calls and text messages assessing health problems or concerns regarding the infant

Frequency and intensity: mobile phone contact and texting on posthospital discharge days 3, 7, 14, 21, then monthly for 6 months. Additionally, intervention group mothers were able to contact the NPs by mobile phone or texting Monday to Saturday

Control: standard care/no intervention. Routine hospital discharge care

Co-interventions: if a healthcare concern was stated, the NP followed US paediatric protocols to implement care. If the mother voiced a serious infant complaint such as fever, excessive crying, vomiting, lethargy, or seizure-like activity, the mother was instructed to contact the 911 emergency systems. Backup paediatricians were available to the NPs for consultation on infant health concerns.

Outcomes

1. Immunisations up to date (at 2, 4, 6 months – we included data for the longest time point (6 months)); 2. Infant emergency room attendance (6 months)

Outcomes reported but not included in review:

1. Infant hospitalisations (6 months) (outcome not included, for healthcare attendance we included immunisations and emergency department visits); 2. Infant urgent care seeking (6 months) (outcome not included, for healthcare attendance we included immunisations and emergency department visits); 3. Time in days to first well visit (postnatal care appointment) – outcome not eligible for inclusion; 4. Mothers' perceived stress (posthospital discharge) – outcome not eligible for inclusion; 5. Mothers' perceived social support (posthospital discharge) – outcome not eligible for inclusion; 6. Infant well visits received late (first visit, month 1, month 2, month 4, month 6) – outcome not eligible for inclusion

Outcome assessment time point: specified above after each outcome

Funding/declarations of interest

Funding: funded by MBRS Score National Institute of Health; Eunice Kennedy Shriver National Institute of Child Health & Human Development.

Conflicts of interest: author(s) declared no potential conflicts of interest.

Notes

Trial ID: not reported

Risk of bias
Bias

Authors' judgement **Support for judgement**

Hannan 2016 (Continued)

Random sequence generation (selection bias)	Low risk	Quote: "mothers were randomized to a control or intervention group using a table of random numbers."
Allocation concealment (selection bias)	Unclear risk	No information on allocation concealment.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants, no information on blinding of study personnel.
Blinding of subjective outcome assessment (detection bias)	High risk	Measures collected from mothers were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	12 mothers were lost to contact postdischarge due to disconnected mobile phones. It is not reported how many from each group.
Selective reporting (reporting bias)	Unclear risk	All expected outcomes and all outcomes listed in the methods section were reported, but no protocol or online trial record available to check against.
Other bias	Unclear risk	There were no significant differences in the demographic characteristics between the groups except for the number of years in the US.

Hofstetter 2015a
Study characteristics

Methods	<p>Aim: assessed the impact of text message reminders to parents on timely MMR vaccination.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: parents/children identified through the hospital registration system and its immunisation registry. The registration system included demographic and visit data for participants, and all vaccine doses administered to participants at the hospital and affiliated clinics through the electronic health record.</p> <p>Study duration: 2.5–3.5 months</p> <p>Study dates: June 2011 to October 2012</p>
Participants	<p>Inclusion criteria: parents were eligible for participation if their child: was age 9.5–10.5 months, had a participating clinic visit in the past 6 months, had a mobile phone number listed in the hospital registration system.</p> <p>Sample size (n): 2054 (SMS only: 686; scheduling + SMS: 686; usual care: 682)</p> <p>Age: parents' age not reported, children: 9.5–10.5 months at enrolment</p> <p>Sex (female): SMS only: 48.1%; scheduling + SMS: 49.4%; usual care: 49.0%</p> <p>Country: USA (high-income)</p> <p>Setting: community, linked to 4 paediatric practices in an ambulatory care network affiliated with a large academic medical centre in New York City.</p>

Hofstetter 2015a (Continued)

Interventions

Intervention: reminder SMS message 2 days before scheduled appointment. Scheduled SMS messages not sent if a previously arranged appointment was detected in the system.

Content: SMS message reminded parents 2 days before a scheduled 1-year appointment, informed them that doctor would discuss needed vaccines, and asked them to remember to bring the child's vaccination card. SMS messages included the clinic contact information and mentioned the child's need for important vaccines such as measles following the first birthday.

Frequency and intensity: 1 reminder SMS 2 days before child's 1-year vaccination. Up to 3 weekly SMS after randomisation (at child age of 9.5–10.5 months) followed by 1 reminder 2 days before child's 1-year vaccination.

Control: standard care/no intervention. The usual care arm received no text message reminders.

Co-interventions: children in all arms received "usual care," which included a routine automated telephone appointment reminder provided directly from the clinic network 1 day before existing appointments.

Outcomes

1. Child receives MMR vaccine between 361 days and 13 months of age; 2. Client satisfaction (13 months)

Outcomes reported but not included in the review:

1. 1-year preventive care visit scheduled at 11–13 months of age (outcome not eligible for inclusion); 2. 1-year preventive care visit attended 11–13 months of age (outcome not eligible for inclusion); 3. MMR vaccination by 16 months of age (only overall proportion and P value were reported – not included in review, 13-month time point was included)

Outcome assessment time point: 13 months

Funding/declarations of interest

Funding: supported by National Center for Immunization and Respiratory Diseases of the Centers for Disease Control and Prevention.

Conflicts of interest: unclear: under 'Funding' the authors have listed personal research funding, apparently linked to a different study: quote: "Dr. Hofstetter receives research support for an investigator initiated study funded by the Pfizer Medical Education Group."

Notes **Trial ID:** NCT01199666

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomised (quote): "Using a random sample algorithm generator in SPSS 19.0."
Allocation concealment (selection bias)	Low risk	Randomisation and delivery of intervention automated and integrated within the hospital registration system and its immunisation registry.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants were not aware that they were in a study. Quote: "The study was approved by the Columbia University Medical Center Institutional Review Board with a waiver of consent." Intervention delivered by an automated "text messaging platform integrated with the hospital registration system and its immunization registry."
Blinding of objective outcome assessment (detection bias)	Low risk	Data were retrieved electronically from hospital and registry records. Quote: "Study analysts were blinded to group assignments."

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Hofstetter 2015a (Continued)

Blinding of subjective outcome assessment (detection bias)	Low risk	Participants and study analysts were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Medical and vaccination records available for all participants.
Selective reporting (reporting bias)	Unclear risk	Online trial record available, but all outcomes listed in methods were reported.
Other bias	Low risk	Baseline demographic characteristics were similar between arms.

Jareethum 2008
Study characteristics

Methods	<p>Aim: to compare the satisfaction levels of ANC between healthy pregnant women who received SMS via mobile phone for antenatal support, and those who did not.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: all pregnant women who received ANC and planned to give birth at the study site.</p> <p>Study duration: from 28 weeks of gestation until delivery</p> <p>Study dates: May 2007 to October 2007</p>
Participants	<p>Inclusion criteria: aged > 18 years, no medical diseases or obstetric complications, singleton pregnancy, gestational age < 28 weeks (confirmed by ultrasound) when enrolled in the present study, owner of a mobile phone, able to receive and understand SMS messages</p> <p>Sample size (n): 68 (intervention: 34; standard care: 34)</p> <p>Age (mean): intervention: 28.72 (SD 4.9) years; standard care: 25.97 (SD 6.1) years</p> <p>Sex (female): 100%</p> <p>Country: Thailand (upper middle-income)</p> <p>Setting: Siriraj Hospital, Bangkok</p>
Interventions	<p>Intervention: SMS messages appropriate to the women's gestational age</p> <p>Content: information and warnings relating to abnormal symptoms</p> <p>Frequency and intensity: twice a week (Mondays and Thursdays during the daytime)</p> <p>Control: standard care/no intervention</p> <p>Co-interventions: all participants received phone calls at 32 weeks' gestation to check both groups were still contacted by the Siriraj antenatal clinic and to confirm that participants could receive and understand SMS messages. All participants received routine antenatal and perinatal care.</p>
Outcomes	<p>1. Preterm delivery; 2. Birth weight; 3. Gestational age at birth; 4. Patient satisfaction with intervention (antenatal and perinatal periods)</p> <p><u>Outcomes reported but not included in the review:</u></p>

Jareethum 2008 (Continued)

1. Self-reported level of confidence (outcome not eligible for inclusion); 2. Anxiety (scale not verified, not eligible for inclusion); 3. Route of delivery (vaginal or Caesarean) (outcome not eligible for inclusion)

Outcome assessment time point: postpartum

Funding/declarations of interest	Funding: not reported Conflicts of interest: not reported
----------------------------------	--

Notes	Trial ID: not reported
-------	-------------------------------

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The pregnant women who participated were randomly allocated into 2 groups using a table of random numbers.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No attempts to blind the participants or personnel were reported. Not possible to blind participants due to the nature of the intervention.
Blinding of objective outcome assessment (detection bias)	Low risk	It was not reported whether outcome assessors were blinded, but objective pregnancy outcomes (preterm delivery, birth weight, route of delivery, etc.) were collected from the obstetric records at the postpartum ward and were thus considered at low risk of bias.
Blinding of subjective outcome assessment (detection bias)	High risk	Subjective outcomes (client satisfaction, confidence, anxiety) were self-reported by unblinded participants using a questionnaire.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Available-case analysis with low attrition (10%). 2 (6.3%) women in study group and 2 (6.9%) women in control group changed to deliver at another hospital. 2 (6.9%) women were lost during the follow-up and 1 (3.4%) woman had her pregnancy aborted before 28 weeks of gestation. The remaining 32 women in study group and 29 in control group were included in analyses.
Selective reporting (reporting bias)	Low risk	All relevant outcomes in the method section were reported in the results section.
Other bias	Low risk	No baseline differences between groups.

Joshi 2015
Study characteristics

Methods	Aim: to conduct an RCT to test and document the effectiveness of 2 mHealth interventions provided to target women. Study design: cluster RCT
---------	---

Joshi 2015 (Continued)

Recruitment: a list of all villages in Osmanabad, Solapur, and Washim districts was sent to biostatisticians to randomly select 100 villages each from Osmanabad and Solapur (being larger districts) and 50 villages from Washim district. Then 1743 pregnant women from the 250 villages were enrolled and divided into 2 groups.

Study duration: 11–39 weeks of pregnancy

Study dates: not reported

Participants

Inclusion criteria: women in project area were enrolled from the third or fourth month of pregnancy.

Sample size: 1743 (intervention: 1162 participants, 166 clusters; control: 581 participants, 84 clusters*)

Age: not reported

Sex (female): 100%

Country: India (lower middle-income)

Setting: home-based care

Interventions

Intervention: twice-a-week mobile phone voice messaging service in local dialect that disseminates targeted, timely, and culturally sensitive preventive information directly to the pregnant women. Content of voice messages as well as animation films were culturally appropriate; timed and targeted as per beneficiary's gestational age; sent in a user specified language and time slot. The range of functions served by mHealth tools included: client education and counselling; diagnostic alerts; information giving; actionable tips to pregnant women for self-care and foetal health; and encouragement to clients for behaviour change.

Content: the voice messaging service was supported by short educational animation film clips displayed on mobile phones for reinforcement of preventive health information.

Frequency and intensity: twice a week from 11–39 weeks of pregnancy

Control: standard care/no intervention.

Co-interventions: none reported

Outcomes

1. Proportion of pregnant women who took iron and folic acid tablets for 100 days; 2. Proportion of births attended by skilled birth assistant/skilled health personnel; 3. Client satisfaction

Outcomes reported but not included in the review:

1. Knowledge gain (outcome not eligible for inclusion)

Outcome assessment timepoint: not reported

Funding/declarations of interest

Funding: Department for International Development (DFID), UK

Conflicts of interest: not reported

Notes

*number of clusters in the control group was not reported but we assumed 84 since the total was 250 and there were 166 intervention clusters.

Trial ID: not reported

Cluster features: 250 villages were randomised. Mean cluster size: 6.97. The study did not adjust for clustering effect; we used ICC 0.041 for skilled birth attendance and ICC 0.154 for iron and folate tablet adherence (reported in [Pagel 2011](#)) to adjust for cluster effects in the analyses.

Risk of bias
Bias
Authors' judgement
Support for judgement

Joshi 2015 (Continued)

Random sequence generation (selection bias)	Unclear risk	Villages were randomly selected by a biostatistician, but it is unclear how the participants were randomised.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Selective cluster recruitment	Unclear risk	Unclear whether participants were enrolled before or after clusters had been randomised. Quote: "In the next step, 1743 pregnant women from 250 villages enrolled in RCT were divided into two groups."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The flow of participants and clusters through the study was not clearly reported.
Selective reporting (reporting bias)	Unclear risk	No protocol or trial registration was mentioned and few details were reported in the paper to determine the risk of reporting bias.
Other bias	Unclear risk	Very few details were mentioned to determine any other risk of bias.

Kamau-Mbuthia 2013
Study characteristics

Methods	<p>Aim: to investigate pathways through which mobile phone-based support may promote exclusive breastfeeding.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: women in third trimester attending ANC</p> <p>Study duration: up to 3 months postpartum</p> <p>Study dates: not reported</p>
Participants	<p>Inclusion criteria: low-income women in the third trimester attending ANC at a large hospital</p> <p>Sample size (n): 505 (intervention: 153; control: 179)*</p> <p>Age: not reported</p> <p>Sex (female): 100%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: facility based</p>
Interventions	<p>Intervention: mobile phone-based peer support provided by trained peer leaders</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Kamau-Mbuthia 2013 (Continued)

Content: not reported

Frequency and intensity: not reported

Control: standard care/no intervention. existing facility-based standard of care

Co-interventions: not reported

Outcomes

1. Receiving postpartum help with breastfeeding; 2. Neonatal morbidity: infant diarrhoeal disease; 3. Infant length and weight (no SD reported); 4. Mastitis: breast pain and engorgement

Outcomes reported but not included in the review:

1. Any infant illness (outcome not included, for neonatal morbidity we included 'infant diarrhoeal disease' (see above)); 2. Receiving infant feeding advice from HCWs or others (outcome not included, we included 'receiving postpartum help with breastfeeding' (see above)); 3. Give medicines, vitamins, and minerals (outcome not eligible for inclusion); 5. Return to work by 3 months postpartum (results were not reported)

Outcome assessment time point: 3 months postpartum (1 time point for all outcomes)

Funding/declarations of interest

Funding: Bill & Melinda Gates Foundation through the Alive and Thrive small grants programme

Conflicts of interest: not reported

Notes

Trial ID: not reported

*1 study arm (peer support group) was not eligible for inclusion in the review.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Randomized", no further details reported.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-report by mothers who were not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not reported.
Selective reporting (reporting bias)	High risk	Not all outcomes were reported (return to work by 3 months postpartum).
Other bias	Unclear risk	The study was reported in 2 conference abstracts with limited information to determine any other risk of bias.

Kassaye 2016
Study characteristics

Methods	<p>Aim: to determine the utility of text messages to improve uptake of antenatal and PMTCT service.</p> <p>Study design: cluster RCT</p> <p>Recruitment: HIV-positive women presenting for ANC</p> <p>Study duration: from enrolment (< 32 weeks' gestation) to 6–8 weeks postpartum</p> <p>Study dates: June 2012 to June 2013</p>
Participants	<p>Inclusion criteria: HIV-positive pregnant women, < 32 weeks' gestational age, were not currently receiving ARV therapy, were planning to remain in the area for the duration of the study period, and agreed to follow-up of their infants until 6 weeks following delivery</p> <p>Sample size: 550 (intervention: 280 participants, 13 clusters; control: 270 participants, 13 clusters)</p> <p>Age (median): intervention: 25.5 (IQR 21–29) years; control: 25.6 (IQR 22–29) years</p> <p>Sex (female): 100%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: community, health facilities</p>
Interventions	<p>Intervention: automated text messages</p> <p>Content: PMTCT services including appointment reminders and adherence support, motivational messages, male-partner involvement and engagement in delivery planning and essential child health messages including warning signs and nutrition</p> <p>Frequency and intensity: 3–6 text messages each week (< 32 weeks' gestation to 6–8 weeks postpartum)</p> <p>Control: standard care/no intervention</p> <p>Co-interventions: none</p>
Outcomes	<p>1. ARV adherence – any maternal missed doses in the past week (6–8 weeks postpartum, we did not include earlier time points 36 weeks of gestation and at delivery); 2. Birth in health facility; 3. infant ARV treatment adherence (6 weeks postpartum); 4. infant HIV test (6 weeks postpartum); 5. infant HIV DNA PCR test positive (6 weeks postpartum); 6. number of face-to-face or mobile phone communications (delivery)</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Mother on combination ARV therapy (time points: 34–36 weeks' gestation, 7 days, and 6–8 weeks postpartum, outcome not eligible for inclusion)</p> <p>Outcome assessment time point: see above</p>
Funding/declarations of interest	<p>Funding: WHO through a grant to the Elizabeth Glaser Pediatric AIDS foundation and the Elizabeth Glaser Pediatric AIDS Foundation.</p> <p>Conflicts of interest: no competing interests to report</p>
Notes	<p>Trial ID: NCT01645865</p> <p>Cluster features: 26 healthcare facilities were randomised. Mean cluster size: 21.15. The study adjusted some outcomes (mother taking antenatal and postnatal ARV) for cluster effect and confounding variables (participant age, gestational age, whether the woman was newly diagnosed with HIV, and disclosure of HIV status to her partner and family). For the remaining outcomes, we used ICCs reported in</p>

Kassaye 2016 (Continued)

Page1 2011: ICC 0.127 (birth in health facility), ICC 0.055 (infant ARV, infant HIV test uptake, infant HIV test positive), and ICC 0.030 (communications between HCW and participant) to adjust for cluster effect in the analysis.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "All health facilities were randomly allocated to be an intervention or control site, stratified by high volume (hospitals) and medium and low volumes (health centers and dispensaries)." Comment: information on sequence generation were not reported.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Selective cluster recruitment	Unclear risk	Unclear whether participants were enrolled before or after clusters were allocated to intervention arms. Quote: "Consecutive eligible HIV-positive pregnant women presenting for ANC were invited to participate."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Unable to blind participants.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-report by unblinded participants.
Incomplete outcome data (attrition bias) All outcomes	High risk	Attrition at each stage of follow-up, no reasons provided, dropouts not included in analysis.
Selective reporting (reporting bias)	Unclear risk	Trial registry includes an additional secondary outcome that was not reported in the publication – time to initiation of ARV (clinicaltrials.gov/ct2/show/NCT01645865).
Other bias	High risk	There were baseline differences between groups.

Kebaya 2014
Study characteristics

Methods	<p>Aim: to compare self-reported adherence to infant nevirapine prophylaxis and retention in care in HIV-exposed infants randomised to 2-weekly mobile phone call vs control (no phone calls).</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: recruited from postnatal wards of 3 health facilities and randomised within 24 hours of birth</p> <p>Study duration: not reported</p> <p>Study dates: not reported</p>
---------	---

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Kebaya 2014 (Continued)

Participants	<p>Inclusion criteria: HIV-positive women and their infants</p> <p>Sample size (n): 150 (intervention: 75; control: 75)</p> <p>Age: not reported</p> <p>Sex (female): 100%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: recruited from postnatal wards of 3 health facilities in Kisumu, treatment was home based.</p>
Interventions	<p>Intervention: mobile phone calls</p> <p>Content: reminders on PMTCT messages</p> <p>Frequency and intensity: fortnightly</p> <p>Control: digital, non-targeted communication: "standard health care messages (no calls)"</p> <p>Co-interventions: infants received nevirapine prophylaxis for PMTCT</p>
Outcomes	<p>1. Attendance for postpartum care appointment (10 weeks); 2. Adherence to newborn PMTCT treatment (nevirapine) (6 weeks)</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Mode of infant feeding (mother report) (results were not reported); 2. Early HIV testing (results were not reported)</p> <p>Outcome assessment time point: 6, 10 weeks</p>
Funding/declarations of interest	<p>Funding: not reported</p> <p>Conflicts of interest: not reported</p>
Notes	<p>Limited information reported in 2 abstracts, no author contact details in abstracts.</p> <p>Trial ID: not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Report states the study was randomised but no further details provided.
Allocation concealment (selection bias)	Unclear risk	Details of allocation concealment not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Open-label RCT.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Limited information reported in 2 abstracts.

Kebaya 2014 (Continued)

Blinding of subjective outcome assessment (detection bias)	Unclear risk	Limited information reported in 2 abstracts.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Limited information reported in 2 abstracts.
Selective reporting (reporting bias)	High risk	Limited information reported in 2 abstracts. However, safe infant feeding and HIV testing were outcome measures mentioned for which results were not reported in any of the abstracts.
Other bias	Unclear risk	The study was reported in 2 conference abstracts with limited information to determine any other risk of bias.

Lund 2012
Study characteristics

Methods	<p>Aim: to examine the effect of the wired mothers' intervention on ANC, skilled delivery attendance, access to emergency obstetric care, and perinatal mortality.</p> <p>Study design: cluster RCT</p> <p>Recruitment: women who attended ANC at 1 of the 24 selected healthcare facilities were included in their first ANC visit and followed until 42 days after delivery.</p> <p>Study duration: until 6 weeks after delivery</p> <p>Study dates: March 2009 to March 2010</p>
Participants	<p>Inclusion criteria: 1. Clusters: health facilities with the highest number of ANC clients in 2008; availability of ≥ 1 midwife in the facility; mobile phone network coverage; 2. Participants: women aged 14–45 years; pregnant women attending ANC at 1 of 24 selected primary healthcare facilities</p> <p>Sample size: 2637 (intervention: 1351 participants, 12 clusters; control: 1286 participants, 12 clusters)</p> <p>Age group: < 19 years: intervention: 9%; control: 10%; 20–24 years: intervention: 25%; control: 26%; 25–29 years: intervention: 29%; control: 26%; 30–34 years: intervention: 20%; control: 22%; ≥ 35 years: intervention: 18%; control: 17%</p> <p>Sex (female): 100%</p> <p>Country: Tanzania (low-income)</p> <p>Setting: primary healthcare facilities</p>
Interventions	<p>Intervention: SMS + voucher. The frequency of the messages varied depending on the stage of the pregnancy.</p> <p>Content: simple health education and appointment reminders to encourage attendance to routine ANC, skilled delivery attendance, and postnatal care</p> <p>Frequency and intensity: early in the pregnancy, women received 2 messages a month, after gestational week 36, the intensity increased to 2 a week, until 6 weeks after delivery</p> <p>Control: standard care/no intervention. Standard antenatal, delivery, and postnatal services consisted of ≥ 4 ANC visits, skilled attendance at delivery, and a postnatal visit within the first 48 hours for deliveries taking place outside healthcare facilities.</p>

Lund 2012 (Continued)

Co-interventions: if the women could not provide a phone number, they received only the mobile phone voucher component. All enrolled women were offered the standard antenatal, delivery, and postnatal services.

Outcomes 1. Maternal deaths; 2. Skilled attendant at birth (adjusted ORs split by rural and urban populations); 3. Neonatal death; 4. Attendance \geq 4 ANC appointments (outcome was selected among attendance at 0 to $>$ 5 appointments because it is the local and global WHO recommendation); 5. Antenatal antitetanus vaccinations; 6. Antenatal preventive treatment for malaria; 7. Premature deliveries; 8. Severe obstetric complications (mother); 9. Client satisfaction with intervention (only reported in intervention group)

Outcomes reported but not included in the review:

1. Gestational age at last ANC visit (outcome not eligible for inclusion); 2. Antepartum referrals (outcome not eligible for inclusion); 3. Timing of the mentioned services (outcome not eligible for inclusion); 4. Caesarean section (outcome not eligible for inclusion); 5. Perinatal death (combination of neonatal death and stillbirth) (outcome not eligible for inclusion, we included neonatal deaths)

Outcome assessment time point: 6 weeks after delivery (1 time point for all outcomes)

Funding/declarations of interest **Funding:** Danish International Development Cooperation

Conflicts of interest: none declared

Notes **Trial ID:** NCT01821222

Cluster features: 24 primary healthcare facilities were randomised. Mean cluster size: 110. The study adjusted some outcomes (skilled attendant at birth, attendance at $>$ 4 ANC appointments, perinatal death, antenatal antitetanus vaccinations, antenatal preventive treatment for malaria) for cluster design effect. For the remaining outcomes, we used ICCs reported in [Pagel 2011](#): ICC 0.003 (maternal deaths, premature delivery, severe maternal complications) to adjust for cluster effect in the analyses.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Primary healthcare facilities were assigned by simple random allocation to the mobile phone intervention. Method not described.
Allocation concealment (selection bias)	Unclear risk	No details on allocation concealment reported.
Selective cluster recruitment	Low risk	All eligible women from the selected clusters were included. Quote: "Women who attended antenatal care at one of the 24 selected healthcare facilities were included on their first antenatal care visit and followed until 42 days after delivery. Women were eligible for study participation irrespective of their mobile phone and literacy status."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Neither study participants nor clinic staff blinded because of the nature of the intervention requiring overt participation.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Antenatal, intrapartum, and neonatal outcomes measured objectively.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No clusters were lost to follow-up and rate of participant loss to follow-up was low: 3.0% (40/1351) participants in the intervention group and 3.7% (47/1286) in the control group were lost to follow-up.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Lund 2012 (Continued)

Selective reporting (reporting bias)	Low risk	All outcomes stated in the online trial register were reported in 3 different publications.
Other bias	Low risk	No significant differences between intervention and control groups with regard to baseline characteristics. Cluster sizes were reported and some analyses were adjusted for clustering effects.

Maslowsky 2016
Study characteristics

Methods	<p>Aim: to evaluate the effects of a mobile phone-based intervention, providing postpartum health education and support + 30 days of telephone access to a dedicated on-call nurse, on postnatal maternal health behaviour, and maternal and infant health.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: study enrolled consecutive inpatient mothers after delivery of their newborn at a large public hospital or a community clinic with a birthing centre</p> <p>Study duration: 1 call + 30-day access to call nurse</p> <p>Study dates: 1 June 2012 (start of recruitment) to 31 August 2012 (end of recruitment)</p>
Participants	<p>Inclusion criteria: mothers aged ≥ 15 years, spoke Spanish, newborn not admitted to neonatal intensive care unit</p> <p>Sample size (n): 178 (intervention: 102; control: 76)</p> <p>Age (mean): intervention: 27.0 (SD 5.4) years; control: 25.5 (SD 6.2) years</p> <p>Sex (female): 100%</p> <p>Country: Ecuador (upper middle-income)</p> <p>Setting: community served by a large public hospital or a community clinic with a birthing centre.</p>
Interventions	<p>Intervention: telephone-based education intervention in 2 parts: 1. educational session administered by nurse via phone within 48 hours of hospital discharge; 2. Access to a nurse on-call during the first 30 days of the newborn's life</p> <p>Content: initial session: advice on newborn feeding, safe sleeping, environment, health red flags, recommended appointments, and excessive crying + advice on mother recognising/preventing infection, bleeding, pain, voiding, contraception, and mood. Second part: mothers could call the nurse to ask questions regarding their own or their newborn's health and care, as needed.</p> <p>Frequency and intensity: 1 call and 30 days of telephone access to receive support from a nurse</p> <p>Control: standard care/no intervention. Brief discharge instructions delivered by a nurse at the time of hospital or clinic discharge, typically including a newborn check-up within 1 week, a maternal follow-up visit within 6 weeks, and initiation of a family planning regimen. If mothers required additional support during the postnatal period, they were instructed to attend a local health centre or emergency department.</p> <p>Co-interventions: none</p>
Outcomes	<p>1. Self-reported breastfeeding; 2. Self-reported use of contraception; 3. Infant attended newborn check-up (reported by mother); 4. Self-reported attendance for postpartum care appointment (narrative result); 5. Acute episodes of illness in infants and mothers (reported by mother); 6. Client satisfaction (only intervention group)</p>

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Maslowsky 2016 (Continued)

Outcomes reported but not included in the review:

1. Infant attended 2-month check-up (outcome not eligible for inclusion)

Outcome assessment time point: 3 months (1 time point for all outcomes)

Funding/declarations of interest	<p>Funding: pilot grant from the University of Michigan Center for Global Health; 2 authors' salaries supported by Eunice Kennedy Shriver National Institute of Child Health and Human Development; 1 author supported in part by grants from the Breast Cancer Research Foundation and the National Cancer Institute.</p> <p>Conflicts of interest: none.</p>
Notes	Trial ID: not reported.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "... participants were assigned via a random number generator to either the intervention or the control group."
Allocation concealment (selection bias)	Unclear risk	No information on allocation concealment.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Blinding of participants not possible, no information on blinding of personnel.
Blinding of subjective outcome assessment (detection bias)	High risk	Outcomes were self-reported by unblinded participants. No information on blinding of study personnel.
Incomplete outcome data (attrition bias) All outcomes	High risk	High attrition rate: intervention: 27/102 (26%); control: 16/76 (21%).
Selective reporting (reporting bias)	Unclear risk	All outcomes in methods reported, but no protocol available.
Other bias	Low risk	Intervention and control groups did not significantly differ in any demographic (Table 2 in paper) or obstetric (Table 3 in paper) characteristic.

McConnell 2016
Study characteristics

Methods	<p>Aim: to pilot a programme implemented 3 days after delivery in which a checklist was used by a CHW to assess the health of mother and newborn and targeted health education was offered.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: women attending selected health facilities were approached by a CHW for recruitment after a normal delivery just prior to their discharge home.</p> <p>Study duration: 1 week (at day 7 after birth)</p>
---------	---

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

McConnell 2016 (Continued)

Study dates: April 2014 to October 2014

Participants	<p>Inclusion criteria: women aged 18–40 years, safe normal delivery of a live infant at a Jacaranda Health facility, reside within a 20 km radius from the Jacaranda Health's hospital facility in peri-urban Nairobi, provide 2 phone numbers</p> <p>Sample size (n): 104 (phone call: 41; home visit: 32; standard care: 31)</p> <p>Age (mean): phone call: 25.6 years; home visit: 25.9 years; standard care: 26.7 years</p> <p>Sex (female): 100%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: facility-based, i.e. at Jacaranda Health's 10-inpatient bed hospital in a peri-urban setting just outside Nairobi</p>
Interventions	<p>Intervention: phone/mobile call</p> <p>Content: CHWs call to screen for maternal and newborn danger signs, to deliver targeted postnatal health education, and to refer mothers and their newborns to facility-based care if necessary using a checklist to guide them through the process that was available in English and Kiswahili.</p> <p>Frequency and intensity: once (at day 3 after delivery)</p> <p>Control: 1. Standard care/no intervention. Standard care from Jacaranda clinic and did not receive any additional postpartum check-in from CHW. 2. Home visit. Same health screening and education as the intervention, but during CHW house calls</p> <p>Co-interventions: none</p>
Outcomes	<p>1. Exclusive breastfeeding (9 weeks postpartum, the study also reported on this outcome at 10 days postpartum, but we only included the longest follow-up); 2. Newborn postpartum care sought (10 days postpartum); 3. Maternal postpartum care sought (10 days postpartum); 4. Newborn health problems (10 days postpartum); 5. Maternal health problems (10 days postpartum); 6. Use of family planning method (9 weeks postpartum); 7. Newborn vaccination: ≥ 1 dose of OPV and pentavalent vaccines (9 weeks postpartum)</p> <p><u>Outcomes reported but not included in review:</u></p> <p>1. Days postpartum maternal care sought, among maternal care-seekers, mean (no SD was reported, outcome not eligible for inclusion) (10 days postpartum); 2. Days postpartum newborn care sought, among maternal care-seekers, mean (no SD was reported, outcome not eligible for inclusion) (10 days postpartum); 3. Postnatal knowledge (10 days postpartum, outcome not eligible for inclusion); 4. Breastfed ≥ 3 times in past 8 hours (10 days postpartum, 9 weeks postpartum) – outcome not included, we included exclusive breastfeeding (see above); 5. Combined outcome of health knowledge (10 days postdelivery, outcome not eligible for inclusion); 6. Combined outcome of health practices (10 days' and 9 weeks postdelivery, outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 10 days and 9 weeks postpartum</p>
Funding/declarations of interest	<p>Funding: Grand Challenges Canada</p> <p>Conflicts of interest: none declared</p>
Notes	<p>Trial ID: NCT02104635</p>
Risk of bias	
Bias	Authors' judgement Support for judgement

McConnell 2016 (Continued)

Random sequence generation (selection bias)	Low risk	Random assignment of patient identifiers was done using a randomisation sequence generated by the principal investigators.
Allocation concealment (selection bias)	Unclear risk	Participants were individually randomised prior to enrolment using numeric patient identifiers assigned by Jacaranda Health. A unique identifier was given to each Jacaranda Health client seeking any service (including ANC, delivery, postnatal care, and child wellness care) during the client's first visit to Jacaranda. No further details provided.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants to phone call, home visit, or no intervention.
Blinding of subjective outcome assessment (detection bias)	High risk	All outcomes collected through a survey with the mothers who were not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Quote: "We test for differences in outcomes across study arms on an intention-to-treat basis, with the treatment arms defined as participants' randomized treatment assignment and the sample including all respondents where data is available (not just those reached by a day 3 intervention)." Many participants could not be reached for outcome surveys, 16–22% for 10-day survey and 41–45% for 9-week survey.
Selective reporting (reporting bias)	Low risk	All study outcomes were explained and reported. See also ClinicalTrials.gov Identifier: NCT02104635.
Other bias	Unclear risk	There were some baseline differences. Quote: "Most of the differences in characteristics across arms are small and not statistically significant, except for a somewhat lower rate of marriage in the call arm (85 % (35/41) vs. 97 % (30/31) and 97 % (30/31) in the standard of care and home visit arms, respectively) and a lower employment rate in the visit arm (41 % (11/27) vs. 66 % (21/32) to 70 % (16/23) in the other arms)."

Moniz 2013
Study characteristics

Methods	<p>Aim: to estimate whether text messages encouraging influenza vaccination sent to an ambulatory obstetric population could improve influenza vaccination rates among women unsure about or unwilling to receive the vaccine.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: participants recruited and enrolled at routine obstetric visits to a Women's Hospital's outpatient clinic</p> <p>Study duration: 12 weeks</p> <p>Study dates: September 2010 (recruitment) to February 2012 (recruitment)</p>
Participants	<p>Inclusion criteria: pregnant women, < 28 weeks' gestation, aged 14–50 years, owned a personal mobile phone with text messaging capabilities, reported not receiving that season's influenza vaccine, electronic record lacked documentation of influenza vaccine administration in that season</p>

Moniz 2013 (Continued)

Sample size (n): 216 (intervention SMS: 104; control SMS: 100)

Age group: 13–17 years: control/general: 2 (2%), intervention/influenza: 3 (3%); 18–25 years: control/general: 67 (67%), intervention/influenza: 77 (74%); 26–49 years: control/general: 31 (31%), intervention/influenza: 24 (23%)

Sex (female): 100%

Country: USA (high-income)

Setting: community, served by a single women's hospital

Interventions	<p>Intervention: SMS messages</p> <p>Content: SMS messages regarding general preventive health in pregnancy plus the importance of influenza vaccination during pregnancy</p> <p>Frequency and intensity: 12 weekly messages</p> <p>Control: non-targeted, client communication: participants received 12 weekly text messages regarding general preventive health in pregnancy. General preventive health messages received by all participants covered topics such as the importance of antenatal vitamins, nutritional foods, and seat belt use during pregnancy</p> <p>Co-interventions: participants in both groups received usual ANC in the outpatient clinic, where ANC providers (nurses, midlevel HCPs, physicians) verbally recommend and offer influenza vaccination at each antenatal visit.</p>
Outcomes	<p>1. Attendance for antenatal influenza vaccination; 2. Acceptability of text messaging</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Self-reported reasons for declining influenza vaccination (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 12 weeks</p>
Funding/declarations of interest	<p>Funding: grant from the Amy Roberts Health Promotion Foundation</p> <p>Conflicts of interest: authors did not report any potential conflicts of interest.</p>
Notes	<p>Trial ID: NCT01248520</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Participants were randomised to the 2 study arms with equal frequency using a permuted block design with random block sizes of 2, 4, and 6. A researcher uninvolved in participant recruitment or clinical care generated the randomisation sequence.
Allocation concealment (selection bias)	Low risk	A researcher uninvolved in participant recruitment or clinical care generated the randomisation sequence and placed group assignments in sequentially numbered, sealed, opaque envelopes.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Both groups received SMS messages, but only 1 group's messages contained information about influenza vaccination. Uncertain to what extent women were cognisant that the influenza vaccine content was the intervention. HCPs were blind to the groups to which participants were randomised.

Moniz 2013 (Continued)

Blinding of objective outcome assessment (detection bias)	Low risk	Vaccine receipt was verified by review of the clinic's electronic medical record, which automatically documents date of vaccination at the time of vaccine administration. Record review was conducted after exit surveys were completed by a researcher unaware of participants' randomisation allocation.
Blinding of subjective outcome assessment (detection bias)	Unclear risk	Unclear whether participants were blinded, thus unclear whether self-reported measures were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Data complete for ITT analysis, all exclusions accounted for in per-protocol analysis.
Selective reporting (reporting bias)	Low risk	No reason to suspect selective reporting, online trial record was available and results for listed outcomes were reported.
Other bias	Low risk	No baseline differences between groups.

Naughton 2017
Study characteristics

Methods	<p>Aim: to 1. estimate the effectiveness of pregnancy smoking cessation support delivered by SMS text message, and 2. key parameters needed to plan a definitive trial.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: research midwives identified potential participants in antenatal clinics via their clinic notes or a screening questionnaire, and interested women were provided with participant information sheets.</p> <p>Study duration: 12 weeks</p> <p>Study dates: February 2014 (start of recruitment) to September 2014 (end recruitment)</p>
Participants	<p>Inclusion criteria: pregnant and < 25 weeks' gestation, smoking ≥ 5 cigarettes per day prepregnancy, smoking ≥ 1 cigarette on a typical day during pregnancy, aged ≥ 16 years, agreed to accept information to assist cessation, had own or had primary use of a mobile phone, familiar with sending and receiving text messages, able to understand written English, able to give informed consent</p> <p>Sample size (n): 407 (intervention: 2013; control: 204)</p> <p>Age (mean): intervention: 26.6 (SD 5.7) years; control: 26.4 (SD 5.7) years</p> <p>Sex (female): 100%</p> <p>Country: England (high-income)</p> <p>Setting: English NHS hospital antenatal clinics</p>
Interventions	<p>Intervention: SMS (MiQuit)</p> <p>Content: motivational messages, advice about preparing for a quit attempt, how to manage cravings and withdrawal, dealing with trigger situations, information about how smoking affects babies, and general encouragement</p> <p>Frequency and intensity: daily, according to a delivery schedule (0, 1, or 2 daily texts). Women were able to alter support frequency by texting the keywords MORE or LESS</p>

Naughton 2017 (Continued)

Control: standard care/no intervention

Co-interventions: all participants were given a standard NHS booklet on smoking cessation for mothers-to-be and could access smoking cessation information, advice, or support for stopping smoking offered as part of routine ANC.

Outcomes

1. Continuous abstinence (i.e. no more than 5 cigarettes in total) between 4 weeks postrandomisation until late pregnancy (approximately 36 weeks' gestation) (both self-report and objectively verified, we only included the objectively verified measure); 2. Client acceptability (SMS quite or extremely useful, SMS quite or extremely annoying); 3. Economic analysis: total per participant cost (of intervention); 4. Economic analysis: incremental cost per additional quitter

Outcomes reported but not included in the review:

1. 7-day point abstinence at 4 weeks postrandomisation and 36 weeks' gestation (both self-report and objectively verified) – we excluded these measures and included the long-term continuous abstinence outcome instead (see above); 2. Use of smoking cessation support during the trial period (outcome not eligible for inclusion); 3. Number of quit smoking attempts between baseline and late pregnancy (we excluded this smoking cessation outcome and included the long-term continuous abstinence outcome instead (see above))

Outcome assessment time point: assessed at approximately 36 weeks' gestation unless otherwise stated above.

Funding/declarations of interest

Funding: National Institute for Health Research (NIHR) under the Programme Grants for Applied Research programme; the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, and the Department of Health, under the auspices of the UK Clinical Research Collaboration.

Conflicts of interest: on 2 occasions since 2008, TC paid to attend and present at symposia arranged by Pierre Fabre Laboratories (PFL); PFL is a manufacturer of nicotine replacement therapy. All other authors had no competing interests.

Notes

Trial ID: NCT02043509

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomised individually to usual care or the MiQuit intervention in a 1:1 ratio using the Nottingham Clinical Trials Unit web-based system.
Allocation concealment (selection bias)	Unclear risk	Randomisation used a computer-generated pseudo-random code with random permuted blocks of randomly varying size, and stratification was by study site and gestation (< 16 vs ≥ 16 weeks). Following randomisation, unblinded trial team members sent arm-specific information packs to participants, which included the usual care booklet.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants.
Blinding of objective outcome assessment (detection bias)	Low risk	Trial staff involved in follow-up remained unaware of participants' treatments until questions on the intervention were asked at the end of the study, after smoking outcome data had been collected.

Naughton 2017 (Continued)

Blinding of subjective outcome assessment (detection bias)	High risk	Subjective outcomes self-reported by unblinded participants.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT analysis used. Participants lost to follow-up or with missing outcome data (around 30–40% depending on outcome and time point) were assumed to be smoking. Quote: "At 4 weeks, 295 (72%) participants provided smoking outcome data (68% MiQuit, 77% usual care). Further attrition in late pregnancy was fairly minimal, with 261 (64%) participants providing these outcome data (64% MiQuit, 65% usual care). Two hundred and thirty (57%) provided smoking outcome data at both time-points (55% MiQuit, 58% usual care) and 254 (62%) gave data used for smoking outcome 1 on abstinence between 4 weeks and late pregnancy (61% MiQuit, 64% usual care)."
Selective reporting (reporting bias)	Low risk	No evidence that outcomes were selectively reported.
Other bias	Unclear risk	Quote: "Participants' characteristics were similar in both groups, apart from that women randomized into the usual care group were more likely to reside in the most deprived (e.g. lower income) areas and have a non-smoking partner." Comment: for the main smoking cessation outcome. Quote: "We obtained validation samples for 37 of 64 (58%) of participants who reported abstinence at 36 weeks gestation (56% MiQuit, 61% usual care); with two (3.1%) and 15 (23%) participants providing only CO [carbon monoxide] or cotinine readings, respectively."

Niederhauser 2015
Study characteristics

Methods	<p>Aim: to examine the effect text messages immunisation reminders have on immunisation rates in the first 7 months of life.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: enrolment at health facilities</p> <p>Study duration: 5 months</p> <p>Study dates: not reported</p>
Participants	<p>Inclusion criteria: parent of a child aged 1–28 days, able to speak and read English, having a mobile telephone with text message capabilities, willingness to participate in study for 6 months</p> <p>Sample size (n): 57 (intervention: 30; control: 27)</p> <p>Age (mean): intervention: 31.5 (SD 6.9) years; control: 30.7 (SD 5.7) years</p> <p>Sex: 98% females (mothers). Just 1 father was included in the study.</p> <p>Country: USA (high-income)</p> <p>Setting: 4 health facilities: 1 large federally qualified health centre, 1 women/infants and children clinic, 1 private paediatric clinic, and the Honolulu Baby Expo, Hawaii</p>

Niederhauser 2015 (Continued)

Interventions	<p>Intervention: SMS</p> <p>Content: child immunisation reminders</p> <p>Frequency and intensity: 4 and 2 weeks prior to the due date for the infant's 2, 4, and 6-month vaccinations. From 2 to 7 months of age, i.e. 5 months' duration</p> <p>Control: digital, non-targeted communication. Messages were age-appropriate and based on well-baby information found in Bright Futures in addition to their routine reminders that they received from their HCPs.</p> <p>Co-interventions: 10% of the parents were randomly selected to receive phone calls verifying the receiving of the messages.</p>
Outcomes	<p>1. Immunisation compliance with 5 vaccines (DTaP; hepatitis B; haemophilus influenzae type B; pneumococcal conjugate vaccine; and polio) at 7 months of age</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Barriers to immunisation Survey (SHOTS survey), total score and subscales (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 2 months and 7 days, 2 months and 14 days, 4 months and 7 days, 4 months and 14 days, 6 months and 7 days, 6 months and 14 days, and 7 months of age – we reported on the longest (7 months) time point only.</p>
Funding/declarations of interest	<p>Funding: not reported</p> <p>Conflicts of interest: none declared</p>
Notes	<p>Trial ID: not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	After obtaining a written consent and contact information, using a random number table, parents were assigned to the intervention (30) or control group (27).
Allocation concealment (selection bias)	Unclear risk	Allocation concealment not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Blinding of participants and personnel not reported.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Not reported.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since the study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	The loss to follow-up of 39% of the intervention group compared to a loss of 10% of the control group, only per-protocol analysis was done.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Niederhauser 2015 (Continued)

Selective reporting (reporting bias)	Unclear risk	All outcomes described in the Measures section were included in the Results section, but no protocol or online trial record was available.
Other bias	Unclear risk	In comparing the 2 groups, the only difference in the 2 groups that nearly reached significance ($P = 0.07$), was a higher percentage of intervention (58%) than control (30%) were not married.

Odeny 2014
Study characteristics

Methods	<p>Aim: evaluated effect of an interactive, individually tailored, 2-way text messaging system informed by behavioural theory on maternal postpartum clinic attendance and infant HIV testing within 8 weeks after birth.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: not reported</p> <p>Study duration: from pregnancy to 6 weeks after delivery</p> <p>Study dates: April 2012 to March 2013</p>
Participants	<p>Inclusion criteria: aged ≥ 18 years, between 28 weeks' gestation and delivery, enrolled in PMTCT, planning to remain in study area, access to a mobile phone, and reported ability to read or had someone who read SMS on their behalf. Women who shared phones were eligible only if they had disclosed their HIV status to the person with whom the phone was shared.</p> <p>Sample size (n): 388 (intervention: 195; standard care: 193)</p> <p>Age (n): 18–24 years: intervention: 60 (30.8%); control: 65 (33.7%); 25–34 years: intervention: 111 (56.9%); control: 111 (57.5%); ≥ 35 years: intervention: 24 (12.3%); control: 17 (8.8%)</p> <p>Sex (female): 100%</p> <p>Country: Kenya (lower middle-income)</p> <p>Setting: ANC or HIV clinics at 5 health facilities, including a mix of rural and urban settings</p>
Interventions	<p>Intervention: text messages</p> <p>Content: educational or motivational messages tailored to gestation week or newborn age</p> <p>Frequency and intensity: 14 messages. Up to 8 were sent during pregnancy (weeks 28, 30, 32, 34, 36, 38, 39, and 40). Additional messages were sent weekly for the first 6 weeks after delivery.</p> <p>Control: standard care/no intervention</p> <p>Co-interventions: participants in both arms were allowed to call or send SMS to the study nurse at any time. Baseline treatments (n): on ART for own health: 101 (51.8%); 102 (52.8%); received ZDV prophylaxis: 85 (43.6%); 81 (42.0%); received ZDV + 3TC + nevirapine (delivery pack): 60 (30.8%); 53 (27.5%); received ZDV + 3TC (postdelivery pack): 60 (30.8%); 51 (26.4%); nevirapine prophylaxis for baby issued: 139 (71.3%); 133 (68.9%)</p>
Outcomes	<p>1. Infant HIV testing; 2. Infant positive virological HIV test results; 3. Maternal postnatal appointment attendance within 8 weeks; 4. Neonatal death/stillbirth</p> <p>Outcome assessment time point: 8 weeks postpartum</p>

Odeny 2014 (Continued)

Funding/declarations of interest

Funding: the National Institutes of Health Office of the Director, Fogarty International Center, Office of AIDS Research, National Cancer Center, National Eye Institute, National Heart, Blood, and Lung Institute, National Institute of Dental and Craniofacial Research, National Institute On Drug Abuse, National Institute of Mental Health, National Institute of Allergy and Infectious Diseases, and National Institutes of Health Office of Women's Health and Research through the Fogarty International Clinical Research Scholars and Fellows Program at Vanderbilt University and the American Relief and Recovery Act. TAO was a Fogarty International Clinical Research Fellow. CSC was supported by a Research Scientist Development Award from the National Institute of Mental Health, KY was supported by the University of Washington Center for AIDS Research (CFAR), an NIH funded programme.

Conflicts of interest: financial competing interests include but are not limited to paid employment or consultancy: CRC served as a paid consultant for CerMed Inc. to help them develop a barrier contraceptive/HIV prevention device. This consultancy ended in 2012. Research grants (from any source, restricted or unrestricted): CRC has active grants from the US NIH, CDC, and Bill & Melinda Gates Foundation. RSM has active grants from the US NIH and Hologic/Gen-Probe. Travel grants and honoraria for speaking or participation at meetings: CRC received a travel grant to consult with Gynuity on a study they conducted to investigate infections following medical abortion in the United States. Nonfinancial (professional): acting as an expert witness – CRC has served as an expert witness on a case in New York City involving a malpractice suit of a woman who died after delivery due to infectious complications. Membership in a government or other advisory board: CRC was a non-paid consultant on a WHO panel to assess the risk of hormonal contraception and HIV acquisition in women. RSM has received a donation of study product for treatment of vaginal infections from Embil Pharmaceutical Company.

Notes

Trial ID: not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomisation scheme with variable block sizes. Investigators and study staff were unaware of block numbers, sizes, or sequences.
Allocation concealment (selection bias)	Low risk	Intervention groups assigned using sealed, opaque envelopes.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Labelled as unblinded.
Blinding of objective outcome assessment (detection bias)	Low risk	Women's return visits and infant HIV testing data extracted from clinic records.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Analyses were ITT. There were few withdrawals or losses to follow-up.
Selective reporting (reporting bias)	Unclear risk	Expected outcomes reported, but no protocol or online trial register to check against.
Other bias	Low risk	Maternal characteristics balanced between study arms at baseline.

Omole 2018
Study characteristics
Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Copyright © 2020 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Omole 2018 (Continued)

Methods	<p>Aim: to determine the impact of pregnancy information and appointment reminder SMSs on maternal health behaviour.</p> <p>Study design: cluster RCT</p> <p>Recruitment: public launch of the project to sensitise health workers and select stakeholders followed by recruitment of random selection of pregnant women in their first trimester who were attending antenatal clinics in selected health facilities in the Ife-Ijesa zone.</p> <p>Study duration: participants followed up to delivery and completion of immunisation for their children</p> <p>Study dates: December 2013 (start of enrolment) to December 2014 (end of enrolment)</p>
Participants	<p>Inclusion criteria: pregnant women residing in Ife-Ijesa zone, owner of a mobile phone, able to read and write in English or Yoruba (local dialect) language</p> <p>Sample size: 508 (targeted SMS: 260 participants, 2 clusters; non-targeted SMS: 248 participants, 2 clusters)</p> <p>Age group: 15–24 years: 27.8%; 25–34 years: 59.3%; > 35 years: 13%</p> <p>Sex (female): 100%</p> <p>Country: Nigeria (lower middle-income)</p> <p>Setting: public health facilities in Ife-Ijesa area, Osun State, south-west Nigeria. These are secondary health facilities with the highest number of ANC attendees and delivery within the zone.</p>
Interventions	<p>Intervention: pregnancy information and appointment reminder text messages</p> <p>Content: messages for the intervention group included pregnancy-related information such as birth preparedness, complication readiness, and reminders of antenatal visits. Study participants in the intervention group also had the opportunity of sending text messages to the project team to seek for health information.</p> <p>Frequency and intensity: once a week</p> <p>Control: digital, non-targeted communication. General health tips, which excluded pregnancy-related health information and clinic schedule reminders</p> <p>Co-interventions: none</p>
Outcomes	<p>1. Place of birth (health facility); 2. Opinion of respondents about SMS intervention (only intervention group)</p> <p>Outcome assessment time point: at delivery (place of birth) or 6 weeks postpartum (self-reported questionnaire)</p>
Funding/declarations of interest	<p>Funding: The Madiro Fund: Gillian and Adrian Schauer Foundation, Montreal, Canada.</p> <p>Conflicts of interest: none declared</p>
Notes	<p>Trial ID: not reported</p> <p>Cluster features: 4 government health facilities were randomised. Mean cluster size: 127. Study did not adjust for clustering effect for the relevant outcome; we used ICC 0.127 (reported in Pagel 2011) to adjust for cluster effect in the analysis.</p>
Risk of bias	
Bias	Authors' judgement Support for judgement

Omole 2018 (Continued)

Random sequence generation (selection bias)	Unclear risk	Random sequence generation not mentioned.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment not mentioned/reported.
Selective cluster recruitment	High risk	Participants were enrolled after clusters had been allocated to intervention arms. Quote: "This phase started with the public launch of the project to sensitize health workers and select stakeholders. This was followed by recruitment of participants."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible to blind participants.
Blinding of objective outcome assessment (detection bias)	Unclear risk	Blinding of objective outcome assessment not mentioned/reported.
Blinding of subjective outcome assessment (detection bias)	Unclear risk	Blinding of subjective outcome assessment not mentioned/reported.
Incomplete outcome data (attrition bias) All outcomes	High risk	Table 4 shows primary outcome with high attrition.
Selective reporting (reporting bias)	High risk	Primary outcome of study was complete ANC defined as "attendance of at least 4 antenatal clinic visits and delivery in a health facility." The authors only reported on the proportions of delivery in a health facility for a subgroup of pregnant women.
Other bias	Unclear risk	Groups differed significantly in their occupational distribution, occupational category of spouses, proportion that had delivered in a health facility in their previous pregnancy, and in reasons for the choice of place of delivery.

Sharma 2011
Study characteristics

Methods	<p>Aim: to evaluate the effect of providing oral health education by 2 different modes (text messages and pamphlets), on the knowledge, attitude, and practices of mothers of preschool children and plaque scores of their children.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: not reported</p> <p>Study duration: 4 weeks</p> <p>Study dates: not reported</p>
Participants	<p>Inclusion criteria: mothers proficient in English, owners of a personal mobile phone, familiar with text messaging</p>

Sharma 2011 (Continued)

Sample size: 150 children enrolled, 143 completed study (SMS: 71; control: 72 at completion, numbers per group at enrolment not reported)

Age (mean): mothers: not reported; children: text messages: 3.6 (SD 0.5) years; pamphlet: 3.3. (SD 0.5) years

Sex: mothers: not reported; children: 83 boys/60 girls (among those who completed the study)

Country: India (lower middle-income)

Setting: not reported

Interventions	<p>Intervention: SMS</p> <p>Content: messages covered topics that would help the mothers to maintain an optimum oral health for their children and themselves</p> <p>Frequency and intensity: total of 21 messages, either in the form of text messages or pamphlets, were sent in 7 days (3 messages per day). Messages repeated every week for 4 weeks</p> <p>Control: non-digital, targeted communication. Pamphlets sent daily to the mothers through the children; 3 messages printed on the pamphlet and attached to the daily work diary of each child belonging to the pamphlet group</p> <p>Co-interventions: not reported</p>
Outcomes	<p>1. Visible Plaque Index in children</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>1. Knowledge, attitude, and practices of the mothers (outcome not eligible for inclusion)</p> <p>Outcome assessment time point: 4 weeks (1 time point for all outcomes)</p>
Funding/declarations of interest	<p>Funding: not reported</p> <p>Conflicts of interest: not reported</p>
Notes	<p>Trial ID: not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Participants were randomly allocated to the text message group or the pamphlets group. Method of generating sequence not described.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment not reported/mentioned.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Given nature of intervention, participants could not be blinded. The examiner and statistician who analysed data were blinded.
Blinding of objective outcome assessment (detection bias)	Low risk	Examiner and statistician who analysed the data were blinded.
Incomplete outcome data (attrition bias)	High risk	143/150 preschool children and their mothers completed the study. It was not reported which groups the dropouts belonged to.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Sharma 2011 (Continued)

All outcomes

Selective reporting (reporting bias)	Unclear risk	All outcomes described in the Measures section were included in the Results section, but no protocol or online trial record was available.
Other bias	Low risk	No apparent baseline differences between groups.

Stockwell 2015
Study characteristics

Methods	<p>Aim: to compare the effectiveness of text message vaccine reminders with and without vaccine education vs written reminder-only on receipt of second dose of influenza vaccine in young, low-income children.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: study sites were part of an ambulatory-care network staffed by 1 centrally administered paediatric group practice, all eligible children seen when research assistants were on-site were approached.</p> <p>Study duration: 42 days</p> <p>Study dates: August 2012 to March 2013</p>
Participants	<p>Inclusion criteria: parenting adult of child age 6 months to 8 years, child received care at study site (visit in last 12 months), child received 1 dose influenza vaccine and was in need of a second dose in same season season*, parent had mobile phone with text message capability, parent spoke English or Spanish and could read text messages</p> <p>Sample size (n): 660 (conventional SMS: 225; educational SMS: 216; usual care: 219)</p> <p>Age: parents: not reported; children: age group: < 5 years: 83% (550/660); 5–8 years: 17% (110/660)</p> <p>Sex (female): children: 49.5%</p> <p>Country: USA (high-income)</p> <p>Setting: 3 community-based paediatric clinics affiliated with New York-Presbyterian Hospital/Columbia University Medical Center in Northern Manhattan in New York City</p>
Interventions	<p>Intervention: 1. SMS vaccine reminders only; 2. SMS vaccine reminders and educational messages</p> <p>Content: SMS reminders contained the date after which the next dose was due and clinic-specific walk-in hours. Those in the educational text message arm additionally received educational information that included that the child was not protected until he or she received the second dose, that reaching full protection could take 2 weeks after second dose administration, and that doctors recommended a second dose. In addition, in 1 interactive message, parents could select to receive more information via text message. Messages were sent in English or Spanish based on the participant's request at enrolment.</p> <p>Frequency and intensity: 5 messages – 3 dates before dose was due (day 7, day 21, and day 25 after first influenza vaccine dose), on the day it was due (day 28), and 2 weeks after it was due (day 42)</p> <p>Control: standard care/no intervention. Parents of children randomly assigned to receive usual care did not receive any further intervention beyond the written reminder.</p> <p>Co-interventions: all families received a written reminder with the date the next influenza vaccine dose was due (routine practice)</p>

Stockwell 2015 (Continued)

Outcomes 1. Attendance for second vaccination before end of influenza season; 2. Receipt of a second influenza vaccine dose by 42 days postvaccination (i.e. on time); 3. Client satisfaction

Outcome assessment time point: see above

Funding/declarations of interest

Funding: institutional career development grant funded by the National Institutes of Health.

Conflicts of interest: Dr Hofstetter received support from the Pfizer Medical Education Group for a different investigator-initiated study; Dr Stockwell is a co-investigator but received no financial support; Dr Hofstetter also received support through the Investigator-Initiate.

Notes

Trial ID: NCT01662583

*Children in need of a second dose were those who had not received 2 doses of vaccine since July 2010 (the first season the 2009 H1N1-strain was included in the seasonal vaccine) or those who had not received 2 previous seasonal influenza vaccinations + $\geq 1 \times$ 2009 H1N1-containing vaccination, either as a seasonal or monovalent pandemic vaccine.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information on sequence generation. Quote: "Subjects were randomly assigned centrally with a 1:1:1 allocation at an individual level by using a permuted block design with a block size of 9, stratified by age and clinic site."
Allocation concealment (selection bias)	Low risk	Central allocation. Quote: "Subjects were randomly assigned centrally with a 1:1:1 allocation at an individual level by using a permuted block design with a block size of 9, stratified by age and clinic site."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Blinding of participants and personnel not reported/mentioned. However, due to the nature of the intervention participants (at least) could not be blinded to the intervention.
Blinding of objective outcome assessment (detection bias)	Low risk	Vaccination status retrieved from medical records and study analyst was blinded to individual group assignment.
Blinding of subjective outcome assessment (detection bias)	High risk	Self-reported measures were at high risk of detection bias since study was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Analyses were ITT, there was only 2 exclusions after randomisation but this was before the launching of the interventions.
Selective reporting (reporting bias)	Low risk	All outcomes listed in the online trial record (NCT01662583) were reported.
Other bias	Low risk	No significant baseline differences.

Yudin 2017

Study characteristics

Methods	<p>Aim: to evaluate if text message reminders increase the likelihood of receiving the influenza vaccine among pregnant women.</p> <p>Study design: parallel individual RCT</p> <p>Recruitment: women were approached in the waiting room by research personnel who were not part of the clinical care team and were asked if they were interested in enrolling in a study investigating the use of text messages in pregnancy. Women were told the messages would be about health-related behaviour in pregnancy, but no mention was made of influenza or vaccination at the time of recruitment.</p> <p>Study duration: 4 weeks</p> <p>Study dates: November 2014 (start recruitment) to March 2014 (end recruitment)</p>				
Participants	<p>Inclusion criteria: pregnant woman, aged ≥ 18 years, working mobile phone with ability to receive text messages, ability to speak or understand English</p> <p>Sample size (n): 317 (SMS: 153; standard care: 164)</p> <p>Age (mean): intervention: 32.2 (SD 4.5) years; control: 32.4 (SD 4.9) years</p> <p>Sex (female): 100%</p> <p>Country: Canada (high-income)</p> <p>Setting: hospital-based antenatal clinic at St Michael's Hospital, which is a women's health ambulatory care clinic in downtown Toronto</p>				
Interventions	<p>Intervention: SMS</p> <p>Content: SMS messages reinforcing that the influenza vaccine is recommended for all pregnant women and safe during pregnancy and breastfeeding</p> <p>Frequency and intensity: weekly (2 SMS/week)</p> <p>Control: standard care/no intervention. All women attending the clinic were given a pamphlet containing information about the risks of influenza during pregnancy, the importance of the vaccine for pregnant women, and the fact that the vaccine is safe in pregnancy and breastfeeding.</p> <p>Co-interventions: all women received usual ANC, including the verbal recommendation for influenza vaccination.</p>				
Outcomes	<p>1. Attendance for influenza vaccination; 2. Satisfaction with receiving SMS and opinion on timing and number of SMS received</p> <p><u>Outcomes reported but not included in the review:</u></p> <p>Outcome assessment time point: 6 weeks postpartum (1 time point for all outcomes)</p>				
Funding/declarations of interest	<p>Funding: St Michael's Hospital Innovation Fund Grant</p> <p>Conflicts of interest: not reported</p>				
Notes	<p>Trial ID: NCT02428738 (registered as a case control study)</p>				
Risk of bias					
Bias	<table border="1"> <thead> <tr> <th>Authors' judgement</th> <th>Support for judgement</th> </tr> </thead> <tbody> <tr> <td> </td> <td> </td> </tr> </tbody> </table>	Authors' judgement	Support for judgement		
Authors' judgement	Support for judgement				

Yudin 2017 (Continued)

Random sequence generation (selection bias)	Low risk	Computerised random number generator.
Allocation concealment (selection bias)	Low risk	Group allocation was assigned using sequentially numbered, sealed, opaque envelopes which were opened at the time of randomisation by study staff.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Women were told the messages would be about health-related behaviour in pregnancy, but no mention was made of influenza or vaccination at the time of recruitment. Medical and nursing staff caring for the women were blinded to study group allocation and were not involved in any aspects of the study.
Blinding of subjective outcome assessment (detection bias)	Low risk	Self-reported subjective outcomes among participants who were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	The final analysis was based on the actual treatment received and included 281/317 randomised participants. All analyses using ITT groupings were repeated and there were no changes in results when compared to the analyses based on treatment received (data available upon request).
Selective reporting (reporting bias)	Low risk	Outcomes listed in the online trial registration were reported.
Other bias	Unclear risk	No significant baseline differences.

3TC: lamivudine; ANC: antenatal care; ART: antiretroviral therapy; ARV: antiretroviral; BCG: Bacillus Calmette-Guérin; CHW: community health worker; DPT: diphtheria, pertussis (whooping cough), and tetanus; DTaP: diphtheria, tetanus, and whooping cough (pertussis); EPI: Expanded Programme on Immunisations; HCP: healthcare provider; HCW: healthcare worker; ICC: intracluster correlation coefficient; IQR: interquartile range; ITT: intention to treat; KES: Kenyan shilling; LMIC: low- to middle-income country; n: number of participants; NHS: National Health Service; NP: nurse practitioner; OPV: oral polio vaccine; OR: odds ratio; PCR: polymerase chain reaction; PMTCT: prevention of mother-to-child transmission; RCT: randomised controlled trial; SD: standard deviation; SMS: short message service; WHO: World Health Organization; ZDV: zidovudine.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Atukunda 2017	Irrelevant intervention – including digital tracking component.
Bracken 2014	Irrelevant intervention – not exclusively using mobile device.
Broberg 2013	Irrelevant intervention – not exclusively using mobile device.
Carlsen 2013	Irrelevant intervention – not exclusively using mobile device.
Collier 2005	Irrelevant intervention – not exclusively using mobile device.
Gallegos 2014	Irrelevant study design – not randomised controlled trial.
Haberer 2016	Irrelevant intervention – including digital tracking component.
Hashemian 2015	Irrelevant intervention – targeted communication used in conjunction with other interventions.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

101

Study	Reason for exclusion
Herring 2016	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Hofstetter 2015b	Irrelevant population – parents of children older than 5 years.
Irons 2015	Irrelevant study design – not randomised controlled trial.
Jimenez 2017	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Kofinas 2014	Irrelevant intervention – not targeted client communication.
Lau 2013	Irrelevant study design – not randomised controlled trial.
Lau 2014	Irrelevant study design – not randomised controlled trial.
Lewis 2012	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Maduka 2013	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Mauriello 2016	Irrelevant intervention – not exclusively using mobile device.
Milani 2015	Irrelevant intervention – not exclusively using mobile device.
Moore 2013	Irrelevant intervention – including digital tracking component.
Moore 2015	Irrelevant intervention – including digital tracking component.
Murthy 2019	Quasi-randomised study.
Mwapasa 2017	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Oakley-Girvan 2016	Irrelevant study design – not randomised controlled trial.
Patel 2014	Irrelevant intervention – not exclusively using mobile device.
Peitzmeier 2016	Irrelevant intervention – not exclusively using mobile device.
Pollak 2014	Irrelevant intervention – including digital tracking component.
Prieto 2016	Irrelevant study design – not randomised controlled trial.
Pérez-Ferre 2010	Irrelevant intervention – including digital tracking component.
Rampersaud 2016	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Rand 2015	Irrelevant population – parents of adolescents receiving vaccination reminders.
Rand 2017	Irrelevant population – parents of adolescents receiving vaccination reminders.
Reeder 2014	Irrelevant intervention – not exclusively using mobile device.
Reid 2014	Irrelevant study design – not randomised controlled trial.
Richman 2016	Irrelevant intervention – not exclusively using mobile device.
Robbins 2013	Irrelevant intervention – not exclusively using mobile device.

Study	Reason for exclusion
Sridhar 2013	Irrelevant intervention – not exclusively using mobile device.
Sridhar 2014	Irrelevant intervention – not targeted communication.
Stern 2013	Irrelevant intervention – not exclusively using mobile device.
Stockwell 2012	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Szilagyi 2013	Irrelevant intervention – not exclusively using mobile device.
Takeuchi 2016	Irrelevant intervention – targeted communication used in conjunction with other interventions.
Tarrant 2014	Irrelevant intervention – not exclusively using mobile device.
Trent 2013	Irrelevant study design – not randomised controlled trial.
Trent 2015	Irrelevant study design – not randomised controlled trial.
Van Ryswyk 2015	Irrelevant comparison group – control group also received targeted communication.
Wright 2012	Irrelevant intervention – not exclusively using mobile device.
Young 2013	Irrelevant intervention – not targeted communication.
Young 2014	Irrelevant intervention – not targeted communication.

Characteristics of studies awaiting classification *[ordered by study ID]*

Abroms 2017a

Methods	<p>Aim: to test whether an interactive and intensive text messaging programme, Quit4baby, can promote smoking cessation for pregnant women already enrolled in a health text messaging programme, Text4baby.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: July 2015 to February 2016</p>
Participants	497 pregnant smokers recruited among 'Text4baby' subscribers in the USA (except CA, OK, OH, and LA)
Interventions	<p>TCC: Quit4baby text messages + Text4baby text messages</p> <p>Control: Text4baby text messages</p>
Outcomes	Smoking cessation outcomes
Notes	Trial ID: NCT02412865

Abroms 2017b

Methods	Aim: to test the acceptability and feasibility of SmokefreeMOM, a national smoking cessation text-messaging programme for pregnant smokers.
---------	---

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Abroms 2017b (Continued)

	Study design: randomised controlled trial
	Study dates: September 2014 to May 2016 (recruitment)
Participants	99 pregnant smokers recruited from obstetrics-gynaecology clinics in Washington DC, USA.
Interventions	TCC: SmokefreeMOM text messaging programme Control: mailed self-help materials on quitting smoking while pregnant + 1 text message referring to telephone quit line
Outcomes	1. Programme acceptability and feasibility; 2. Use of treatments and resources for quitting at 1 month; 3. Smoking-related outcomes
Notes	Trial ID: NCT02412956

Altazan 2019

Methods	Aim: to assess the effect of a behavioural intervention targeting excess gestational weight gain. Study design: randomised controlled trial. Study dates: February 2013 to October 2015
Participants	54 pregnant women who were overweight or obese recruited through advertisements and targeted emails and referrals from local obstetricians in Baton Rouge, LA, USA.
Interventions	TCC 1: SmartMoms through mobile application TCC 2: SmartMoms in person Control: usual care
Outcomes	1. Weight; 2. Mood; 3. Quality of life
Notes	Trial ID: NCT01610752

Bangal 2018

Methods	Aim: to assess the influence of mobile communication between health facility and pregnant women on utilisation and outcome of maternal health services. Study design: randomised controlled trial Study dates: not reported
Participants	400 pregnant women in India (region not reported)
Interventions	TCC: mobile phone calls as reminders about next visit and SMS on maternal health Control: usual care
Outcomes	1. Antenatal care visits; 2. Institutional deliveries; 3. Perinatal mortality; 4. Complications during pregnancy; 5. Postnatal follow-up

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Bangal 2018 *(Continued)*

Notes	Trial ID: not reported
-------	------------------------

Domek 2019

Methods	Randomised controlled trial
Participants	Carers of children aged 6 weeks to 6 months who received first dose of primary vaccination series
Interventions	TCC: SMS message reminders 1 week before scheduled vaccination Control: usual care
Outcomes	1. Completion of vaccine primary series; 2. Factors associated with immunisation delays; 3. Acceptability of SMS vaccine reminders; 4. Timeliness of immunisation
Notes	Trial ID: NCT02567006

Forinash 2018

Methods	Aim: to evaluate the impact of text messaging on smoking cessation rates among pregnant women. Study design: randomised controlled trial Study dates: May 2014 to January 2016
Participants	49 pregnant women in the preparation stage of quitting smoking recruited from a maternal foetal care centre in Saint Louis, USA
Interventions	TCC: text messaging Control: usual care
Outcomes	1. Self-reported cessation rates at 1 month, as verified by exhaled carbon monoxide levels
Notes	Trial ID: NCT03024606

Gibson 2019

Methods	Aim: to determine if text message reminders, with or without mobile phone-based incentives, can improve measles immunisation coverage and timeliness in rural western Kenya. Study design: randomised controlled trial Study dates: enrolment to March 2017
Participants	537 carers of infants aged 6–8 months in Siaya County, Kenya
Interventions	TCC 1: SMS reminders for measles vaccine TCC 2: SMS reminders plus a KES 150 incentive Control: usual care

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Gibson 2019 *(Continued)*

Outcomes	1. Measles vaccination coverage; 2. Time to measles vaccination
Notes	Trial ID: NCT02904642

Harari 2018

Methods	<p>Aim: to test the acceptability and feasibility of the Lactation Advice thru Texting Can Help intervention.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: not reported</p>
Participants	58 pregnant women that intended to breastfeed recruited from 2 women, infants, and children breastfeeding peer counselling programmes in CT, USA
Interventions	<p>TCC: peer counselling support with texting (breastfeeding education and support from peer counsellors)</p> <p>Control: peer counselling support without texting</p>
Outcomes	1. Breastfeeding status; 2. Early postpartum contact; 3. Feasibility and acceptability
Notes	Trial ID: NCT02214849

Harrington 2019

Methods	<p>Aim: to assess the effect of 2-way SMS with a nurse on postpartum contraceptive use among individual women and couples.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: July 2016 to September 2017</p>
Participants	260 pregnant women (with the option to include male partners) recruited from 2 public hospitals in western Kenya
Interventions	<p>TCC: automated family planning-focused SMS messages</p> <p>Control: usual care</p>
Outcomes	Contraceptive use
Notes	Trial ID: NCT02781714

Hmone 2017

Methods	<p>Aim: to implement a breastfeeding promotion intervention using mobile phone text messages in Yangon, Myanmar, and evaluate its impact on breastfeeding practices.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: January to March 2015 (recruitment)</p>
---------	---

Hmone 2017 (Continued)

Participants	358 pregnant women of 28–34 weeks' gestation recruited from the Central Women's Hospital in Yangon, Myanmar
Interventions	TCC: breastfeeding promotional SMS messages 3 times a week Control: maternal and child healthcare messages (excluding breastfeeding-related messages) once a week
Outcomes	1. Breastfeeding outcomes; 2. Other infant feeding practices; 3. Client satisfaction
Notes	Trial ID: ACTRN12615000063516

Jasemzadeh 2018

Methods	Aim: to investigate the impact of extended parallel process model (EPPM)-based SMS on protective behaviours of pregnant women in reducing diseases caused by air pollution. Study design: randomised controlled trial Study dates: May 2015 to March 2016
Participants	130 pregnant women in Ahvaz, Iran
Interventions	TCC: SMS intervention Control: usual care
Outcomes	1. Perceived severity of air pollution; 2. Response efficacy; 3. Self-efficacy; 4. Protective behaviours against air pollution
Notes	Trial ID: IRCT2016102810804N8

Oliveira-Ciabati 2017

Methods	Aim: to determine whether PRENACEL (a bi-directional, mobile-phone based, SMS) increases the coverage of recommended antenatal care practices. Study design: cluster-randomised controlled trial Study dates: April to June 2015
Participants	1210 pregnant women invited through leaflets and posters in Brazil
Interventions	TCC: text messages with health education and health promotion content related to pregnancy and childbirth Control: usual care
Outcomes	Antenatal care outcomes
Notes	Trial ID: RBR-54zf73

Ortiz 2018

Methods	<p>Aim: to explore the safety and feasibility of using text messages in Colombia as an alternative to in-person follow-up after medication abortion for women with no clinical indication for an in-person visit.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: 2014</p>
Participants	Women after medication abortion recruited from a medical health centre in Bogota, Colombia (number of participants not reported in this conference abstract)
Interventions	<p>TCC: SMS over 14 days containing clinical information and supportive messaging</p> <p>Control: usual care</p>
Outcomes	1. Abortion complications; 2. Client satisfaction
Notes	Trial ID: not reported

Palacios 2018

Methods	<p>Aim: to test the effects of weekly SMS for improving infant feeding practices and infant weight.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: not reported</p>
Participants	202 carers of healthy term infants aged 0–2 months participating in the women, infants and children programme in Puerto Rico and Hawaii, USA
Interventions	<p>TCC: text messages on breastfeeding, preventing overfeeding, delaying introduction of solid foods, and delaying and reducing baby juice consumption</p> <p>Control: text messages related to general infant's health issues related to sleeping, bathing, teething, travelling in a car, medications, handling baby, smoking, information related to immunisation, and care of common illnesses</p>
Outcomes	1. Infant feeding practices; 2. Infant weight status
Notes	Trial ID: NCT02903186

Patel 2018

Methods	<p>Aim: to assess effectiveness of mobile phones for personalised lactation consultation to improve breastfeeding practices.</p> <p>Study design: cluster-randomised controlled trial</p> <p>Study dates: August 2010 to June 2012</p>
Participants	1036 pregnant women residing in urban slums recruited from 4 urban, public, maternity hospitals in Nagpur, India
Interventions	TCC: weekly mobile phone counselling and daily text messages

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Patel 2018 (Continued)

	Control: usual care
Outcomes	1. Breastfeeding outcomes; 2. Other infant feeding practices; 3. Client satisfaction; 4. Cost effectiveness
Notes	Trial ID: CTRI/2011/06/001822

Seth 2018

Methods	<p>Aim: to evaluate the role of compliance-linked incentives vs mobile phone messaging to improve childhood immunisations.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: July 2016 to July 2017</p>
Participants	608 carers of children aged ≤ 24 months and pregnant women from a rural community in Mewat region, Haryana state, India
Interventions	<p>TCC: automated text and voice reminders</p> <p>TCC 2: automated text and voice reminders + incentives (mobile-phone minutes)</p> <p>Control: usual care</p>
Outcomes	1. Immunisation coverage; 2. Timeliness of immunisations
Notes	Trial ID: NCT03180138

Shinde 2018

Methods	<p>Aim: to evaluate the effectiveness of the vaccine reminder system among nursing mothers.</p> <p>Study design: randomised controlled trial</p> <p>Study dates: December 2016 and March 2017 (recruitment)</p>
Participants	125 nursing mothers of newborn babies aged 0–3 weeks recruited from maternity wards of a tertiary care teaching hospital in South Canara district, Karnataka state, India
Interventions	<p>TCC: SMS vaccination reminders</p> <p>Control: usual care</p>
Outcomes	1. Vaccination coverage; 2. Client satisfaction
Notes	Trial ID: not reported

Shorey 2017

Methods	Aim: to examine the effectiveness of Home-but not Alone, a postnatal psychoeducational programme delivered via a mobile-health application for parents during the early postpartum period to improve parenting outcomes.
---------	--

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Shorey 2017 *(Continued)*

	Study design: randomised controlled trial
	Study dates: December 2015 to May 2016
Participants	250 parents recruited from a local tertiary hospital in Singapore
Interventions	TCC: mobile-health application Control: usual care
Outcomes	1. Parental self-efficacy; 2. Social support; 3. Postnatal depression; 4. Parenting satisfaction
Notes	Trial ID: ISRCTN99092313

Unger 2018

Methods	Aim: to assess the effect of SMS communication on facility delivery, exclusive breastfeeding, and contraceptive use. Study design: randomised controlled trial Study dates: August 2013 to April 2014 (recruitment)
Participants	300 pregnant women attending antenatal care recruited from a public sector maternal child health clinic in Nairobi, Kenya
Interventions	TCC 1: 1-way SMS TCC 2: 2-way SMS Control: usual care
Outcomes	1. Facility delivery; 2. Exclusive breastfeeding; 3. Contraceptive use
Notes	Trial ID: NCT01894126

Wang 2018

Methods	Aim: to evaluate the effectiveness of daily text messages as a means to improve carers' adherence to infant micronutrient powder in rural Shaanxi Province of China. Study design: cluster-randomised controlled trial Study dates: April to July 2013
Participants	638 carers of infants aged 6–11 months in randomly selected villages in rural Shaanxi Province, China
Interventions	TCC: daily text messages + free micronutrient powder packets Control: usual care + free micronutrient powder packets
Outcomes	1. Adherence to infant micronutrient powder
Notes	Trial ID: ISRCTN44149146

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Xie 2018

Methods	<p>Aim: to evaluate the potential benefits of implementing the World Health Organization maternal education programme using text messaging in a remote area in China.</p> <p>Study design: cluster-randomised controlled trial</p> <p>Study dates: October 2011 to August 2012 (recruitment)</p>
Participants	13,937 pregnant women aged 16–45 years who were registered by local Maternal Child Health unit during the study period in Hunan, China
Interventions	<p>TCC: mobile phone text messages containing maternal and newborn healthcare education</p> <p>Control: usual care</p>
Outcomes	Neonatal and maternal mortality and morbidity
Notes	Trial ID: NCT01775150

SMS: short message service; TCC: targeted client communication.

Characteristics of ongoing studies [ordered by study ID]

ACTRN12618000636257

Study name	Investigating the impact of short automated text message reminder system in improving influenza vaccine uptake in children with chronic lung conditions: a randomised controlled trial.
Methods	Randomised controlled trial
Participants	Parents of children aged 6 months to 18 years with chronic lung diseases attending the Sydney Children's Hospital, Australia
Interventions	<p>TCC: SMS text reminder + education flyer</p> <p>Control: usual care</p>
Outcomes	1. Vaccination coverage; 2. Parental attitudes towards vaccination; 3. Client satisfaction
Starting date	April 2018
Contact information	Dr Nusrat Homaira; n.homaira@unsw.edu.au
Notes	Trial ID: ACTNR12618000636257

Adam 2019

Study name	The Philani MOVIE study: a cluster-randomised controlled trial of a mobile video entertainment-education intervention to promote exclusive breastfeeding in South Africa.
Methods	Cluster-randomised controlled trial
Participants	Pregnant women within the under-resourced settlements of the Western Cape Province in South Africa

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Adam 2019 *(Continued)*

Interventions	TCC: 13 short (2–5 minutes), educational videos shared via smartphones Control: usual care
Outcomes	1. Exclusive breastfeeding (1 and 5 months); 2. Other breastfeeding and infant feeding outcomes
Starting date	November 2018
Contact information	Maya Adam; madam@stanford.edu and Charles Prober; cprober@stanford.edu
Notes	Trial ID: NCT03688217

CTRI/2018/04/013510

Study name	Effectiveness of mHealth for improving fetal outcome: a community based intervention trial.
Methods	Cluster-randomised controlled trial
Participants	Pregnant women gestation > 16 weeks in urban slums of Jaipur, India
Interventions	TCC: daily audio calls via interactive voice response system Control: usual care
Outcomes	1. Birth weight; 2. Post-term/preterm; 3. Healthcare service utilisation
Starting date	February 2016
Contact information	Manisha Malik (drmanishamalik@gmail.com); Vaseem Naheed Baig (drvaseemnaheed@yahoo.com)
Notes	Trial ID: CTRI/2018/04/013510

Cyan 2016

Study name	Affordable technology for saving maternal and infant lives
Methods	Cluster-randomised controlled trial
Participants	Pregnant women enrolled during the first trimester of pregnancy in villages in Pakistan
Interventions	Trial has 5 arms to test the effectiveness of higher vs lower frequency of messages, messages timed with progression of pregnancy and messages linked with small financial incentives. A concurrent intervention provides health literacy support to LHWs.
Outcomes	1. Adoption of intrapartum care; 2. Health literacy scores; 3. Health outcomes of mother and infant
Starting date	Not reported
Contact information	R Cyan, Georgia State University, Atlanta
Notes	Trial ID: not reported

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Gelano 2018

Study name	Effect of mobile-health on maternal health care service utilization in Eastern Ethiopia: study protocol for a randomised controlled trial.
Methods	Cluster-randomised controlled trial
Participants	Pregnant women in Haramaya and Kombolcha districts, Oromia Regional State, Eastern Ethiopia
Interventions	TCC: voice messages Control: usual care
Outcomes	1. Antenatal care visits; 2. Institutional delivery; 3. Postnatal care visits; 4. Pregnancy outcomes
Starting date	Not reported
Contact information	Tilayie Feto, gelanotilaye@gmail.com
Notes	Trial ID: PACTR201704002216259

Gul 2019

Study name	An mHealth trial to promote the use of postpartum contraception (PPFP).
Methods	Randomised controlled trial
Participants	Married, pregnant women with gestational age up to 20 weeks
Interventions	TCC 1: text and voice messages regarding antenatal and postnatal care and family planning services TCC 2: interactive phone calls regarding antenatal and postnatal care and family planning services Control: usual care
Outcomes	1. Postpartum contraceptive uptake; 2. Skilled birth; 3. Immunisation; 4. Intention to adopt modern contraception
Starting date	September 2018
Contact information	Ishaque Sheikh (ishaque.sheikh@mariestopes.org.pk); Junaid-ur-Rehman Siddiqui (junaidrehman1994@hotmail.com)
Notes	Trial ID: NCT03612518

IRCT20180520039728N1

Study name	Effectiveness of the distance education program on the mothers' empowerment in breast-feeding.
Methods	Randomised controlled trial

IRCT20180520039728N1 (Continued)

Participants	72 pregnant women referred to midwifery clinics of educational hospitals affiliated to Babol University of Medical Sciences, Iran
Interventions	TCC: text messages Control: standard care
Outcomes	1. Mothers' breastfeeding empowerment; 2. Exclusive breastfeeding
Starting date	June 2018
Contact information	Fatemeh Bakouei, f.bakouei@mubabol.ac.ir
Notes	Trial ID: IRCT20180520039728N1

ISRCTN13224744

Study name	The effectiveness of theory based intervention using social media to reduce urinary incontinence among postpartum women in Hebron city hospitals.
Methods	Randomised controlled trial
Participants	Women with urinary incontinence after child delivery in Hebron City, Palestinian Territory
Interventions	TCC: WhatsApp messages Control: usual care
Outcomes	1. Severity of urinary incontinence; 2. Level of practice of pelvic floor muscle exercise
Starting date	August 2018
Contact information	zinat.mesk@gmail.com
Notes	Trial ID: ISRCTN13224744

ISRCTN15017499

Study name	A trial exploring the feasibility of using telephone support (SMS and call) as a means of supporting young mothers (12–19 years) in Western Kenya soon after giving birth.
Methods	Randomised controlled trial
Participants	New mothers aged 12–19 and qualified midwives at the study hospitals in western Kenya
Interventions	TCC: weekly text messages and telephone calls every 3 weeks for 10 weeks in total Control: usual care
Outcomes	1. Feasibility and acceptability of the intervention; 2. Quality of data available; 3. Maternal social support; 4. Maternal self-esteem; 5. Mother–infant bonding; 6. Postnatal depression
Starting date	September 2015

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

ISRCTN15017499 *(Continued)*

Contact information	Mr Elijah Kirop (elproptich@gmail.com)
Notes	Trial ID: ISRCTN15017499

Kazi 2019

Study name	Mobile phone SMS messages and automated calls in improving vaccine coverage among children in Pakistan.
Methods	Cluster-randomised controlled trial
Participants	Carers of children aged < 14 days
Interventions	TCC 1: 1-way SMS messages related to routine immunisation TCC 2: 2-way SMS messages related to routine immunisation with the option to reply and receive more information through text messages TCC 3: 1-way automated calls related to routine immunisation TCC 4: 2-way automated calls related to routine immunisation with the option to reply and receive more information through phone call Control: usual care
Outcomes	1. Number of children who got vaccinated for routine immunisation scheduled at 6, 10, and 14 weeks of life; 2. Mean improvement in on-time vaccination for routine immunisation scheduled at 6, 10 and 14 weeks of life
Starting date	January 2018
Contact information	Abdul M Kazi (momin.kazi@aku.edu)
Notes	Trial ID: NCT03341195

Lefevre 2019

Study name	Impact evaluation of maternal health information messaging in India.
Methods	Randomised controlled trial
Participants	Women 5–7 months pregnant
Interventions	TCC: mobile phone health information messages Control: usual care
Outcomes	1. Exclusive breastfeeding; 2. Immediate breastfeeding
Starting date	July 2018
Contact information	Amnesty LeFevre (aelefevre@gmail.com); Aarushi Bhatnagar (aarushi.bhatnagar@btspmle.com)
Notes	Trial ID: NCT03576157

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Maycock 2015

Study name	Parent Infant Feeding Initiative (PIFI)
Methods	Randomised controlled trial
Participants	Mothers and their male partners attending antenatal classes at selected public and private hospitals with maternity departments in Perth, Western Australia
Interventions	Fathers will be randomly allocated to either the usual care control group, 1 of 2 medium intensity (MI1 and MI2) interventions, or a high intensity (HI) intervention. MI1 will include a specialised antenatal breastfeeding education session for fathers with supporting print materials. MI2 will involve the delivery of an antenatal and postnatal social support intervention delivered via a smartphone application and HI will include both the specialised antenatal class and the social support intervention.
Outcomes	1. Duration of any and exclusive breastfeeding; 2. Age of introduction of formula and complementary foods; 3. Infant feeding attitudes of both partners
Starting date	August 2015
Contact information	Jane A Scott (jane.scott@curtin.edu.au)
Notes	Trial ID: ACTRN12614000605695

NCT02371213

Study name	Social networking on mobile phone to improve maternal and neonatal outcomes (HISONET).
Methods	Randomised controlled trial
Participants	Thai women who attend antenatal care clinic and have intention to deliver at the study hospital.
Interventions	TCC: audio-video media via social networking on mobile phone to antenatal women from the first antenatal care visit 4 times every month and 4 times biweekly + usual antenatal care group-health education Control: usual care
Outcomes	1. Rates of premature birth; 2. Rate of respiratory distress syndrome; 3. Rate of stillbirth; 4. Rate of perinatal mortality
Starting date	April 2015
Contact information	Krissada Tomyabatra, Nopparatrajathanee Hospital
Notes	Trial ID: NCT02371213

NCT03023033

Study name	Supporting attendance for facility delivery and infant health (SAFI).
Methods	Randomised controlled trial

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

NCT03023033 (Continued)

Participants	HIV positive and HIV negative pregnant women in Tanzania
Interventions	TCC 1: SMS health promotion and reminder messages TCC 2: SMS health promotion and reminder messages + payment scaled to reflect typical transport costs to facility Control: usual care
Outcomes	1. Attendance for early infant diagnosis of HIV; 2. Early identification of HIV exposed infants at reproductive and child health clinic; 3. Antenatal care visits; 4. Facility delivery; 5. Postnatal care visits; 6. Nevirapine at delivery
Starting date	October 2014
Contact information	Godfrey Woelk, Elizabeth Glaser Pediatric AIDS Foundation
Notes	Trial ID: NCT03023033

NCT03135288

Study name	Cell-phone assisted postpartum counseling on the use of long-acting reversible contraceptives.
Methods	Randomised controlled trial
Participants	Women that gave birth at > 28 weeks' gestation and desire birth spacing for > 1 year
Interventions	TCC: mobile phone assistance regarding postpartum family planning, including reminders and 2 follow-up phone calls Control: usual care
Outcomes	Initiation of long acting reversible contraception method
Starting date	July 2017
Contact information	Dr Ahmed Mohamed Abbas, Assiut University, Egypt
Notes	Trial ID: NCT03135288

NCT03332108

Study name	Novel approach to improving lactation support with mobile health technology.
Methods	Randomised controlled trial
Participants	Women having given birth with the intention to breastfeed
Interventions	TCC: EpxBreastfeeding, a phone, and text message-based system that query patients via their personal phones and subsequently collect response data, allowing clinically relevant responses to trigger alerts to designated healthcare providers + baby book survey Control: baby book survey

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

NCT03332108 (Continued)

Outcomes	1. Exclusive breastfeeding duration; 2. Time to transition feeding status; 3. Time to reported problems with: latching, concern regarding deficiency in milk production, concern for inadequate child's weight gain; 4. Time from event to provider intervention; 5. Time to nursing status change; 6. Engagement; 7. Patient satisfaction with provider, service, and survey; 8. Breastfeeding status at 6 weeks postpartum and at 3 months postpartum; 9. Proportion of mothers exclusively breastfeeding at 6 months postpartum
Starting date	September 2017
Contact information	Camaryn Chrisman Robbins, Washington University School of Medicine, Saint Louis, MO, USA
Notes	Trial ID: NCT03332108

NCT03355989

Study name	Evaluation of conditional cash transfers (CCTs) for immunization.
Methods	Randomised controlled trial
Participants	Carers of children aged 0–23 months visiting the Expanded Programme on Immunization Center for vaccination, residing within Korangi Town in Karachi, Pakistan
Interventions	TCC: SMS reminder Control: usual care Study includes 10 more study arms of SMS reminders with various combinations of different incentives and lotteries.
Outcomes	1. Proportion of fully immunised children aged 2 years; 2. Penta3, Polio3, PCV3, or Measles1 coverage at 12–23 months; 3. Penta3, Polio3, PCV3, or Measles1 coverage at 12 months; 4. Proportion of fully immunised children at 12 months; 5. Proportion timely receipt of vaccine doses; 6. Immunisation system utilisation (dropout rate)
Starting date	November 2017
Contact information	Aamir Khan and Subhash Chandir, Interactive Research and Development
Notes	Trial ID: NCT03355989

NCT03379467

Study name	Use of SMS and interactive reminders to improve timely immunisation coverage.
Methods	Randomised controlled trial
Participants	Carers of children presenting for pentavalent 1 vaccine residing in Korangi Town in Karachi, Pakistan
Interventions	TCC 1: SMS reminder TCC 2: interactive SMS reminder Control: usual care

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

NCT03379467 (Continued)

Outcomes	1. Measles; 2. Completion
Starting date	May 2011
Contact information	Subhash Chandir, Interactive Research and Development
Notes	Trial ID: NCT03379467

NCT03399058

Study name	Innovative nutrition and mHealth Evidence Building Project.
Methods	Cluster-randomised controlled trial
Participants	Carers of underweight (weight for age z-score < -1) children aged 6–23 months in Cambodia
Interventions	TCC 1: messages on child-feeding practices TCC 2: messages on child-feeding practices with follow-up phone calls Control: usual care Co-intervention: all participants receive the basic health and nutrition service package in Cambodia called 5+5+5
Outcomes	1. Change in prevalence of underweight children over time; 2. Change in mean of weight (grams) over time; 3. Change in percentage of carers correctly answering questions on child feeding, hygiene, health-seeking, and caring practices over time; 4. Change in percentage of carers who have adopted proper child feeding, hygiene, health-seeking, and caring practices; 5. Change in percentage of carers self-reporting confidence in ability to adopt proper child feeding, hygiene, health-seeking, and caring practices over time; 6. Percentage of siblings of enrolled children in each group with a weight for height z-score < -2
Starting date	November 2017
Contact information	Oy Sreymom (sreymomoy@gmail.com); Chhea Chhorvann (cchhorvann@niph.org.kh)
Notes	Trial ID: NCT03399058

NCT03402646

Study name	Mobile phone reminders (and photovoice) for routine immunisation in Nigeria – the MOPING study.
Methods	Cluster-randomised controlled trial
Participants	Carers of healthy infants aged 0–12 months
Interventions	TCC: automated SMS and phone call reminders for routine immunisation Other: photovoice: participants will be shown photographs of debilitating consequences of non-immunisation, which will form the basis of group discussions, knowledge sharing and consensus-building sessions Control: usual care

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

NCT03402646 (Continued)

Outcomes	1. Immunisation coverage; 2. Timeliness of receipt of scheduled immunisation; 3. Incidence of any childhood vaccine-preventable disease
Starting date	May 2018
Contact information	Surajudeen A Abdulrahman (abdulsuraj@gmail.com)
Notes	Trial ID: NCT03402646

NCT03516682

Study name	Reducing delay in vaccination of children: logistic barriers (REDIVAC-LB).
Methods	Randomised controlled trial
Participants	Parents of children aged < 11 months with missing recommended vaccinations currently enrolled at Kaiser Permanente Colorado
Interventions	TCC: automated vaccination reminders (text, phone, email, or a combination) Control: usual care
Outcomes	1. Vaccines received; 2. Vaccine dose
Starting date	June 2018
Contact information	Jason Glanz, Kaiser Permanente, Denver, CO
Notes	Trial ID: NCT03516682

NCT03705455

Study name	Immunization schedule alert platform: determining ISAP SMS efficacy in improving childhood immunization timeliness and completeness in Nigeria.
Methods	Randomised controlled trial
Participants	Carer with child due for birth dose vaccination or at 6 weeks
Interventions	TCC: text messages that provides immunisation schedule and vaccine availability at the nearest immunisation clinic Control: usual care
Outcomes	1. Proportion of timely and complete vaccine uptake
Starting date	November 2018
Contact information	Emmanuel Ihedioha (emmanuel.ihedioha@lifespanhcr.com)
Notes	Trial ID: NCT03705455

NCT03797950

Study name	Mobile nudges to increase early vaccination coverage in rural areas: a pilot investigation in Ghana's northern region.
Methods	Randomised controlled trial
Participants	Women who have given birth to a live-born, surviving infant in the last 2 weeks
Interventions	TCC: voice reminder Other: cash incentive Control: usual care
Outcomes	1. Full on time early vaccination coverage (OPV0 and BCG); 2. On time BCG coverage; 3. On time OPV coverage; 4. Birth documentation and reporting coverage
Starting date	November 2018
Contact information	Guenther Fink (guenther.fink@swisstph.ch); Gillian Levine (gillian.levine@swisstph.ch)
Notes	Trial ID: NCT03797950

NCT03890978

Study name	Improving exclusive breastfeeding via mobile phone text messages: a randomized controlled trial in southern Jordan.
Methods	Randomised controlled trial
Participants	Women who had an uncomplicated singleton pregnancy and express interest in breastfeeding
Interventions	TCC: promotional exclusive breastfeeding text messages will be sent to women via mobile phone Non-TCC: child healthcare-related text messages (except breastfeeding messages) will be sent to women via mobile phone
Outcomes	1. Rate of exclusive breastfeeding; 2. Median duration of exclusive breastfeeding; 3. Rates of early initiation of breastfeeding
Starting date	January 2018
Contact information	Reham M Khresheh, Mutah University, Jordan
Notes	Trial ID: NCT03890978

Odeny 2018

Study name	Maximizing adherence and retention for women living with HIV and their infants in Kenya (MOTIVATE! study).
Methods	2 × 2 factorial cluster-randomised controlled trial

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Odeny 2018 *(Continued)*

Participants	HIV-infected pregnant women and their HIV-exposed infants attending antenatal care clinic at 1 of the study sites in Migori, Kisumu, and Homa Bay, Kenya
Interventions	TCC: mobile phone text messaging intervention TCC 2: mobile phone text messaging intervention + community mentor mother home visits TCC 3: community mentor mother home visits Control: usual care
Outcomes	1. Self-reported adherence on antiretroviral therapy; 2. Retention in care; 3. Maternal CD4 count change; 4. Uptake of intervention services; 5. Mother-to-child-transmission; 6. Male partner involvement
Starting date	May 2014
Contact information	Thomas A Odeny (taodeny@gmail.com)
Notes	Trial ID: NCT02491177

PACTR201703002093382

Study name	Effect of enhanced reminders on postnatal clinic attendance in Addis Ababa: a cluster randomized control trial.
Methods	Cluster-randomised controlled trial
Participants	Women who gave birth during the study period in public health centres in Ibadan, Ethiopia
Interventions	TCC: SMS or voice call appointment reminders Control: usual care
Outcomes	1. Number of postnatal visits; 2. Acceptability
Starting date	June 2017
Contact information	Abraham Sahilemichael Kebede (abrishya@yahoo.com)
Notes	Trial ID: PACTR201703002093382

PACTR201711002737120

Study name	Zinc adherence: a follow-up study of under-fives with acute watery diarrhoea using mobile phones: a randomised controlled trial.
Methods	Randomised controlled trial
Participants	Parents of children aged < 5 years with acute watery diarrhoea recruited from a health centre in Dar es Salaam, Tanzania
Interventions	TCC: voice calls and text messages

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

PACTR201711002737120 (Continued)

Control: usual care

Outcomes	1. 10-day adherence to zinc sulphate prescribed; 2. Clinic attendance for follow-up visit; 3. Acceptability of mHealth; 4. Dependability of mHealth
Starting date	December 2016
Contact information	Fatimah Zahra Karim (fzahrakarim@gmail.com)
Notes	Trial ID: PACTR201711002737120

PACTR201801002231314

Study name	Utilization of short message service to enhance uptake of focused ante-natal care among women in Tharaka Nithi County, Kenya.
Methods	Randomised controlled trial
Participants	Pregnant women attending rural health centres in Kenya
Interventions	TCC: SMS reminder to attend antenatal clinic Non-digital TCC: date reminder in a book to attend antenatal clinic
Outcomes	1. Enhance uptake of focused antenatal care; 2. Skilled birth attendance at birth; 3. Postnatal care at 2 weeks
Starting date	January 2016
Contact information	Eliphas Gitonga (gitonga.eliphas@ku.ac.ke)
Notes	Trial ID; PACTR201801002231314

PACTR201806003369302

Study name	Efficacy of text messages, voice calls and community visits on developmental milestones of children from teenage pregnancies: a randomized intervention trial.
Methods	Cluster-randomised controlled trial
Participants	Teenage mother, aged < 20 years with a child aged < 3 months
Interventions	TCC 1: 1 voice call weekly, 3 text messages every week TCC 2: 3 messages every week TCC 3: messages to participants' phones, 1 voice call every week and visits by a community health volunteer every month Control: usual care
Outcomes	1. Head circumference as a marker for overall brain development (6 and 12 months); 2. Diarrhoea incidence, wasting, stunting, motor and language milestones (6 and 12 months)

PACTR201806003369302 (Continued)

Starting date	April 2019
Contact information	Valerian Mwenda (valmwenda@gmail.com)
Notes	Trial ID: PACTR201806003369302

Rossing 2016

Study name	mHealth to improve measles immunization in Guinea-Bissau.
Methods	Randomised controlled trial
Participants	Mothers of children receiving measles vaccination in the Republic of Guinea-Bissau
Interventions	TCC 1: scheduled SMS reminder of the measles vaccination TCC 2: scheduled SMS reminder + voice call reminder of the measles vaccination Control: usual care
Outcomes	1. Measles vaccination coverage and timeliness when children reach 12 months of age; 2. Mean number of health centre visits (with intention to obtain the measles vaccination) required before successful administration
Starting date	March 2016
Contact information	Emil Rossing (emro@ssi.dk)
Notes	NCT02662595

Salam 2018

Study name	Saving lives with better gestational age estimation: improving the accuracy of recall and reporting of the date of last menstrual period (LMP) in Rural Bangladesh.
Methods	Cluster-randomised controlled trial
Participants	3360 adolescent girls and recently married women with no or a single child selected from the Demographic surveillance system database in Mirzapur subdistrict of Tangail district, Bangladesh
Interventions	TCC 1: counselling and a mobile phone-based SMS alert system TCC 2: counselling and smart-phone application TCC 3: counselling and a paper-based calendar Control: usual care
Outcomes	Accuracy and certainty of last menstrual period – recall dates
Starting date	January 2017
Contact information	Shumona Sharmin Salam (shumona@icddr.org)

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Salam 2018 (Continued)

Notes

Trial ID: NCT02944747

Tobe 2018

Study name	Mobile-health tool to improve maternal and neonatal health care in Bangladesh.
Methods	Cluster-randomised controlled trial
Participants	Pregnant women in Lohagora of Narail District and Dhamrai of Dhaka District, Bangladesh
Interventions	TCC: mobile short messaging and audio system Control: usual care
Outcomes	1. Neonatal mortality; 2. Maternal mortality; 3. Stillbirth; 4. Miscarriage; 5. Preterm birth; 6. Low birth weight; 7. Maternal morbidities; 8. Frequency of antenatal care visits; 9. Accessibility to skilled birth attendants for delivery; 10. Referral for identified complications; 11. Utilisation of postpartum care; 12. Status of initiating breastfeeding
Starting date	February 2017
Contact information	Ruoyan Gai Tobe (gai-r@ncchd.go.jp); Syed Emdadul Haque (emdad91@gmail.com)
Notes	Trial ID: UMIN000025628

BCG: Bacillus Calmette-Guérin; OPV: oral polio vaccine; SMS: short message service; TCC: targeted client communication.

DATA AND ANALYSES
Comparison 1. Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Health behaviour change – exclusive breastfeeding in short term (up to 3 months)	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1.1 Low-risk setting	1	40	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.79, 1.08]
1.1.2 Moderate-risk setting	1	135	Risk Ratio (M-H, Random, 95% CI)	1.30 [1.06, 1.59]
1.2 Health behaviour change – receiving postpartum help with breastfeeding (3 months postpartum)	1	332	Risk Ratio (M-H, Random, 95% CI)	2.15 [1.79, 2.58]
1.3 Health behaviour change – taking iron and folate tablets during pregnancy	1	908	Risk Ratio (M-H, Random, 95% CI)	1.71 [1.42, 2.07]
1.4 Health behaviour change – contraceptive use (3 months postpartum)	2	175	Risk Ratio (M-H, Random, 95% CI)	1.35 [0.75, 2.46]


Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

125

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.5 Health behaviour change – smoked in the last 30 days	1	459	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.17, 1.10]
1.6 Health behaviour change – smoking cessation (objectively verified continuous abstinence) (36 weeks' gestation)	1	407	Risk Ratio (M-H, Random, 95% CI)	2.76 [0.89, 8.54]
1.7 Health behaviour change – no alcohol consumption during pregnancy	1	459	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.97, 1.03]
1.8 Service utilisation – attendance at ≥ 4 antenatal care appointments	1	2550	Odds Ratio (IV, Random, 95% CI)	1.54 [0.80, 2.96]
1.9 Service utilisation – attendance for antenatal vaccination	2	714	Odds Ratio (IV, Random, 95% CI)	1.36 [0.90, 2.06]
1.9.1 Influenza vaccine	1	281	Odds Ratio (IV, Random, 95% CI)	1.22 [0.73, 2.04]
1.9.2 Tetanus vaccine	1	433	Odds Ratio (IV, Random, 95% CI)	1.67 [0.84, 3.33]
1.10 Service utilisation – attendance at antenatal preventive treatment for malaria	1	2550	Odds Ratio (IV, Random, 95% CI)	1.69 [0.82, 3.48]
1.11 Service utilisation – skilled attendant at birth	2		Odds Ratio (IV, Random, 95% CI)	Subtotals only
1.12 Service utilisation – newborn postpartum care	2	191	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.69, 1.87]
1.12.1 Low-risk setting	1	56	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.87, 1.11]
1.12.2 High-risk setting	1	135	Risk Ratio (M-H, Random, 95% CI)	1.35 [1.02, 1.78]
1.13 Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)	1	56	Risk Ratio (M-H, Random, 95% CI)	1.50 [0.30, 7.52]
1.14 Service utilisation – attendance for newborn vaccination	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.14.1 Pentavalent and polio vaccine at up to 9 weeks	1	40	Risk Ratio (M-H, Random, 95% CI)	1.08 [0.89, 1.32]
1.15 Health status and well-being – maternal mortality and morbidity	3		Risk Ratio (IV, Random, 95% CI)	Subtotals only
1.15.1 Maternal mortality up to 6 weeks postpartum	1	2637	Risk Ratio (IV, Random, 95% CI)	2.86 [0.30, 27.40]
1.15.2 Maternal morbidity – severe obstetric complications up to 6 weeks postpartum	1	2550	Risk Ratio (IV, Random, 95% CI)	0.86 [0.70, 1.07]
1.15.3 Maternal morbidity – any maternal health problem up to 10 days postpartum	1	56	Risk Ratio (IV, Random, 95% CI)	0.50 [0.09, 2.76]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.15.4 Maternal morbidity – mastitis: breast pain up to 3 months postpartum	1	332	Risk Ratio (IV, Random, 95% CI)	0.28 [0.09, 0.80]
1.15.5 Maternal morbidity – mastitis: breast engorgement up to 3 months postpartum	1	332	Risk Ratio (IV, Random, 95% CI)	0.58 [0.31, 1.10]
1.16 Health status and well-being – maternal mortality and morbidity	1	135	Mean Difference (IV, Random, 95% CI)	0.06 [-0.19, 0.31]
1.16.1 Acute episodes requiring clinic visit up to 3 months postpartum	1	135	Mean Difference (IV, Random, 95% CI)	0.06 [-0.19, 0.31]
1.17 Health status and well-being – neonatal mortality and morbidity	3	2870	Odds Ratio (IV, Random, 95% CI)	1.00 [0.61, 1.64]
1.17.1 Neonatal mortality up to 6 weeks after delivery	1	2482	Odds Ratio (IV, Random, 95% CI)	0.85 [0.37, 1.95]
1.17.2 Neonatal diarrhoea up to 3 months postpartum	1	332	Odds Ratio (IV, Random, 95% CI)	1.05 [0.53, 2.11]
1.17.3 Any newborn health problem reported up to 10 days postpartum	1	56	Odds Ratio (IV, Random, 95% CI)	1.27 [0.36, 4.51]
1.18 Health status and well-being – neonatal mortality and morbidity	1		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.18.1 Acute episodes requiring clinic visit up to 3 months postpartum	1	135	Mean Difference (IV, Random, 95% CI)	-0.53 [-0.92, -0.14]
1.19 Health status and well-being – neonatal health	2		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.19.1 Gestational age at birth (weeks)	1	61	Mean Difference (IV, Random, 95% CI)	0.10 [-0.45, 0.65]
1.19.2 Birth weight (g)	1	61	Mean Difference (IV, Random, 95% CI)	-173.00 [-448.87, 102.87]
1.19.3 Infant weight (kg) at 3 months	1	332	Mean Difference (IV, Random, 95% CI)	Not estimable
1.19.4 Infant length (cm) at 3 months	1	332	Mean Difference (IV, Random, 95% CI)	Not estimable
1.20 Health status and well-being – preterm birth	2	2557	Risk Ratio (IV, Random, 95% CI)	0.85 [0.31, 2.33]
1.21 Sensitivity analysis (cluster-RCTs: health status and well-being – preterm birth)	1		Risk Ratio (IV, Random, 95% CI)	0.18 [0.01, 3.64]


Analysis 1.1. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 1: Health behaviour change – exclusive breastfeeding in short term (up to 3 months)




Footnotes

- (1) 9 weeks postpartum
- (2) 3 months postpartum

Analysis 1.2. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 2: Health behaviour change – receiving postpartum help with breastfeeding (3 months postpartum)




Analysis 1.3. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 3: Health behaviour change – taking iron and folate tablets during pregnancy



Footnotes

- (1) cluster RCT adjusted using design effect=1.92, calculated with ICC=0.154 reported in Pagel 2011; original sample size=1743


Analysis 1.4. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 4: Health behaviour change – contraceptive use (3 months postpartum)



Footnotes

- (1) 3 months postpartum
- (2) 9 weeks post-partum


Analysis 1.5. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 5: Health behaviour change – smoked in the last 30 days




Footnotes

- (1) 4 weeks after starting intervention in pregnant and postpartum population

Analysis 1.6. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 6: Health behaviour change – smoking cessation (objectively verified continuous abstinence) (36 weeks' gestation)




Analysis 1.7. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 7: Health behaviour change – no alcohol consumption during pregnancy



Footnotes

(1) 4 weeks after starting intervention in pregnant and postpartum population


Analysis 1.8. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 8: Service utilisation – attendance at ≥ 4 antenatal care appointments



Footnotes

(1) cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)


Analysis 1.9. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 9: Service utilisation – attendance for antenatal vaccination



Footnotes

(1) Nullipara subgroup only; cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)


Analysis 1.10. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 10: Service utilisation – attendance at antenatal preventive treatment for malaria



Footnotes

(1) cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)


Analysis 1.11. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 11: Service utilisation – skilled attendant at birth




Footnotes

- (1) cluster RCT adjusted using design effect=1.24, calculated from ICC 0.041 reported in Pagel 2011
- (2) rural population; cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)
- (3) urban population; cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)


Analysis 1.12. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 12: Service utilisation – newborn postpartum care




Analysis 1.13. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 13: Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)



Analysis 1.14. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 14: Service utilisation – attendance for newborn vaccination




Analysis 1.15. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 15: Health status and well-being – maternal mortality and morbidity




Footnotes

(1) cluster RCT adjusted using design effect=1.33, calculated with ICC=0.003 reported in Pagel 2011

Analysis 1.16. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 16: Health status and well-being – maternal mortality and morbidity




Analysis 1.17. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 17: Health status and well-being – neonatal mortality and morbidity




Footnotes

(1) cluster RCT with study-adjusted estimate (adjusted for within-cluster effect)


Analysis 1.18. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 18: Health status and well-being – neonatal mortality and morbidity



Analysis 1.19. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 19: Health status and well-being – neonatal health




Analysis 1.20. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 20: Health status and well-being – preterm birth



Footnotes

(1) cluster RCT adjusted using design effect=1.33, calculated with ICC=0.003 reported in Pagel 2011


Analysis 1.21. Comparison 1: Digital targeted client communication (TCC) compared to standard care (pregnant and postpartum women), Outcome 21: Sensitivity analysis (cluster-RCTs: health status and well-being – preterm birth)




Comparison 2. Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Health behaviour change – exclusive breastfeeding (9 weeks postpartum)	1	42	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.79, 1.07]
2.2 Health behaviour change – contraceptive use (9 weeks postpartum)	1	42	Risk Ratio (M-H, Random, 95% CI)	1.45 [0.78, 2.69]
2.3 Service utilisation – newborn postpartum care (10 days after delivery)	1	59	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.09]
2.4 Service utilisation – attendance for newborn vaccination	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.4.1 Pentavalent and polio vaccine at 9 weeks	1	42	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.88, 1.16]
2.5 Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)	1	59	Risk Ratio (M-H, Random, 95% CI)	0.56 [0.18, 1.79]
2.6 Health status and well-being – maternal mortality and morbidity	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.6.1 Any maternal health problem reported up to 10 days postpartum	1	59	Risk Ratio (M-H, Random, 95% CI)	0.19 [0.04, 0.79]
2.7 Health status and well-being – neonatal mortality and morbidity	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.7.1 Any newborn health problem reported up to 10 days postpartum	1	59	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.25, 1.06]


Analysis 2.1. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 1: Health behaviour change – exclusive breastfeeding (9 weeks postpartum)




Analysis 2.2. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 2: Health behaviour change – contraceptive use (9 weeks postpartum)




Analysis 2.3. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 3: Service utilisation – newborn postpartum care (10 days after delivery)




Analysis 2.4. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 4: Service utilisation – attendance for newborn vaccination




Analysis 2.5. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 5: Service utilisation – attendance for postpartum care appointment (mother) (10 days postpartum)



Analysis 2.6. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 6: Health status and well-being – maternal mortality and morbidity




Analysis 2.7. Comparison 2: Digital targeted client communication (TCC) compared to non-digital targeted client communication (pregnant and postpartum women), Outcome 7: Health status and well-being – neonatal mortality and morbidity




Comparison 3. Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant and postpartum women)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.1 Service utilisation – attendance for antenatal influenza vaccination	1	204	Risk Ratio (M-H, Random, 95% CI)	1.05 [0.71, 1.58]
3.2 Service utilisation – birth at health facility	1	16	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.69, 1.45]

Analysis 3.1. Comparison 3: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant and postpartum women), Outcome 1: Service utilisation – attendance for antenatal influenza vaccination



Analysis 3.2. Comparison 3: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant and postpartum women), Outcome 2: Service utilisation – birth at health facility



Footnotes


(1) cluster RCT adjusted using design effect=17, calculated with ICC=0.127 reported in Pagel 2011; original sample size=274

Comparison 4. Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4.1 Health behaviour change – mother taking any type of anti-retroviral (ARV) (34–36 weeks' gestation)	1	503	Risk Ratio (IV, Random, 95% CI)	1.04 [0.91, 1.19]
4.2 Health behaviour change – mother taking any type of ARV (6–8 weeks postpartum)	1	471	Risk Ratio (IV, Random, 95% CI)	0.87 [0.61, 1.24]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4.3 Health behaviour change – infant ARV/prevention of mother-to-child transmission treatment adherence (6 weeks postpartum)	1	223	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.98, 1.04]
4.4 Health behaviour change – infant HIV tested (6–8 weeks postpartum)	2	838	Risk Ratio (IV, Random, 95% CI)	1.04 [0.95, 1.13]
4.5 Service utilisation – postnatal care: attendance at postpartum care appointment (6–8 weeks postpartum)	1	381	Risk Ratio (M-H, Random, 95% CI)	1.66 [1.02, 2.70]
4.6 Service utilisation – intrapartum care: birth in health facility	1	134	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.62, 1.15]
4.7 Service utilisation – antenatal care: mean number of face-to-face or mobile communications with healthcare workers	1	297	Mean Difference (IV, Random, 95% CI)	1.50 [-0.36, 3.36]
4.8 Health status and well-being – neonatal health: neonatal death/stillbirth	1	381	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.39, 3.28]
4.9 Health status and well-being – neonatal health: infant HIV test positive (6–8 weeks postpartum)	2	852	Risk Ratio (IV, Random, 95% CI)	0.54 [0.11, 2.56]


Analysis 4.1. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 1: Health behaviour change – mother taking any type of antiretroviral (ARV) (34–36 weeks' gestation)



Footnotes

(1) cluster RCT with study-adjusted RR (adjusted for cluster residuals and confounding variables including participant age, gestational age, whether the woman was newly diagnosed


Analysis 4.2. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 2: Health behaviour change – mother taking any type of ARV (6–8 weeks postpartum)



Footnotes

(1) cluster RCT with study-adjusted RR (adjusted for cluster residuals and confounding variables including participant age, gestational age, whether the woman was newly diagnosed)


Analysis 4.3. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 3: Health behaviour change – infant ARV/ prevention of mother-to-child transmission treatment adherence (6 weeks postpartum)



Footnotes

(1) cluster RCT adjusted using design effect=2.1085, calculated with ICC=0.055 reported in Pagel 2011; original sample size=470


Analysis 4.4. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 4: Health behaviour change – infant HIV tested (6–8 weeks postpartum)




Footnotes

(1) cluster RCT adjusted using design effect=2.1085, calculated with ICC=0.055 reported in Pagel 2011

Analysis 4.5. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 5: Service utilisation – postnatal care: attendance at postpartum care appointment (6–8 weeks postpartum)




Analysis 4.6. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 6: Service utilisation – intrapartum care: birth in health facility



Footnotes

(1) cluster RCT adjusted using design effect=3.560, calculated with ICC=0.127 reported in Pagel 2011; original sample size=479


Analysis 4.7. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 7: Service utilisation – antenatal care: mean number of face-to-face or mobile communications with healthcare workers




Footnotes

(1) cluster RCT adjusted using design effect=1.605, calculated with ICC=0.030 reported in Pagel 2011; original sample size=476

Analysis 4.8. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 8: Health status and well-being – neonatal health: neonatal death/stillbirth



Analysis 4.9. Comparison 4: Digital targeted client communication (TCC) compared to standard care (pregnant women with HIV), Outcome 9: Health status and well-being – neonatal health: infant HIV test positive (6–8 weeks postpartum)




Footnotes

(1) cluster RCT adjusted using design effect=2.1085, calculated with ICC=0.055 reported in Pagel 2011


Comparison 5. Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant women with HIV)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.1 Health behaviour – infant antiretroviral/prevention of mother-to-child transmission adherence (6 weeks after delivery)	1	150	Risk Ratio (M-H, Random, 95% CI)	1.26 [1.07, 1.48]
5.2 Service utilisation – postnatal care: attendance at postpartum care appointment (10 weeks postpartum)	1	150	Risk Ratio (M-H, Random, 95% CI)	1.86 [1.34, 2.58]

Analysis 5.1. Comparison 5: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant women with HIV), Outcome 1: Health behaviour – infant antiretroviral/prevention of mother-to-child transmission adherence (6 weeks after delivery)




Analysis 5.2. Comparison 5: Digital targeted client communication (TCC) compared to digital non-targeted client communication (pregnant women with HIV), Outcome 2: Service utilisation – postnatal care: attendance at postpartum care appointment (10 weeks postpartum)



Comparison 6. Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
6.1 Attendance for necessary healthcare	10	5660	Risk Ratio (IV, Random, 95% CI)	1.21 [1.08, 1.34]
6.1.1 Vaccinations at up to 6 months	5	1586	Risk Ratio (IV, Random, 95% CI)	1.14 [1.01, 1.28]
6.1.2 Vaccinations at 12 months	4	3832	Risk Ratio (IV, Random, 95% CI)	1.24 [1.02, 1.52]
6.1.3 HIV medical appointment 2 days after reminder	1	242	Risk Ratio (IV, Random, 95% CI)	1.63 [1.26, 2.11]
6.2 Timeliness of vaccination	4	2400	Risk Ratio (IV, Random, 95% CI)	1.18 [1.04, 1.34]
6.3 Service utilisation – no emergency department attendance (6 months)	1	129	Risk Ratio (M-H, Random, 95% CI)	1.32 [1.03, 1.70]


Analysis 6.1. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 1: Attendance for necessary healthcare



Footnotes

- (1) cluster RCT adjusted using design effect=6.9901, calculated with ICC=0.0487 reported in Gibson 2017
- (2) cluster RCT adjusted using design effect=8.37805, calculated with ICC=0.0487 reported in Gibson 2017
- (3) cluster RCT with study-adjusted RR accounting for correlation within clusters
- (4) Combined three intervention groups (text messages, voice call, or text messages + voice call)

Analysis 6.2. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 2: Timeliness of vaccination



Footnotes

- (1) cluster RCT with study-adjusted RR accounting for correlation within clusters
- (2) Combined two intervention groups (educational and conventional SMS)

Analysis 6.3. Comparison 6: Digital targeted client communication (TCC) compared to standard care (parents of children aged under five years), Outcome 3: Service utilisation – no emergency department attendance (6 months)

Study or Subgroup	TCC via mobile devices		Standard care		Weight	Risk Ratio	
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI
Hannan 2016	48	63	38	66	100.0%	1.32 [1.03, 1.70]	
Total (95% CI)		63		66	100.0%	1.32 [1.03, 1.70]	
Total events:		48	38				
Heterogeneity: Not applicable							
Test for overall effect: Z = 2.21 (P = 0.03)							
Test for subgroup differences: Not applicable							

Comparison 7. Digital targeted client communication (TCC) compared to non-digital targeted client communication (parents of children aged under five years)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.1 Health behaviour change – oral health in children (Visible Plaque Index, [0–100%], low = good)	1	143	Mean Difference (IV, Random, 95% CI)	-2.10 [-7.54, 3.34]
7.2 Service utilisation – attendance for vaccinations at 14 weeks	1	744	Risk Ratio (IV, Random, 95% CI)	1.13 [1.00, 1.28]

Analysis 7.1. Comparison 7: Digital targeted client communication (TCC) compared to non-digital targeted client communication (parents of children aged under five years), Outcome 1: Health behaviour change – oral health in children (Visible Plaque Index, [0–100%], low = good)

Study or Subgroup	TCC via mobile devices			Non-digital TCC			Weight	Mean Difference	
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	IV, Random, 95% CI
Sharma 2011 (1)	33.5	17	71	35.6	16.2	72	100.0%	-2.10 [-7.54, 3.34]	
Total (95% CI)			71			72	100.0%	-2.10 [-7.54, 3.34]	
Heterogeneity: Not applicable									
Test for overall effect: Z = 0.76 (P = 0.45)									
Test for subgroup differences: Not applicable									

Footnotes

(1) 4 weeks after intervention

Analysis 7.2. Comparison 7: Digital targeted client communication (TCC) compared to non-digital targeted client communication (parents of children aged under five years), Outcome 2: Service utilisation – attendance for vaccinations at 14 weeks

Study or Subgroup	log[RR]	SE	TCC via mobile devices		Non-digital TCC		Weight	Risk Ratio	Risk Ratio
			Total	Total	IV, Random, 95% CI	IV, Random, 95% CI			
Haji 2016 (1)	0.1252	0.064	372	372	100.0%	1.13 [1.00, 1.28]			
Total (95% CI)			372	372	100.0%	1.13 [1.00, 1.28]			

Heterogeneity: Not applicable
Test for overall effect: Z = 1.96 (P = 0.05)
Test for subgroup differences: Not applicable

Footnotes

(1) cluster RCT adjusted using design effect=6.9901, calculated with ICC=0.0487 reported in Gibson 2017

Comparison 8. Digital targeted client communication (TCC) compared to digital non-targeted client communication (parents of children aged under five years)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
8.1 Service utilisation – attendance for vaccinations at 7 months	1	40	Risk Ratio (M-H, Random, 95% CI)	0.63 [0.33, 1.20]

Analysis 8.1. Comparison 8: Digital targeted client communication (TCC) compared to digital non-targeted client communication (parents of children aged under five years), Outcome 1: Service utilisation – attendance for vaccinations at 7 months

Study or Subgroup	TCC via mobile device		Non-TCC via mobile device		Weight	Risk Ratio	Risk Ratio
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI
Niederhauser 2015	7	17	15	23	100.0%	0.63 [0.33, 1.20]	
Total (95% CI)		17	15	23	100.0%	0.63 [0.33, 1.20]	

Total events: 7
Heterogeneity: Not applicable
Test for overall effect: Z = 1.40 (P = 0.16)
Test for subgroup differences: Not applicable

ADDITIONAL TABLES

Table 1. Setting and income group of included studies

Study ID	Setting	Country	Income group ^a
Population: pregnant and post-partum women			
Evans 2014	Community	USA	High income
Jareethum 2008	Healthcare	Thailand	Upper middle income
Joshi 2015	Community	India	Lower middle income

Table 1. Setting and income group of included studies (Continued)

Kamau-Mbuthia 2013	Healthcare	Kenya	Lower middle income
Lund 2012	Healthcare	Tanzania	Low income
Maslowsky 2016	Healthcare and community	Ecuador	Upper middle income
McConnell 2016	Healthcare	Kenya	Lower middle income
Moniz 2013	Community	USA	High income
Naughton 2017	Healthcare	England	High income
Omole 2018	Healthcare	Nigeria	Lower middle income
Yudin 2017	Healthcare	Canada	High income
Population: pregnant and post-partum women living with HIV			
Kassaye 2016	Healthcare and community	Kenya	Lower middle income
Kebaya 2014	Community	Kenya	Lower middle income
Odeny 2014	Healthcare	Kenya	Lower middle income
Population: parents of children under five years			
Ahlers-Schmidt 2012	Healthcare	USA	High income
Bangure 2015	Healthcare	Zimbabwe	Low income
Bigna 2015	Healthcare	Cameroon	Lower middle income
Brown 2016	Healthcare and community	Nigeria	Lower middle income
Domek 2016	Healthcare and community	Guatemala	Lower middle income
Eze 2015	Healthcare	Nigeria	Lower middle income
Gibson 2017	Community	Kenya	Lower middle income
Haji 2016	Community	Kenya	Lower middle income
Hannan 2016	Community	USA	High income
Hofstetter 2015a	Community	USA	High income
Niederhauser 2015	Healthcare	Hawaii	High income
Sharma 2011	Not reported	India	Lower middle income
Stockwell 2015	Healthcare and community	USA	High income

^aIncome group according to World Bank list of economies, June 2017 (iccmoot.com/wp-content/uploads/2017/07/World-Bank-List-of-Economies.pdf).

Table 2. Interventions in included studies

Study ID	Intervention type		Theory	Phone compat- ibility	Delivery mechanism	Personalisation	Data security	Assessment of fidelity
	Remind/re- call	Inform/ed- ucate or support						
Population: pregnant and post-partum women								
Evans 2014	—	Inform/edu- cate or sup- port	Based on: – health belief model – social in- fluence and dif- fusion of infor- mation within a target popula- tion – so- cial cog- nitive theory.	Not reported	Text mes- sages	Messages tailored to date of enrolment and baby's gestational age. Thus, a woman enrolling in her 10 week of pregnancy would begin receiving week 10 messages.	Quote: "Voxi- va [commer- cial company providing tex- ting services] did not have access to any data collected ... or any pa- tient infor- mation stored at Madigan."	Not reported
Jareethum 2008	—	Inform/edu- cate or sup- port	Not men- tioned	Not reported	Text mes- sages; phone call at 32 weeks' gestation	Not reported	Not reported	Not reported
Joshi 2015	—	Inform/edu- cate or sup- port	Not re- ported	Simple mobile phone (SMS and call func- tions only)	MMS, in- cluding video and audiovisual messages; voice calls	Timed and targeted as per beneficiary's gestational age; sent in a user spec- ified language and time slot.	Not reported	Not reported
Ka- mau-Mbuthia 2013	—	Inform/edu- cate or sup- port	Not re- ported	Simple mobile phone (SMS and call func- tions only)	Text mes- sages	Not reported	Not reported	Not reported

Table 2. Interventions in included studies (Continued)

Lund 2012	Remind/re-call	Inform/educate or support	Not mentioned	Simple mobile phone (SMS and call functions only)	Text messages	The content of the messages varied depending on the stage of the pregnancy. Message content provided as simple text in the local language of Swahili.	Not reported	Not reported
Maslowsky 2016	—	Inform/educate or support	Not mentioned	Simple mobile phone (SMS and call functions only); feature phone (can run java apps); smartphones (android, iOS, Symbian)	Voice calls	Each participant also indicated the days of the week and times of day that were best to reach her.	Not reported	All intervention participants completed the 48-hour post-discharge education session. Only 3 participants called the nurse to ask for additional advice.
McConnell 2016	—	Inform/educate or support	The checklist used by CHWs was developed using international guidelines and academic publications (Table 1, page 3).	Simple mobile phone (SMS and call functions only)	Voice calls	Process was available in English and Kiswahili.	Not reported	Not reported
Moniz 2013	—	Inform/educate or support	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Not reported	All text messages were sent through a password-protected online service (www.ez-texting.com).	13/104 received no messages.

Table 2. Interventions in included studies (Continued)

Naughton 2017	—	Inform/educate or support	MiQuit objectives are informed by Social Cognitive Theory Perspectives on Change Theory, the Elaboration Likelihood Model of Persuasion and several additional cognitive determinants of quitting smoking in pregnancy.	Simple mobile phone (SMS and call functions only)	Text messages, phone call, email just 4 weeks after randomisation	Tailoring characteristics of the intervention include gestation, motivation to quit, the hardest situation to avoid smoking, cessation self-efficacy, cigarette dependence and partner's smoking status. Women were encouraged to set and send a quit date to MiQuit to enable them to receive additional support orientated around when their quit attempt begins.	Not reported	Intervention fidelity was high, 98% of MiQuit recipients recalled receiving text message support.
Omole 2018	Remind/recall	Inform/educate or support	Not mentioned	Simple mobile phone (SMS and call functions only)	Text messages	Pregnancy reminders for ANC appointments and health information structured by the age of the pregnancy, delivered once a week, at a time chosen by the participant.	Not reported	<ol style="list-style-type: none"> 1. Recording of all messages sent by the system and archiving of messages received from registered participants. In addition, the timestamp when the SMS was received and the response given to each SMS received were collected. 2. Periodic contact (based on ANC appointments) with study participants to confirm receipt of messages.

3. All responses to clients' messages were stored in a database.

Table 2. Interventions in included studies (Continued)

Yudin 2017	—	Inform/educate or support	The messages were developed using principles from the Health Belief Model of preventive health behaviour.	Simple mobile phone (SMS and call functions only)	Text messages	Not reported	Messages were sent via a password-protected online service, Memo-text (Memo-text LLC).	Not reported
Population: pregnant and post-partum women living with HIV								
Kassaye 2016	Remind/recall	Inform/educate or support	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Messages targeted based on pregnancy stage.	Not reported	Not reported
Kebaya 2014	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Voice calls	Not reported	Not reported	Not reported
Odeny 2014	Remind/recall	Inform/educate or support	Health Belief Model	Simple mobile phone (SMS and call functions only)	Text messages	SMS preferences were recorded by sending a text message from their phone. This message included date of last normal menstrual period and preferred language (English, Kiswahili, or Dholuo). For participants randomised to the intervention, the message also included preferred time for receiving	Not reported	Not reported

Table 2. Interventions in included studies (Continued)

Population: parents of children under five years								
Ahlers-Schmidt 2012	Remind/recall	—	Not reported	Not reported	Text messages	Not reported	Not reported	Not reported
Bangure 2015	Remind/recall	—	Not reported	Not reported	Text messages	None	Not reported	Not reported
Bigna 2015	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Text messages, voice calls	Not reported	Text message did not contain information about the name of the child or adult, or the health status of the child. Name of adult and child not mentioned in phone call.	57/121 (47%) participants did not receive intervention.
Brown 2016	Remind/recall	Inform/educate or support	Not reported	Simple mobile phone (SMS and call functions only)	Voice calls	None	Not reported	Not reported
Domek 2016	Remind/recall	—	Not reported	Not reported	Text messages	SMS messages were autotopopulated with the child's name and the name of the appropriate clinic.	Not reported	Not reported
Eze 2015	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Each message is tagged with the name of the health facility. No other personalisation reported.	Not reported	No SMS delivery log. Authors stated that "The majority of respondents, [93.1%] accepted the option of being sent reminder messages." Among those who preferred not to be sent SMS reminders, self-confidence in not forgetting appointments (61%) and the

ing SMS, and preferred name.

Table 2. Interventions in included studies (Continued)

								<p>fear of giving out phone numbers (28.8%) were the greatest barriers to mHealth uptake in this study.</p>
Gibson 2017	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Baby's first name, relevant vaccine name, and relevant district were personalised in the messages.	Not reported	The vast majority of carers reported receiving ≥ 1 SMS reminder or incentive during the study.
Haji 2016	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Not reported	Not reported	1488 messages were sent to the participants in the SMS group.
Hannan 2016	Remind/recall	Inform/educate or support	Not reported	Simple mobile phone (SMS and call functions only); feature phone (can run java apps); smartphones (android, iOS, Symbian)	Text messages, voice calls	Not reported	Not reported	681 mobile phone calls; 630 by NPs to mothers and 51 calls to NPs by mothers. The NPs were contacted via texting 29 times by the mothers.
Hofstetter 2015a	Remind/recall	—	Not reported	Simple mobile phone (SMS and call functions only)	Text messages	Messages were sent in either English or Spanish depending on the primary language specified in the electronic health record.	This study utilised a customised text messaging platform integrated with the hospital registration system and its immunisation registry, EzVac. The registration system included demographic and visit data for participants, while the EzVac registry	Of 1254 parents in the text messaging arms who were sent ≥ 1 text message, 7.1% experienced ≥ 1 undelivered message.

Table 2. Interventions in included studies (Continued)

							automatical-ly captured from the insti-tution's elec-tronic health record all vac-cine doses ad-ministered to subjects at the hospital and affiliated clinics.	
Nieder-hauser 2015	Remind/re-call	—	Not re-ported	Simple mobile phone (SMS and call func-tions only)	Text mes-sages	The SMS included word-ing such as: "your ba-by," "your appoint-ment," "your health care provider."	Not reported	To validate that the partici-pants were receiving the text messages, 10% of the parents were randomly selected to re-ceive phone calls verifying the receiving of the messages.
Sharma 2011	—	Inform/edu-cate or sup-port	Not re-ported	Simple mobile phone (SMS and call func-tions only)	Text mes-sages	Messages targeted moth-ers by saying: "you should," "your child teeth, your child tongue," etc.	Not reported	Not reported
Stockwell 2015	Remind/re-call	Inform/edu-cate or sup-port	5 mes-sages were se-lected based on previous studies.	Simple mobile phone (SMS and call func-tions only)	Text mes-sages	SMS were sent in English or Spanish based on the participant's request at enrolment.	Not reported	Although no undeliver-able SMS were registered, 1 postintervention survey re-vealed that some families did not remember receiving SMS.

ANC: antenatal care; CHW: community healthcare worker; MMS: multimedia service; NP: nurse practitioner; SMS: short message service.

APPENDICES

Appendix 1. CENTRAL search strategy

- #1 MeSH descriptor: [Cell Phones] this term only 535
- #2 MeSH descriptor: [Smartphone] this term only 73
- #3 MeSH descriptor: [MP3-Player] this term only 19
- #4 MeSH descriptor: [Computers, Handheld] this term only 203
- #5 ((cell* or mobile*) near/1 (phone* or telephone* or technolog* or device*)):ti,ab,kw 1601
- #6 (handheld or hand-held):ti,ab,kw 1174
- #7 (smartphone* or smart-phone* or cellphone* or mobiles):ti,ab,kw 749
- #8 ((personal near/1 digital) or (PDA near/3 (device* or assistant*)) or "MP3 player*" or "MP4 player*"):ti,ab,kw 188
- #9 (samsung or nokia):ti,ab,kw 44
- #10 (windows near/3 (mobile* or phone*)):ti,ab,kw 3
- #11 android:ti,ab,kw 155
- #12 (ipad* or i-pad* or ipod* or i-pod* or iphone* or i-phone*):ti,ab,kw 403
- #13 (tablet* near/3 (device* or computer*)):ti,ab,kw 187
- #14 MeSH descriptor: [Telemedicine] this term only 1528
- #15 MeSH descriptor: [Videoconferencing] this term only 134
- #16 MeSH descriptor: [Webcasts as Topic] this term only 16
- #17 MeSH descriptor: [Text Messaging] this term only 428
- #18 MeSH descriptor: [Telenursing] this term only 25
- #19 (mhealth or m-health or "mobile health" or ehealth or e-health or "electronic health"):ti,ab,kw 1305
- #20 (telemedicine or tele-medicine or telehealth or tele-health or telecare or tele-care or telenursing or tele-nursing or telepsychiatry or tele-psychiatry or telemonitor* or tele-monitor* or teleconsult* or tele-consult* or telecounsel* or tele-counsel* or telecoach* or tele-coach*):ti,ab,kw 3320
- #21 (videoconferenc* or video-conferenc* or webcast* or web-cast*):ti,ab,kw 418
- #22 (((text* or short or voice or multimedia or multi-media or electronic or instant) near/1 messag*) or "instant messenger") .ti,ab,kw 48
- #23 (texting or texted or texter* or ((sms or mms) near (service* or messag*)) or "interactive voice response*" or IVR or "voice call*" or callback* or "voice over internet" or VOIP):ti,ab,kw 1011
- #24 (Facebook or Twitter or Whatsapp* or Skyp* or YouTube or "You Tube" or "Google Hangout*"):ti,ab,kw 226
- #25 MeSH descriptor: [Mobile Applications] this term only 151
- #26 "mobile app*":ti,ab,kw 441
- #27 MeSH descriptor: [Social Media] this term only 67
- #28 (social near (media or network*)):ti,ab,kw 967
- #29 MeSH descriptor: [Reminder Systems] this term only 816
- #30 (remind* near/3 (text* or system* or messag*)):ti,ab,kw 1228
- #31 MeSH descriptor: [Electronic Mail] this term only 278

- #32 ("electronic mail*" or email* or e-mail or webmail):ti,ab,kw 1778
- #33 MeSH descriptor: [Medical Informatics] this term only 76
- #34 MeSH descriptor: [Medical Informatics Applications] this term only 28
- #35 MeSH descriptor: [Nursing Informatics] this term only 10
- #36 MeSH descriptor: [Public Health Informatics] this term only 6
- #37 ((medical or clinical or health or healthcare or nurs*) near/3 informatics):ti,ab,kw 265
- #38 MeSH descriptor: [Multimedia] this term only 192
- #39 MeSH descriptor: [Hypermedia] this term only 8
- #40 MeSH descriptor: [Blogging] this term only 14
- #41 (multimedia or multi-media or hypermedia or hyper-media or blog* or vlog* or weblog* or web-log*):ti,ab,kw 728
- #42 MeSH descriptor: [Interactive Tutorial] this term only 0
- #43 MeSH descriptor: [Computer-Assisted Instruction] this term only 1132
- #44 ((interactive or computer-assisted) near/1 (tutor* or technolog* or learn* or instruct* or software or communication)):ti,ab,kw 1322
- #45 {or #1-#44} 13953
- #46 MeSH descriptor: [Family Planning Services] this term only 205
- #47 MeSH descriptor: [Contraception] this term only 289
- #48 MeSH descriptor: [Reproductive Behavior] this term only 11
- #49 MeSH descriptor: [Contraception Behavior] this term only 187
- #50 MeSH descriptor: [Contraceptive Agents] explode all trees 2318
- #51 MeSH descriptor: [Contraceptive Devices] explode all trees 1411
- #52 (condom* or (OC near pill) or ("depot medroxyprogest*" or NET-EN or "NET EN" or Mesigyna or Cyclofem) or ("intrauterine system" or "intra-uterine system" or IUS or "intrauterine device*" or "intra-uterine device*" or IUD*) or (vasectomy or sterilisation or sterilization or (tubal near ligation)) or ((vaginal near ring) or cyclotel or cycle-tel or ((abstain or abstinen*) near/2 (sex* or intercourse)) or "lactational amenorr*"):ti,ab,kw 4178
- #53 (contracept* or "family planning" or (birth near (control or regulat* or spacing)) or "planned parenthood" or ((population or fertility) near (regulat* or control))):ti,ab,kw 9149
- #54 MeSH descriptor: [Pregnancy in Adolescence] this term only 185
- #55 (pregnan* near/2 (adolescen* or teen* or schoolchild*)):ti,ab,kw 1541
- #56 MeSH descriptor: [Pregnancy, Unplanned] this term only 73
- #57 MeSH descriptor: [Pregnancy, Unwanted] this term only 48
- #58 (pregnan* near/3 (prevent* or interrupt* or unplanned or unwanted or mistimed)):ti,ab,kw 2178
- #59 MeSH descriptor: [Sexually Transmitted Diseases] explode all trees and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Epidemiology - EP, Prevention & control - PC, Psychology - PX, Transmission - TM] 8731
- #60 ("sexually transmi*" or STI or STIs or STD or STDs or venereal):ti,ab,kw 2535
- #61 MeSH descriptor: [HIV Infections] explode all trees and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Epidemiology - EP, Prevention & control - PC, Psychology - PX, Transmission - TM] 7638
- #62 MeSH descriptor: [HIV Seropositivity] this term only and with qualifier(s): [Drug therapy - DT, Epidemiology - EP, Psychology - PX, Transmission - TM] 380

- #63 MeSH descriptor: [Anti-HIV Agents] this term only 2641
- #64 MeSH descriptor: [Antiretroviral Therapy, Highly Active] this term only 1246
- #65 MeSH descriptor: [Medication Adherence] this term only 1733
- #66 (#63 or #64) and #65 191
- #67 (hiv or hiv-1* or hiv-2* or hiv1 or hiv2 or "human immunodeficiency virus" or "human immunodeficiency virus" or "human immunodeficiency virus" or "human immune-deficiency virus" or ("human immun*" and "deficiency virus") or "acquired immunodeficiency syndrome" or "acquired immunodeficiency syndrome" or "acquired immuno-deficiency syndrome" or "acquired immune-deficiency syndrome" or ("acquired immun*" and "deficiency syndrome")):ti,ab,kw 16777
- #68 ((antiretroviral* or anti-retroviral* or ARV*) near/2 (compliant* or adheren*)):ti,ab,kw 265
- #69 MeSH descriptor: [Infant, Premature] this term only 3279
- #70 MeSH descriptor: [Infant, Newborn] this term only 14904
- #71 MeSH descriptor: [Infant, Low Birth Weight] this term only 1010
- #72 MeSH descriptor: [Infant, Extremely Low Birth Weight] this term only 103
- #73 MeSH descriptor: [Infant, Small for Gestational Age] this term only 254
- #74 MeSH descriptor: [Infant] this term only 66
- #75 MeSH descriptor: [Infant, Very Low Birth Weight] this term only 802
- #76 MeSH descriptor: [Infant, Postmature] this term only 6
- #77 MeSH descriptor: [Infant, Extremely Premature] this term only 100
- #78 MeSH descriptor: [Child] this term only 225
- #79 MeSH descriptor: [Child, Preschool] this term only 80
- #80 MeSH descriptor: [Adolescent] this term only 91247
- #81 {or #69-#80} and (#63 or #64) 595
- #82 ((antiretroviral* or anti-retroviral* or ARV*) and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)):ti,ab,kw 1265
- #83 MeSH descriptor: [Papillomavirus Infections] this term only and with qualifier(s): [Prevention & control - PC] 285
- #84 MeSH descriptor: [Papillomavirus Vaccines] this term only and with qualifier(s): [Administration & dosage - AD, Therapeutic use - TU] 213
- #85 MeSH descriptor: [Human Papillomavirus Recombinant Vaccine Quadrivalent, Types 6, 11, 16, 18] this term only and with qualifier(s): [Administration & dosage - AD, Therapeutic use - TU] 8
- #86 ((hpv or "papilloma virus*" or papillomavirus*) near/2 (vaccinat* or revaccinat* or immuniz* or immunis* or immunother* or inoculat* or innoculat* or prophyla*)):ti,ab,kw 306
- #87 MeSH descriptor: [Domestic Violence] this term only 159
- #88 MeSH descriptor: [Spouse Abuse] this term only 211
- #89 MeSH descriptor: [Intimate Partner Violence] this term only 34
- #90 MeSH descriptor: [Rape] this term only 105
- #91 (((sexual or domestic or spouse* or "intimate partner") near/3 (violen* or abus*)) or rape):ti,ab,kw 1425
- #92 MeSH descriptor: [Puberty] this term only 292
- #93 (pubert* or pubescen*):ti,ab,kw 1161

- #94 MeSH descriptor: [Menstruation] this term only 467
- #95 (menstruat* or menstrual*):ti,ab,kw 5420
- #96 MeSH descriptor: [Abortion, Legal] this term only 29
- #97 MeSH descriptor: [Abortion, Induced] this term only 936
- #98 (abort* or miscarr* or (pregnan* near/2 terminat*)):ti,ab,kw 4659
- #99 MeSH descriptor: [Infertility] this term only 430
- #100 MeSH descriptor: [Reproductive Techniques, Assisted] this term only 222
- #101 MeSH descriptor: [Fertilization in Vitro] this term only 1927
- #102 (infertil* or "assisted reproductive technolog*" or "in vitro fertili*" or "in-vitro fertili*" or IVF):ti,ab,kw 7649
- #103 MeSH descriptor: [Sexual Behavior] this term only 1721
- #104 MeSH descriptor: [Sex Work] this term only 90
- #105 MeSH descriptor: [Safe Sex] this term only 224
- #106 MeSH descriptor: [Unsafe Sex] this term only 256
- #107 (sex* near (protected or unprotected or safe or unsafe or risk* or behavio*)):ti,ab,kw 7317
- #108 MeSH descriptor: [Contact Tracing] this term only 102
- #109 MeSH descriptor: [Disease Notification] this term only 24
- #110 MeSH descriptor: [Sexual Partners] this term only 542
- #111 (#108 or #109) and #110 31
- #112 (partner* near/3 (notifi* or tracing or report*)):ti,ab,kw 338
- #113 MeSH descriptor: [Prenatal Care] this term only 1322
- #114 (((antenatal or ante-natal or prenatal or pre-natal or antepartum or ante-partum) near/3 (care or service* or counsel* or test*)) or (birth near/3 prepar*)):ti,ab,kw 2581
- #115 MeSH descriptor: [Maternal Health Services] this term only 232
- #116 ((maternal or mother*) near/3 (health or service* or care)):ti,ab,kw 2279
- #117 MeSH descriptor: [Reproductive Health] this term only 64
- #118 (reproductive near/2 (health or care or service*)):ti,ab,kw 545
- #119 MeSH descriptor: [Midwifery] this term only 327
- #120 (midwi* or "skilled birth" or "skilled attendan*"):ti,ab,kw 1200
- #121 MeSH descriptor: [Obstetric Labor Complications] this term only 472
- #122 MeSH descriptor: [Pregnancy Complications] this term only 1493
- #123 ((obstetric* or pregnan* or labour or labor or parturition) near/3 (emergenc* or complication*)):ti,ab,kw 5036
- #124 MeSH descriptor: [Postnatal Care] this term only 391
- #125 MeSH descriptor: [Perinatal Care] this term only 166
- #126 MeSH descriptor: [Postpartum Period] this term only 956
- #127 ((postnatal or post-natal or perinatal or peri-natal or postpartum or post-partum) near/2 (care or service*)):ti,ab,kw 1204

- #128 MeSH descriptor: [Maternal Nutritional Physiological Phenomena] this term only 205
- #129 MeSH descriptor: [Prenatal Nutritional Physiological Phenomena] this term only 121
- #130 MeSH descriptor: [Breast Feeding] this term only 1632
- #131 ("breast feed*" or "breast fed" or breastfeed* or breastfed):ti,ab,kw 3952
- #132 MeSH descriptor: [Early Diagnosis] this term only 581
- #133 #132 and {or #69-#77} 27
- #134 (early near/1 diagnos* near/2 (infant* or neonat* or newborn*)):ti,ab,kw 48
- #135 (diagnos* and (infant* or neonat* or newborn*)):ti,ab,kw 5267
- #136 MeSH descriptor: [Infectious Disease Transmission, Vertical] this term only 586
- #137 (("mother-to-child transmi*" near/3 (prevent* or eliminat*)) or emtct or pmtct or (vertical near transmi*)):ti,ab,kw 835
- #138 MeSH descriptor: [Immunization] explode all trees 4855
- #139 MeSH descriptor: [Immunization Programs] explode all trees 465
- #140 (#138 or #139) and {or #69-#80} 1539
- #141 MeSH descriptor: [Pregnancy] this term only 58
- #142 (#138 or #139) and #141 2
- #143 ((immuniz* or immunis* or vaccinat*) and (infant* or newborn* or neonat* or child* or adolescen* or teen*)):ti,ab,kw 6911
- #144 MeSH descriptor: [Child Health Services] this term only 387
- #145 MeSH descriptor: [Maternal-Child Health Services] explode all trees 11
- #146 MeSH descriptor: [Child Nutrition Disorders] explode all trees 149
- #147 MeSH descriptor: [Infant Nutrition Disorders] explode all trees 99
- #148 MeSH descriptor: [Nutrition Disorders] explode all trees 13509
- #149 MeSH descriptor: [Infant] explode all trees 15237
- #150 MeSH descriptor: [Child, Preschool] this term only 80
- #151 #148 and (#149 or #150) 352
- #152 MeSH descriptor: [Delivery of Health Care, Integrated] this term only 359
- #153 ((integrat* near/3 ("health care" or healthcare or management or treat* or service*) near/3 (child* or schoolchild* or infant* or neonat* or newborn or adolescen* or teen*)) or IMCI or IMNCI):ti,ab,kw 105
- #154 MeSH descriptor: [Guideline Adherence] this term only 999
- #155 MeSH descriptor: [Quality Assurance, Health Care] this term only 806
- #156 (#154 or #155) and (#149 or #150) 52
- #157 (((guideline* or protocol*) near/3 (adher* or observ*)) or "prescribed care") and (infant* or newborn* or neonat* or child*)):ti,ab,kw 303
- #158 MeSH descriptor: [Diarrhea] explode all trees and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Epidemiology - EP, Prevention & control - PC, Therapy - TH] 1754
- #159 #158 and {or #69-#80} 346
- #160 (diarrh* and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)):ti,ab,kw 5413

- #161 MeSH descriptor: [Hand Hygiene] this term only 41
- #162 MeSH descriptor: [Hand Disinfection] this term only 339
- #163 MeSH descriptor: [Water Supply] this term only 163
- #164 MeSH descriptor: [Drinking Water] this term only 81
- #165 MeSH descriptor: [Sanitation] this term only 62
- #166 (handwash* or hand-wash* or (wash* near/1 hand*) or "hand hygiene" or hand-hygiene or soap or "water suppl*" or sanitation or sanitary or "drinking water" or "potable water"):ti,ab,kw 1933
- #167 MeSH descriptor: [Fluid Therapy] this term only 1536
- #168 #167 and {or #69-#79} 107
- #169 ("oral rehydration" near (solution* or salt* or therapy)):ti,ab,kw 587
- #170 MeSH descriptor: [Child Development] this term only 1604
- #171 MeSH descriptor: [Adolescent Development] this term only 78
- #172 ((child* or schoolchild* or adolescen* or teen*) near/2 (develop* or progress*)):ti,ab,kw 4641
- #173 MeSH descriptor: [Breast Neoplasms] this term only and with qualifier(s): [Diagnosis - DI, Diagnostic imaging - DG, Prevention & control - PC] 1770
- #174 MeSH descriptor: [Breast Neoplasms] this term only 10204
- #175 MeSH descriptor: [Mass Screening] this term only 4818
- #176 #173 or (#174 and #175) 1866
- #177 MeSH descriptor: [Uterine Cervical Neoplasms] this term only and with qualifier(s): [Diagnosis - DI, Diagnostic imaging - DG, Prevention & control - PC] 826
- #178 (((breast or cervix or cervical) near (neoplasm* or cancer*)) and (screen* or diagnos*)):ti,ab,kw 6059
- #179 MeSH descriptor: [Folic Acid] this term only and with qualifier(s): [Administration & dosage - AD, Therapeutic use - TU] 1046
- #180 MeSH descriptor: [Folic Acid Deficiency] this term only and with qualifier(s): [Drug therapy - DT, Prevention & control - PC, Therapy - TH] 51
- #181 (folic acid near (fortif* or supplement* or treat* or therap*)):ti,ab,kw 1880
- #182 MeSH descriptor: [Sex Education] this term only 242
- #183 (sex* near (educat* or "health promot*")):ti,ab,kw 1551
- #184 MeSH descriptor: [Pregnancy in Adolescence] this term only 185
- #185 MeSH descriptor: [Kangaroo-Mother Care Method] this term only 42
- #186 (kangaroo near/2 (mother* or infant* or care)):ti,ab,kw 242
- #187 MeSH descriptor: [Anemia] this term only and with qualifier(s): [Drug therapy - DT, Prevention & control - PC] 1051
- #188 MeSH descriptor: [Anemia, Hypochromic] this term only and with qualifier(s): [Drug therapy - DT, Prevention & control - PC] 195
- #189 MeSH descriptor: [Anemia, Iron-Deficiency] this term only and with qualifier(s): [Drug therapy - DT, Prevention & control - PC] 643
- #190 MeSH descriptor: [Pregnancy Complications] this term only 1493
- #191 (#187 or #188 or #189) and #190 49
- #192 ((maternal or mother* or pregnan*) near/2 (nutrition* or folate or folic or iron or anaemi* or anemi*)):ti,ab,kw 1369
- #193 MeSH descriptor: [Malaria] this term only and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Prevention & control - PC] 863

- #194 MeSH descriptor: [Malaria, Falciparum] this term only and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Prevention & control - PC] 1240
- #195 MeSH descriptor: [Malaria, Vivax] this term only and with qualifier(s): [Diagnosis - DI, Drug therapy - DT, Prevention & control - PC] 151
- #196 MeSH descriptor: [Pregnancy] this term only 58
- #197 MeSH descriptor: [Pregnancy Complications, Parasitic] this term only 167
- #198 (#193 or #194 or #195) and (#196 or #197) 124
- #199 ((malaria* or falciparum or vivax) near/3 (pregnan* or mother* or maternal or postpartum or "post partum")):ti,ab,kw 319
- #200 MeSH descriptor: [Smoking Cessation] this term only 3848
- #201 #200 and (#196 or #184) 6
- #202 (((smoking or smoker* or cigarette or tobacco) near/3 (ceas* or cessation or stop* or discontinu*)) and (pregnan* or maternal or mother*)):ti,ab,kw 479
- #203 MeSH descriptor: [Mental Health] this term only 1082
- #204 MeSH descriptor: [Mental Disorders] this term only 2830
- #205 MeSH descriptor: [Mental Health Services] this term only 727
- #206 MeSH descriptor: [Community Mental Health Services] this term only 743
- #207 MeSH descriptor: [Maternal Behavior] this term only 253
- #208 MeSH descriptor: [Mother-Child Relations] this term only 704
- #209 MeSH descriptor: [Paternal Behavior] explode all trees 28
- #210 MeSH descriptor: [Depression, Postpartum] this term only 392
- #211 (((mental or behavio*) near/3 (health or disorder*)) or "postpartum depression" or "post-partum depression" or "postnatal depression" or "post-natal depression"):ti,ab,kw 25906
- #212 {or #46-#62, #66-#68, #81-#107, #111-#131, #133-#137, #140, #142-#147, #151-#153, #156-#157, #159-#166, #168-#172, #176-#186, #191-#192, #198-#199, #201-#211} 115974
- #213 #45 and #212 Publication Year from 2010 to 2017, in Trials 2150 hits

Appendix 2. Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) search strategy

- 1 Family Planning Services/ (24014)
- 2 Contraception/ (18551)
- 3 Reproductive behavior/ or Contraception behavior/ (8119)
- 4 exp Contraceptive agents/ (68445)
- 5 exp Contraceptive Devices/ (23921)
- 6 (condom* or (OC adj pill) or (depot medroxyprogest* or NET-EN or NET EN or Mesigyna or Cyclofem) or (intrauterine system or intra-uterine system or IUS or intrauterine device* or intra-uterine device* or IUD*) or (vasectomy or sterilisation or sterilization or (tubal adj ligation) or ((vaginal adj ring) or cycletel or cycle-tel or ((abstain or abstinen*) adj2 (sex* or intercourse)) or lactational amenorr*))):ti,ab,kw. (55772)
- 7 (contracept* or family planning or (birth adj (control or regulat* or spacing)) or planned parenthood or ((population or fertility) adj (regulat* or control))):ti,ab,kw. (88295)
- 8 Pregnancy in Adolescence/ (7481)
- 9 (pregnan* adj2 (adolescen* or teen* or schoolchild*)):ti,ab,kw. (6460)

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

- 10 Pregnancy, unplanned/ or Pregnancy, unwanted/ (3826)
- 11 (pregnan* adj3 (prevent* or interrupt* or unplanned or unwanted or mistimed)).ti,ab,kw. (12867)
- 12 exp Sexually Transmitted Diseases/di, dt, ep, pc, px, tm [Diagnosis, Drug Therapy, Epidemiology, Prevention & Control, Psychology, Transmission] (203961)
- 13 (sexually transmi* or STI or STIs or STD or STDs or venereal).ti,ab,kw. (43434)
- 14 exp HIV Infections/di, dt, ep, pc, px, tm [Diagnosis, Drug Therapy, Epidemiology, Prevention & Control, Psychology, Transmission] (172773)
- 15 HIV Seropositivity/dt, ep, pc, px, tm [Drug Therapy, Epidemiology, Prevention & Control, Psychology, Transmission] (8281)
- 16 (Anti-HIV Agents/ or Antiretroviral Therapy, Highly Active/) and Medication Adherence/ (1731)
- 17 (hiv or hiv-1* or hiv-2* or hiv1 or hiv2 or human immunodeficiency virus or human immunodeficiency virus or human immuno-deficiency virus or human immune-deficiency virus or (human immun* and deficiency virus) or acquired immunodeficiency syndrome or acquired immunodeficiency syndrome or acquired immuno-deficiency syndrome or acquired immune-deficiency syndrome or (acquired immun* and deficiency syndrome)).ti,ab,kw. (305680)
- 18 ((antiretroviral* or anti-retroviral* or ARV*) adj2 (complian* or adheren*)).ti,ab,kw. (2211)
- 19 (Anti-HIV Agents/ or Antiretroviral Therapy, Highly Active/) and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/ or Adolescent/) (8416)
- 20 ((antiretroviral* or anti-retroviral* or ARV*) and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)).ti,ab,kw. (7849)
- 21 Papillomavirus Infections/pc [Prevention & Control] (4923)
- 22 Papillomavirus Vaccines/ad, tu [Administration & Dosage, Therapeutic Use] (3440)
- 23 Human Papillomavirus Recombinant Vaccine Quadrivalent, Types 6, 11, 16, 18/ad, tu [Administration & Dosage, Therapeutic Use] (66)
- 24 ((hpv or papilloma virus* or papillomavirus*) adj2 (vaccinat* or revaccinat* or immuniz* or immunis* or immunother* or inoculat* or innoculat* or prophyla*)).ti,ab,kw. (4560)
- 25 Domestic Violence/ or Spouse Abuse/ or Intimate Partner Violence/ or Rape/ (18904)
- 26 (((sexual or domestic or spouse* or intimate partner) adj3 (violen* or abus*)) or rape).ti,ab,kw. (29484)
- 27 Puberty/ (12854)
- 28 (pubert* or pubescen*).ti,ab,kw. (35751)
- 29 Menstruation/ (15653)
- 30 (menstruat* or menstrual*).ti,ab,kw. (46842)
- 31 Abortion, Legal/ (7401)
- 32 Abortion, Induced/ (26969)
- 33 (abort* or miscarr* or (pregnan* adj2 terminat*)).ti,ab,kw. (90766)
- 34 Infertility/ (13697)
- 35 Reproductive Techniques, Assisted/ (8232)
- 36 Fertilization in Vitro/ (29054)
- 37 (infertil* or assisted reproductive technolog* or in vitro fertili* or in-vitro fertili* or IVF).ti,ab,kw. (78409)
- 38 Sexual behavior/ or Sex work/ or Safe sex/ or Unsafe sex/ (59074)
- 39 (sex* adj (protected or unprotected or safe or unsafe or risk* or behavio*)).ti,ab,kw. (30964)

- 40 (Contact tracing/ or Disease notification/) and Sexual partners/ (489)
- 41 (partner* adj3 (notifi* or tracing or report*)).ti,ab,kw. (4218)
- 42 Prenatal Care/ (24280)
- 43 (((antenatal or ante-natal or prenatal or pre-natal or antepartum or ante-partum) adj3 (care or service* or counsel* or test*)) or (birth adj3 prepar*)).ti,ab,kw. (24022)
- 44 Maternal Health Services/ (12560)
- 45 ((maternal or mother*) adj3 (health or service* or care)).ti,ab,kw. (23733)
- 46 Reproductive Health/ (2247)
- 47 (reproductive adj2 (health or care or service*)).ti,ab,kw. (11738)
- 48 Midwifery/ (17848)
- 49 (midwi* or skilled birth or skilled attendan*).ti,ab,kw. (21972)
- 50 Obstetric Labor Complications/ (16678)
- 51 Pregnancy Complications/ (85905)
- 52 ((obstetric* or pregnan* or labour or labor or parturition) adj3 (emergenc* or complication*)).ti,ab,kw. (19704)
- 53 Postnatal Care/ (4878)
- 54 Perinatal Care/ (3796)
- 55 Postpartum Period/ (22535)
- 56 ((postnatal or post-natal or perinatal or peri-natal or postpartum or post-partum) adj2 (care or service*)).ti,ab,kw. (5636)
- 57 Maternal Nutritional Physiological Phenomena/ (3299)
- 58 Prenatal Nutritional Physiological Phenomena/ (1591)
- 59 Breast Feeding/ (34451)
- 60 (breast feed* or breast fed or breastfeed* or breastfed).ti,ab,kw. (37063)
- 61 (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/) and Early Diagnosis/ (2277)
- 62 (early adj1 diagnos* adj2 (infant* or neonat* or newborn*)).ti,ab,kw. (378)
- 63 diagnosis.fs. and (infant* or neonat* or newborn*).ti,ab,kw. (84137)
- 64 *"*Infectious Disease Transmission, Vertical"/ (8524)
- 65 ((mother-to-child transmi* adj3 (prevent* or eliminat*)) or emtct or pmtct or (vertical adj transmi*)).ti,ab,kw. (7686)
- 66 (Immunization/ or Immunization, passive/ or Immunization schedule/ or Immunization, secondary/ or Immunization Programs/ or Vaccination/ or Mass vaccination/) and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/ or Adolescent/ or Pregnancy/) (43902)
- 67 ((immuniz* or immunis* or vaccinat*) and (infant* or newborn* or neonat* or child* or adolescen* or teen*)).ti,ab,kw. (45525)
- 68 Child health services/ or Maternal-child health services/ (19947)
- 69 exp child nutrition disorders/ or exp infant nutrition disorders/ or (exp nutrition disorders/ and (exp Infant/ or Child, Preschool/)) (39850)
- 70 "Delivery of Health Care, Integrated"/ (10651)

- 71 ((integrat* adj3 (health care or healthcare or management or treat* or service*) adj3 (child* or schoolchild* or infant* or neonat* or newborn or adolescen* or teen*)) or IMCI or IMNCI).ti,ab,kw. (935)
- 72 (Guideline Adherence/ or Quality Assurance, Health Care/) and (exp Infant/ or Child, Preschool/) (4112)
- 73 (((guideline* or protocol*) adj3 (adher* or observ*)) or "prescribed care") and (infant* or newborn* or neonat* or child*).ti,ab,kw. (1065)
- 74 (Diarrhea/di, dt, ep, pc, th, tm or Diarrhea, Infantile/di, dt, ep, pc, th, tm) and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/ or Adolescent/) (10212)
- 75 (diarrh* and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)).ti,ab,kw. (25541)
- 76 Hand Hygiene/ or Hand Disinfection/ (5743)
- 77 Water Supply/ (31018)
- 78 Drinking Water/ (5487)
- 79 Sanitation/ (6592)
- 80 (handwash* or hand-wash* or (wash* adj1 hand*) or hand hygiene or hand-hygiene or soap or water suppl* or sanitation or sanitary or drinking water or potable water).ti,ab,kw. (79667)
- 81 Fluid Therapy/ and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/) (4522)
- 82 (oral rehydration adj (solution* or salt* or therapy)).ti,ab,kw. (2174)
- 83 Child Development/ or Adolescent Development/ (43982)
- 84 ((child* or schoolchild* or adolescen* or teen*) adj2 (develop* or progress*)).ti,ab,kw. (48454)
- 85 Breast Neoplasms/di, dg, pc or (Breast Neoplasms/ and Mass Screening/) (60213)
- 86 Uterine Cervical Neoplasms/di, dg, pc [Diagnosis, Diagnostic Imaging, Prevention & Control] (22572)
- 87 (((breast or cervix or cervical) adj (neoplasm* or cancer*)) and (screen* or diagnos*)).ti,ab,kw. (70920)
- 88 Folic Acid/ad, tu, th [Administration & Dosage, Therapeutic Use, Therapy] (8376)
- 89 Folic Acid Deficiency/dt, pc, th [Drug Therapy, Prevention & Control, Therapy] (806)
- 90 (folic acid adj (fortif* or supplement* or treat* or therap*)).ti,ab,kw. (3105)
- 91 Sex Education/ (8484)
- 92 (sex* adj (educat* or "health promot*")).ti,ab,kw. (8530)
- 93 Pregnancy in Adolescence/ (7481)
- 94 Kangaroo-Mother Care Method/ (224)
- 95 (kangaroo adj2 (mother or infant or care)).ti,ab,kw. (546)
- 96 (Anemia/dt, pc or Anemia, Hypochromic/dt, pc or Anemia, Iron-Deficiency/dt, pc) and Pregnancy/ (1425)
- 97 ((maternal or mother* or pregnan*) adj2 (nutrition* or folate or folic or iron or anaemi* or anemi*)).ti,ab,kw. (8978)
- 98 (Malaria/di, dt, pc or Malaria, Falciparum/di, dt, pc or Malaria, Vivax/di, dt, pc) and (Pregnancy/ or Pregnancy Complications, Parasitic/) (2012)
- 99 ((malaria* or falciparum or vivax) adj3 (pregnan* or mother* or maternal or postpartum or post partum)).ti,ab,kw. (2199)
- 100 Smoking Cessation/ and (Pregnancy/ or Pregnancy in Adolescence/) (1521)

- 101 (((smoking or smoker* or cigarette or tobacco) adj3 (ceas* or cessation or stop* or discontinu*)) and (pregnan* or maternal or mother*)).ti,ab,kw. (1903)
- 102 Mental health/ or Mental disorders/ or Mental health services/ or Community mental health services/ (207284)
- 103 Maternal behavior/ or Mother-child relations/ or Parenting/ or Paternal behavior/ (39063)
- 104 Depression, Postpartum/ (4500)
- 105 (((mental or behavio*) adj3 (health or disorder*)) or postpartum depression or post-partum depression).ti,ab,kw. (188278)
- 106 or/1-105 (1926054)
- 107 Cell Phones/ (7081)
- 108 Smartphone/ (1248)
- 109 MP3-Player/ (167)
- 110 Computers, Handheld/ (3094)
- 111 ((cell* or mobile*) adj1 (phone* or telephone* or technolog* or device*)).ti,ab,kw. (13097)
- 112 (handheld or hand-held).ti,ab,kw. (9948)
- 113 (smartphone* or smart-phone* or cellphone* or mobiles).ti,ab,kw. (5528)
- 114 ((personal adj1 digital) or (PDA adj3 (device* or assistant*)) or MP3 player* or MP4 player*).ti,ab,kw. (1294)
- 115 (samsung or nokia).ti,ab,kw. (816)
- 116 (windows adj3 (mobile* or phone*)).ti,ab,kw. (43)
- 117 android.ti,ab,kw. (1531)
- 118 (ipad* or i-pad* or ipod* or i-pod* or iphone* or i-phone*).ti,ab,kw. (1959)
- 119 (tablet* adj3 (device* or computer*)).ti,ab,kw. (995)
- 120 Telemedicine/ (16715)
- 121 Videoconferencing/ or Webcasts as topic/ (1495)
- 122 Text Messaging/ (1659)
- 123 Telenursing/ (174)
- 124 (mhealth or m-health or "mobile health" or ehealth or e-health or "electronic health").ti,ab,kw. (15415)
- 125 (telemedicine or tele-medicine or telehealth or tele-health or telecare or tele-care or telenursing or tele-nursing or telepsychiatry or tele-psychiatry or telemonitor* or tele-monitor* or teleconsult* or tele-consult* or telecounsel* or tele-counsel* or telecoach* or tele-coach*).ti,ab,kw. (13879)
- 126 (videoconferenc* or video-conferenc* or webcast* or web-cast*).ti,ab,kw. (2492)
- 127 (((text* or short or voice or multimedia or multi-media or electronic or instant) adj1 messag*) or instant messenger).ti,ab,kw. (3426)
- 128 (texting or texted or texter* or ((sms or mms) adj (service* or messag*)) or interactive voice response* or IVR or voice call* or callback* or voice over internet or VOIP).ti,ab,kw. (2544)
- 129 (Facebook or Twitter or Whatsapp* or Skyp* or YouTube or "You Tube" or Google Hangout*).ti,ab,kw. (4064)
- 130 Mobile Applications/ (2240)
- 131 "mobile app".ti,ab,kw. (1732)
- 132 Social Media/ (3805)
- 133 (social adj (media or network*)).ti,ab,kw. (16003)

- 134 Reminder Systems/ (3057)
- 135 (remind* adj3 (text* or system* or messag*)).ti,ab,kw. (1405)
- 136 Electronic Mail/ (2408)
- 137 (electronic mail* or email* or e-mail or webmail).ti,ab,kw. (11435)
- 138 Medical informatics/ or Medical informatics applications/ (12875)
- 139 Nursing informatics/ or Public health informatics/ (2467)
- 140 ((medical or clinical or health or healthcare or nurs*) adj3 informatics).ti,ab,kw. (5003)
- 141 Multimedia/ (1790)
- 142 Hypermedia/ (399)
- 143 Blogging/ (815)
- 144 (multimedia or multi-media or hypermedia or hyper-media or blog* or vlog* or weblog* or web-log*).ti,ab,kw. (6166)
- 145 Interactive Tutorial/ (248)
- 146 Computer-Assisted Instruction/ (11266)
- 147 ((interactive or computer-assisted) adj1 (tutor* or technolog* or learn* or instruct* or software or communication)).ti,ab,kw. (2214)
- 148 or/107-147 (133860)
- 149 randomized controlled trial.pt. (470403)
- 150 controlled clinical trial.pt. (94471)
- 151 randomized.ab. (403622)
- 152 placebo.ab. (189095)
- 153 drug therapy.fs. (2022689)
- 154 randomly.ab. (280484)
- 155 trial.ab. (423499)
- 156 groups.ab. (1725529)
- 157 or/149-156 (4123563)
- 158 exp animals/ not humans.sh. (4445095)
- 159 157 not 158 (3560201)
- 160 106 and 148 and 159 (4615)
- 161 limit 160 to yr="2010 -Current" (3358)

Appendix 3. Embase Classic + Embase (Ovid) search strategy

- 1 family planning/ (36696)
- 2 contraception/ (46800)
- 3 reproductive behavior/ (919)
- 4 contraceptive behavior/ (2794)
- 5 exp contraceptive agent/ (150341)
- 6 exp contraceptive device/ (40749)

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

- 7 (condom* or (OC adj pill) or (depot medroxyprogest* or NET-EN or NET EN or Mesigyna or Cyclofem) or (intrauterine system or intra-uterine system or IUS or intrauterine device* or intra-uterine device* or IUD*) or (vasectomy or sterilisation or sterilization or (tubal adj ligation)) or ((vaginal adj ring) or cycletel or cycle-tel or ((abstain or abstinen*) adj2 (sex* or intercourse)) or lactational amenorr**)).ti,ab,kw. (66547)
- 8 (concept* or family planning or (birth adj (control or regulat* or spacing)) or planned parenthood or ((population or fertility) adj (regulat* or control))).ti,ab,kw. (95639)
- 9 adolescent pregnancy/ (8690)
- 10 (pregnan* adj2 (adolescen* or teen* or schoolchild*)).ti,ab,kw. (6730)
- 11 unplanned pregnancy/ (4299)
- 12 unwanted pregnancy/ (3137)
- 13 (pregnan* adj3 (prevent* or interrupt* or unplanned or unwanted or mistimed)).ti,ab,kw. (16419)
- 14 exp sexually transmitted disease/di, dt, ep, pc [Diagnosis, Drug Therapy, Epidemiology, Prevention] (36227)
- 15 (sexually transmi* or STI or STIs or STD or STDs or venereal).ti,ab,kw. (56299)
- 16 exp Human immunodeficiency virus infection/di, dt, ep, pc [Diagnosis, Drug Therapy, Epidemiology, Prevention] (161972)
- 17 (hiv or hiv-1* or hiv-2* or hiv1 or hiv2 or human immunodeficiency virus or human immunodeficiency virus or human immuno-deficiency virus or human immune-deficiency virus or (human immun* and deficiency virus) or acquired immunodeficiency syndrome or acquired immunodeficiency syndrome or acquired immuno-deficiency syndrome or acquired immune-deficiency syndrome or (acquired immun* and deficiency syndrome)).ti,ab,kw. (372022)
- 18 ((antiretroviral* or anti-retroviral* or ARV*) adj2 (complian* or adheren*)).ti,ab,kw. (2598)
- 19 (antiretroviral therapy/ or highly active antiretroviral therapy/) and medication compliance/ (703)
- 20 (antiretroviral therapy/ or highly active antiretroviral therapy/) and (child/ or infant/ or adolescent/ or newborn/) (4203)
- 21 ((antiretroviral* or anti-retroviral* or ARV*) and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)).ti,ab,kw. (10214)
- 22 papillomavirus infection/pc [Prevention] (2243)
- 23 Wart virus vaccine/ad, dt [Drug Administration, Drug Therapy] (6083)
- 24 ((hvp or papilloma virus* or papillomavirus*) adj2 (vaccinat* or revaccinat* or immuniz* or immunis* or immunother* or inoculat* or innoculat* or prophyla*)).ti,ab,kw. (5974)
- 25 domestic violence/ or battered woman/ or family violence/ or exp partner violence/ (19311)
- 26 statutory rape/ or acquaintance rape/ or rape/ or marital rape/ (7284)
- 27 (((sexual or domestic or spouse* or intimate partner) adj3 (violen* or abus*)) or rape).ti,ab,kw. (36302)
- 28 puberty/ or menarche/ (36986)
- 29 (pubert* or pubescen*).ti,ab,kw. (50840)
- 30 menstruation/ (22869)
- 31 (menstruat* or menstrual*).ti,ab,kw. (62512)
- 32 abortion/ or imminent abortion/ or recurrent abortion/ or septic abortion/ or spontaneous abortion/ (74082)
- 33 (abort* or miscarr* or (pregnan* adj2 terminat*)).ti,ab,kw. (120224)
- 34 infertility/ (38578)
- 35 infertility therapy/ or in vitro fertilization/ (19053)
- 36 (infertil* or assisted reproductive technolog* or in vitro fertili* or in-vitro fertili* or IVF).ti,ab,kw. (116136)

- 37 sexual behavior/ or adolescent sexual behavior/ or casual sex/ or prostitution/ or exp safe sex/ or sexual practice/ or exp unsafe sex/ (110380)
- 38 (sex* adj (protected or unprotected or safe or unsafe or risk* or behavio*)).ti,ab,kw. (33582)
- 39 contact examination/ (3275)
- 40 (partner* adj3 (notifi* or tracing or report*)).ti,ab,kw. (5304)
- 41 prenatal care/ or prenatal screening/ (40985)
- 42 (((antenatal or ante-natal or prenatal or pre-natal or antepartum or ante-partum) adj3 (care or service* or counsel* or test*)) or (birth adj3 prepar*)).ti,ab,kw. (30891)
- 43 maternal health service/ (467)
- 44 ((maternal or mother*) adj3 (health or service* or care or welfare)).ti,ab,kw. (28175)
- 45 reproductive health/ (13171)
- 46 (reproductive adj2 (health or care or service*)).ti,ab,kw. (15081)
- 47 midwife/ or nurse midwife/ (28959)
- 48 (midwi* or skilled birth or skilled attendan*).ti,ab,kw. (24665)
- 49 labor complication/ (9228)
- 50 pregnancy complication/ (71410)
- 51 ((obstetric* or pregnan* or labour or labor or parturition) adj3 (emergenc* or complication*)).ti,ab,kw. (32301)
- 52 postnatal care/ or newborn care/ (16770)
- 53 perinatal care/ (13024)
- 54 maternal care/ or maternal welfare/ (27759)
- 55 maternal nutrition/ (10080)
- 56 puerperium/ (37622)
- 57 ((postnatal or post-natal or perinatal or peri-natal or postpartum or post-partum) adj2 (care or service*)).ti,ab,kw. (7276)
- 58 breast feeding/ (44726)
- 59 (breast feed* or breast fed or breastfeed* or breastfed).ti,ab,kw. (45447)
- 60 early diagnosis/ and (exp infant/ or newborn/) (6264)
- 61 (early adj1 diagnos* adj2 (infant* or neonat* or newborn*)).ti,ab,kw. (556)
- 62 diagnosis.fs. and (infant* or neonat* or newborn*).ti,ab,kw. (103847)
- 63 vertical transmission/ (12794)
- 64 ((mother-to-child transmi* adj3 (prevent* or eliminat*)) or emtct or pmtct or (vertical adj transmi*)).ti,ab,kw. (9842)
- 65 (immunization/ or mass immunization/ or vaccination/) and (exp infant/ or newborn/ or exp child/ or adolescent/ or pregnancy/) (51587)
- 66 ((immuniz* or immunis* or vaccinat*) and (infant* or newborn* or neonat* or child* or adolescen* or teen* or pregnan*)).ti,ab,kw. (65983)
- 67 child health care/ or early childhood intervention/ or maternal child health care/ (37990)
- 68 exp nutritional disorder/ and (preschool child/ or exp infant/) (58303)
- 69 integrated health care system/ (9211)

- 70 ((integrat* adj3 (health care or healthcare or management or treat* or service*) adj3 (child* or schoolchild* or infant* or neonat* or newborn or adolescen* or teen*)) or IMCI or IMNCI).ti,ab,kw. (1093)
- 71 (protocol compliance/ or health care quality/) and (preschool child/ or exp infant/) (6812)
- 72 (((guideline* or protocol*) adj3 (adher* or observ*)) or "prescribed care") and (infant* or newborn* or neonat* or child*).ti,ab,kw. (1754)
- 73 infantile diarrhea/di, dm, dt, ep, pc, th [Diagnosis, Disease Management, Drug Therapy, Epidemiology, Prevention, Therapy] (1735)
- 74 diarrhea/di, dm, dt, ep, pc, th and (exp infant/ or newborn/ or exp child/ or adolescent/ or pregnancy/) (6447)
- 75 (diarrh* and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*)).ti,ab,kw. (33784)
- 76 hand washing/ or hand disinfection/ (11612)
- 77 water supply/ (34795)
- 78 drinking water/ (42849)
- 79 sanitation/ (13878)
- 80 (handwash* or hand-wash* or (wash* adj1 hand*) or hand hygiene or hand-hygiene or soap or water suppl* or sanitation or sanitary or drinking water or potable water).ti,ab,kw. (106006)
- 81 oral rehydration therapy/ (2412)
- 82 (oral rehydration adj (solution* or salt* or therapy)).ti,ab,kw. (2252)
- 83 child development/ or adolescent development/ (45720)
- 84 ((child* or schoolchild* or adolescen* or teen*) adj2 (develop* or progress*)).ti,ab,kw. (57900)
- 85 breast cancer/di, dm, dt, pc [Diagnosis, Disease Management, Drug Therapy, Prevention] (95730)
- 86 breast cancer/ and cancer screening/ (15585)
- 87 uterine cervix cancer/di, dm, dt, pc [Diagnosis, Disease Management, Drug Therapy, Prevention] (16909)
- 88 (((breast or cervix or cervical) adj (neoplasm* or cancer*)) and (screen* or diagnos*)).ti,ab,kw. (110687)
- 89 folic acid/ad, dt [Drug Administration, Drug Therapy] (11586)
- 90 folic acid deficiency/dm, dt, pc, th [Disease Management, Drug Therapy, Prevention, Therapy] (1174)
- 91 (folic acid adj (fortif* or supplement* or treat* or therap*)).ti,ab,kw. (4146)
- 92 sexual education/ (10956)
- 93 (sex* adj (educat* or "health promot*")).ti,ab,kw. (8934)
- 94 kangaroo care/ (720)
- 95 (kangaroo adj2 (mother or infant or care)).ti,ab,kw. (725)
- 96 (anemia/dt, pc or iron deficiency anemia/dt, pc) and pregnancy/ (1212)
- 97 ((maternal or mother* or pregnan*) adj2 (nutrition* or folate or folic or iron or anaemi* or anemi*)).ti,ab,kw. (10422)
- 98 (malaria/di, dm, dt, pc or malaria, falciparum/di, dm, dt, pc or malaria, vivax/di, dm, dt, pc) and (pregnancy/ or pregnancy complication/) (1493)
- 99 ((malaria* or falciparum or vivax) adj3 (pregnan* or mother* or maternal or postpartum or post partum)).ti,ab,kw. (2728)
- 100 smoking cessation/ and (pregnancy/ or adolescent pregnancy/) (1947)
- 101 (((smoking or smoker* or cigarette or tobacco) adj3 (ceas* or cessation or stop* or discontinu*)) and (pregnan* or maternal or mother*)).ti,ab,kw. (2318)
- 102 mental health/ or community mental health/ or mental health service/ (156252)

- 103 maternal behavior/ or parental behavior/ or paternal behavior/ (22837)
- 104 puerperal depression/ (8364)
- 105 (((mental or behavio*) adj3 (health or disorder*)) or postpartum depression or post-partum depression or postnatal depression or post-natal depression).ti,ab,kw. (246537)
- 106 or/1-105 (2439735)
- 107 mobile phone/ or smartphone/ (16146)
- 108 mp3 player/ (162)
- 109 ((cell* or mobile*) adj1 (phone* or telephone* or technolog* or device*)).ti,ab,kw. (16688)
- 110 (handheld or hand-held).ti,ab,kw. (13317)
- 111 (smartphone* or smart-phone* or cellphone* or mobiles).ti,ab,kw. (7717)
- 112 ((personal adj1 digital) or (PDA adj3 (device* or assistant*)) or MP3 player* or MP4 player*).ti,ab,kw. (1692)
- 113 (samsung or nokia).ti,ab,kw. (1456)
- 114 (windows adj3 (mobile* or phone*)).ti,ab,kw. (67)
- 115 android.ti,ab,kw. (2452)
- 116 (ipad* or i-pad* or ipod* or i-pod* or iphone* or i-phone*).ti,ab,kw. (3612)
- 117 (tablet* adj3 (device* or computer*)).ti,ab,kw. (1571)
- 118 telemedicine/ or telecardiology/ or teleconsultation/ or teledermatology/ or telediagnosis/ or telemonitoring/ or telepathology/ or telepsychiatry/ or teleradiotherapy/ or telesurgery/ or teletherapy/ (27986)
- 119 videoconferencing/ or webcast/ (2824)
- 120 text messaging/ (2877)
- 121 telenursing/ (203)
- 122 (mhealth or m-health or "mobile health" or ehealth or e-health or "electronic health").ti,ab,kw. (19595)
- 123 (telemedicine or tele-medicine or telehealth or tele-health or telecare or tele-care or telenursing or tele-nursing or telepsychiatry or tele-psychiatry or telemonitor* or tele-monitor* or teleconsult* or tele-consult* or telecounsel* or tele-counsel* or telecoach* or tele-coach*).ti,ab,kw. (17704)
- 124 (videoconferenc* or video-conferenc* or webcast* or web-cast*).ti,ab,kw. (3335)
- 125 (((text* or short or voice or multimedia or multi-media or electronic or instant) adj1 messag*) or instant messenger).ti,ab,kw. (4491)
- 126 (texting or texted or texter* or ((sms or mms) adj (service* or messag*)) or interactive voice response* or IVR or voice call* or callback* or voice over internet or VOIP).ti,ab,kw. (3560)
- 127 (Facebook or Twitter or Whatsapp* or Skyp* or YouTube or "You Tube" or Google Hangout*).ti,ab,kw. (5883)
- 128 mobile application/ (4502)
- 129 "mobile app".ti,ab,kw. (2078)
- 130 social media/ (9110)
- 131 (social adj (media or network*)).ti,ab,kw. (20813)
- 132 reminder system/ (2143)
- 133 (remind* adj3 (text* or system* or messag*)).ti,ab,kw. (1962)
- 134 e-mail/ (15606)

- 135 (electronic mail* or email* or e-mail or webmail).ti,ab,kw. (23089)
- 136 medical informatics/ (17801)
- 137 nursing informatics/ (1286)
- 138 ((medical or clinical or health or healthcare or nurs*) adj3 informatics).ti,ab,kw. (7055)
- 139 multimedia/ (3205)
- 140 hypermedia/ (371)
- 141 blogging/ (141)
- 142 (multimedia or multi-media or hypermedia or hyper-media or blog* or vlog* or weblog* or web-log*).ti,ab,kw. (9103)
- 143 teaching/ (85381)
- 144 ((interactive or computer-assisted) adj1 (tutor* or technolog* or learn* or instruct* or software or communication)).ti,ab,kw. (3142)
- 145 or/107-144 (260905)
- 146 106 and 145 (29013)
- 147 crossover procedure/ (53072)
- 148 double blind procedure/ (143697)
- 149 randomized controlled trial/ (465243)
- 150 single-blind procedure/ (28784)
- 151 random\$.tw. (1234071)
- 152 factorial\$.tw. (31351)
- 153 (crossover\$ or cross over\$ or cross-over\$).tw. (90817)
- 154 placebo\$.tw. (263675)
- 155 (doubl\$ adj blind\$).tw. (184763)
- 156 (singl\$ adj blind\$).tw. (19947)
- 157 assign\$.tw. (323334)
- 158 allocat\$.tw. (119980)
- 159 volunteer\$.tw. (228637)
- 160 or/147-159 (1925757)
- 161 146 and 160 (4486)
- 162 limit 161 to yr="2010 -Current" (3567)
- 163 limit 162 to embase (1725)

Appendix 4. POPLINE search strategy

All Fields: ((cell OR cellular OR mobile) AND (phone OR phones OR telephone OR telephones OR technology OR technologies OR device OR devices)) OR smartphone OR smartphones OR smart-phone OR smart-phones OR cellphone OR cellphones OR mobiles OR mhealth OR m-health OR "mobile health" OR ehealth OR e-health OR "electronic health" OR telemedicine OR tele-medicine OR telehealth OR tele-health OR telecare OR tele-care OR telenursing OR tele-nursing OR telepsychiatry OR tele-psychiatry OR telemonitor OR telemonitoring OR tele-monitor OR tele-monitoring OR teleconsult OR teleconsulting OR tele-consult OR tele-consulting OR telecounsel OR telecounseling OR tele-counsel OR tele-counseling OR telecoach OR telecoaching OR tele-coach OR tele-coaching OR videoconference OR videoconferences OR videoconferencing OR video-conference OR video-conferences OR video-conferencing OR webcast OR webcasts OR webcasting OR web-cast OR web-casts OR web-casting OR ((text OR texts OR texting OR short OR voice OR multimedia OR multi-media OR electronic OR instant)

AND (message OR messages OR messaging) OR "instant messenger" OR texting OR texted OR texter OR texters OR ((sms OR mms) AND (service OR services OR message OR messages OR messaging)) OR "interactive voice response" OR "interactive voice responses" OR ivr OR "voice call" OR "voice calls" OR callback OR "voice over internet" OR voip OR "mobile app" OR "mobile apps" OR "mobile application" OR "mobile applications" OR "social media" OR ((medical OR clinical OR health OR healthcare OR nurse OR nurses OR nursing) AND informatics) OR Keyword: × TEXT MESSAGING OR × MOBILE DEVICES OR × INFORMATION COMMUNICATION TECHNOLOGY OR × CELLULAR PHONE

AND

All Fields: (randomised OR randomized OR "randomly allocated" OR "random allocation" OR "controlled trial" OR "control group" OR "control groups" OR trial) OR

Keyword: × QUANTITATIVE RESEARCH OR × QUANTITATIVE EVALUATION OR

× RESEARCH METHODOLOGY OR × CLINICAL TRIALS OR × CONTROL GROUPS - 1006 hits

Appendix 5. WHO Global Health Library search strategy

(tw:(((cell* OR mobile*) AND (phone* OR telephone* OR technolog* OR device*)) OR smartphone* OR smart-phone* OR cellphone* OR mobiles OR mhealth OR m-health OR "mobile health" OR ehealth OR e-health OR "electronic health" OR telemedicine OR tele-medicine OR telehealth OR tele-health OR telecare OR tele-care OR telenursing OR tele-nursing OR telepsychiatry OR tele-psychiatry OR telemonitor* OR tele-monitor* OR teleconsult* OR tele-consult* OR telecounsel* OR tele-counsel* OR telecoach* OR tele-coach* OR videoconferenc* OR video-conferenc* OR webcast* OR web-cast* OR ((text* OR short OR voice OR multimedia OR multi-media OR electronic OR instant) AND messag*) OR "instant messenger" OR texting OR texted OR texter* OR ((sms OR mms) AND (service* OR messag*)) OR "interactive voice response*" OR ivr OR "voice call*" OR callback* OR "voice over internet" OR voip OR "mobile app*" OR (social AND (media OR network*)) OR ((medical OR clinical OR health OR healthcare OR nurs*) AND informatics)) OR (mh:("Telemedicine" OR "Cell Phones" OR "Internet" OR "Mobile Applications" OR "Medical Informatics" OR "Information Technology" OR "Smartphone")) AND (mh:("Controlled Clinical Trials, Randomized" OR "Controlled Clinical Trials as Topic" OR "Controlled Clinical Trial" OR "Clinical Trial")) OR (tw:(randomised OR randomized OR "randomly allocated" OR "random allocation" OR "controlled trial" OR "control group" OR "control groups" OR trial)) – 1121 hits

Appendix 6. WHO ICTRP search strategy

Search 1:

Title: reproductive health OR maternal health OR child health OR adolescent health OR immunization OR immunisation OR pregnancy

AND

Intervention: mobile device OR mobiles OR smartphone OR phone OR cellphone

Result: 80 hits

Search 2:

Title: mobile device OR mobiles OR smartphone OR phone OR cellphone

AND

Intervention: reproductive health OR maternal health OR child health OR adolescent health

Result: 22 hits

Search 3:

Intervention: sexually transmitted OR HIV OR nutrition OR mental health OR family planning OR contraception OR abortion OR prenatal OR postnatal

AND

Title: mobile device OR mobiles OR smartphone OR phone OR cellphone

Result: 240 hits

Search 4:

Title: mobile device OR mobiles OR smartphone OR phone OR cellphone

AND

Intervention: sexually transmitted OR HIV OR nutrition OR mental health OR family planning OR contraception OR abortion OR prenatal OR postnatal

Result: 101 hits

Search 5:

Title: sexual behavior OR sexual behaviour OR sexual health OR safe sex OR unsafe sex OR sex education OR breastfeeding OR integrated delivery

AND

Intervention: mobile device OR mobiles OR smartphone OR phone OR cellphone

Result: 41 hits

Search 6:

Title: mobile device OR mobiles OR smartphone OR phone OR cellphone

AND

Intervention: sexual behavior OR sexual behaviour OR sexual health OR safe sex OR unsafe sex OR sex education OR breastfeeding OR integrated delivery

Result: 90 hits

Amalgamated Results (duplicates removed): 492 hits

Appendix 7. ClinicalTrials.gov search strategy

Search 1: ("reproductive health" OR "maternal health" OR "child health" OR "adolescent health" OR immunization OR immunisation OR pregnancy) AND ("mobile phone" OR "mobile phones" OR "mobile devices" OR mobiles OR smartphone OR smartphones) | Child, Adult | Studies received on or after 01/01/2000 | Studies updated on or before 08/31/2017 – 275 hits

Search 2: ("sexually transmitted" OR HIV OR nutrition OR "mental health" OR "family planning" OR contraception OR abortion OR prenatal OR postnatal) AND ("mobile phone" OR "mobile phones" OR "mobile devices" OR mobiles OR smartphone OR smartphones) | Child, Adult | Studies received on or after 01/01/2000 | Studies updated on or before 08/31/2017 – 481 hits

Search 3: ("sexual behavior" OR "sexual behaviour" OR "sexual health" OR "safe sex" OR "unsafe sex" OR "sex education" OR breastfeeding OR "integrated delivery") AND ("mobile phone" OR "mobile phones" OR "mobile devices" OR mobiles OR smartphone OR smartphones) | Child, Adult | Studies received on or after 01/01/2000 | Studies updated on or before 08/31/2017 – 180 hits

HISTORY

Review first published: Issue 8, 2020

CONTRIBUTIONS OF AUTHORS

Conceiving the protocol: CG, TT, SL, GM.

Designing the protocol: CG, TT, SL, GM, MP, CF, NH.

Designing search strategies: JE, CF, MP.

Writing the protocol: MP, CF, TT.

Providing general advice on the protocol: CG, SL, GV, NH, NM, HB, MF.

Securing funding for the protocol and review: GM, TT, SL, CG, CF.

Data collection for the review: NH, GV, NM, HB, MP.

Data management for the review: NH, GV, NM, HB, MP.

Analysis of data: NH, GV, NM, HB, MP.

Targeted client communication via mobile devices for improving maternal, neonatal, and child health (Review)

Copyright © 2020 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Interpretation of data: MP, CF, GC, TT SL.

Writing the review: MP, CF, CG.

Providing general advice on the review: CG, SL, GV, NH, NM, HB, MF.

DECLARATIONS OF INTEREST

MP: was contracted by the World Health Organization (WHO) to produce this review.

NH: is employed by Cochrane Response, an evidence services unit operated by Cochrane. The WHO contacted Cochrane Response to produce this review.

HB: is employed by Cochrane Response, an evidence services unit operated by Cochrane. The WHO contacted Cochrane Response to produce this review.

GV: is employed by Cochrane Response, an evidence services unit operated by Cochrane. The WHO contacted Cochrane Response to produce this review.

NM: previously worked for Enhanced Reviews Ltd, a company that conducts systematic reviews mostly for the public sector. NM is employed by Cochrane Response, an evidence services unit operated by Cochrane. The WHO contacted Cochrane Response to produce this review.

TT: none known.

GM: owns stock in Apple Computer.

CG: none known.

SL: is the Joint Co-ordinating Editor for the Cochrane Effective Practice and Organisation of Care Review Group.

MF: none known.

CF: was contracted by the WHO to produce this review.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- World Health Organization, Switzerland

This work, including the open access publication costs, was funded by the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a co-sponsored program executed by the World Health Organization (WHO).

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

In the protocol, we specified that the 'Summary of findings' tables for pregnant and postpartum populations would include the following outcomes: maternal morbidity and mortality combined; neonatal morbidity and mortality combined; initiation of breastfeeding; attendance for antenatal care; birth in a health facility; and unintended consequences ([Palmer 2018a](#)). In conducting the review, we decided to include any measure of breastfeeding, instead of only including *initiation* of breastfeeding, and any measure of access to intrapartum care, thereby including births in a health facility *and* births occurring with a skilled attendant present.

We had planned that when a study reported the same outcome measure for multiple time points of follow-up, we would extract data from all time points. However, in the review, we extracted data for the outcome at the longest follow-up point.

For the purpose of pooling data, we planned to categorise lengths of follow-up as follows: short-term follow-up: three months or less; moderate-term follow-up: three to 12 months; long-term follow-up: greater than 12 months. However, given the limited number of studies with the same aim, comparison, and outcome measure that could be pooled, we decided to pool across different lengths of follow-up.

We planned to carry out the following subgroup analyses for the objective outcomes of health status: income region (by World Bank income group) ([World Bank 2017](#)); and delivery mechanisms (i.e. mobile phone messaging only, mobile applications only, combined mobile phone messaging and applications, combined application, and other). However, there were an insufficient number of studies reporting the same objective health status outcomes to conduct these subgroup analyses.

We planned to carry out the following sensitivity analyses: only including studies with low risk of bias on the sequence generation, allocation concealment, and incomplete outcome data domains; and only including studies with *objectively* measured outcomes. However, there were insufficient studies to conduct the second specified sensitivity analyses.

As part of the risk of bias assessments of included studies, we also reported an assessment of 'Other bias'. Under this domain, we considered other potential sources of bias such as the presence of baseline imbalances related to the outcome under study, and evidence of contamination.

In the protocol, we stated we would not pool studies with substantial heterogeneity in meta-analyses. However, some of the pooled analyses do exhibit substantial statistical heterogeneity. We intended to explore possible reasons for variability by conducting our prespecified subgroup analysis; however, there was an insufficient number of studies in the pooled analyses to conduct meaningful subgroup analyses. Where we noted other potential explanations for high heterogeneity (e.g. differing baseline level of risk), and there were a sufficient number of studies, subgroup analyses were conducted to examine these.

NOTES

The protocol for this review is based on standard text and guidance provided by Cochrane Consumers and Communication ([Ryan 2016](#)).

INDEX TERMS

Medical Subject Headings (MeSH)

Breast Feeding [statistics & numerical data]; *Cell Phone; Child Health [*standards] [statistics & numerical data]; *Communication; Delivery, Obstetric [standards]; Health Behavior; *Health Services Needs and Demand; Health Status; HIV Infections [drug therapy]; Infant Health [*standards] [statistics & numerical data]; Maternal Health [*standards] [statistics & numerical data]; Medication Adherence [statistics & numerical data]; Postpartum Period; Prenatal Care [statistics & numerical data]; Quality Improvement; Randomized Controlled Trials as Topic; Text Messaging

MeSH check words

Child, Preschool; Female; Humans; Infant; Infant, Newborn; Pregnancy