

Criterion validity of two physical activity and one sedentary time questionnaire against accelerometry in a large cohort of adults and older adults

Edvard H Sagelv ¹, Laila A Hopstock,² Jonas Johansson,² Bjørge H Hansen,³ Soren Brage,^{4,5} Alexander Horsch,⁶ Ulf Ekelund,^{7,8} Bente Morseth¹

To cite: Sagelv EH, Hopstock LA, Johansson J, *et al*. Criterion validity of two physical activity and one sedentary time questionnaire against accelerometry in a large cohort of adults and older adults. *BMJ Open Sport & Exercise Medicine* 2020;**6**:e000661. doi:10.1136/bmjsem-2019-000661

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2019-000661>).

Accepted 6 February 2020

ABSTRACT

Objectives We compared the ability of physical activity and sitting time questionnaires (PAQ) for ranking individuals versus continuous volume calculations (physical activity level (PAL), metabolic equivalents of task (MET), sitting hours) against accelerometry measured physical activity as our criterion.

Methods Participants in a cohort from the Tromsø Study completed three questionnaires; (1) The Saltin-Grimby Physical Activity Level Scale (SGPALS) (n=4040); (2) The Physical Activity Frequency, Intensity and Duration (PAFID) questionnaire (n=5902) calculated as MET-hours-week⁻¹ and (3) The International Physical Activity questionnaire (IPAQ) short-form sitting question (n=4896). We validated the questionnaires against the following accelerometry (Actigraph wGT3X-BT) estimates: vector magnitude counts per minute, steps-day⁻¹, time (minutes-day⁻¹) in sedentary behaviour, light physical activity, moderate and vigorous physical activity (MVPA) non-bouted and ≥10 min bouted MVPA.

Results Ranking of physical activity according to the SGPALS and quartiles (Q) of MET-hours-week⁻¹ from the PAFID were both positively associated with accelerometry estimates of physical activity (p<0.001) but correlations with accelerometry estimates were weak (SGPALS (PAL): r=0.11 to 0.26, p<0.001) and weak-to-moderate (PAFID: r=0.39 to 0.44, p<0.01). There was 1 hour of accelerometry measured sedentary time from Q1 to Q4 in the IPAQ sitting question (p<0.001) and also weak correlations (r=0.22, p<0.01).

Conclusion Ranking of physical activity levels measured with PAQs appears to have higher validity than energy expenditure calculations. Self-reported sedentary time poorly reflects accelerometry measured sedentary time. These two PAQs can be used for ranking individuals into different physical activity categories supporting previous studies using these instruments when assessing associations with health outcomes.

INTRODUCTION

Physical activity surveillance at population level may support public health initiatives and allow researchers to track physical activity levels and patterns over time.¹ Physical activity

Summary box

What are the new findings

- Ranking of the two included physical activity questionnaires reduces information content but may be the optimal way of processing self-reported physical activity.
- Volume calculations (physical activity level, metabolic equivalents of task hours) allow the biases associated with self-reported physical activity to be more pronounced.
- Self-reported sitting time shows low validity and does not reflect accelerometry measured sedentary time.

is traditionally measured using self-reported methods such as questionnaires.² However, the validity of physical activity questionnaires (PAQ) is threatened by recall and social desirability bias, resulting in imprecise assessments.³⁻⁶ Nevertheless, PAQs have over the years led to valuable knowledge on the effect of physical activity on health outcomes and mortality.⁷⁻¹⁴

Validation of PAQs is crucial to guide researchers when interpreting associations between self-reported physical activity and health outcomes. Moreover, PAQs may inherit different measurement properties. For example, one of the first developed PAQs, by Saltin and Grimby¹⁵ named 'Saltin-Grimby Physical Activity Level Scale' (SGPALS),^{16 17} ranks individuals by physical activity levels. A more recent PAQ, the Physical Activity Frequency, Intensity and Duration (PAFID) questionnaire,¹⁸ allows the answers to be summed up as total physical activity volume (ie, energy expenditure, metabolic equivalents of task (MET)-hours per week). Finally, sedentary behaviour has been suggested as a risk factor for disease and mortality, which is also commonly assessed by PAQs,^{19 20} such as the International Physical



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Edvard H Sagelv;
edvard.h.sagelv@uit.no

Activity Questionnaire (IPAQ) short-form sitting question.²¹ Both PAQs (SGPALS,^{16 22–26} PAFID^{18 27}) and the IPAQ short-form sitting question²¹ have previously been validated, however, the studies that compare these questionnaires against accelerometry are characterised by small sample sizes.^{18 21 23} As population samples are heterogeneous and consequently result in heterogeneous findings, validation studies based on small samples may have limited representability. Furthermore, considering that already established longitudinal population cohorts have implemented PAQs from inception allowing for long follow-up time (SGPALS: >45 years,^{28–30} PAFID: >35 years³¹), validation of PAQs and sitting questionnaires against accelerometry measured physical activity and sitting time from large heterogeneous samples will allow researchers to more accurately interpret results from longitudinal cohort studies where only questionnaires are the physical activity and sedentary time measure.

Moreover, although PAQs can inherit different measurement properties, the methods for processing the PAQs can result in similar expressions (eg, ranking of the SGPALS can be summarised as volume,²⁵ volume calculations can be grouped as quartiles), and thus the processing of questionnaires may also influence the validity differently.

We aimed to assess the validity of two PAQs inheriting different measurement properties; ranking of physical activity levels (SGPALS), volume calculations (PAFID) and one sedentary time questionnaire (IPAQ sitting short-form), by using accelerometry as our criterion, in a large heterogeneous sample of adults and older adults. Additionally, we aimed to assess how ranking and volume calculations of the PAQs reflects accelerometry measured physical activity and sedentary time.

METHODS

Design

We used participants from the seventh wave of the population-based cohort study named The Tromsø Study, which is conducted in Tromsø, Northern Norway. The study includes seven waves of data collection (Tromsø 1: 1974, Tromsø 2: 1979 to 1980, Tromsø 3: 1986 to 1987, Tromsø 4: 1994 to 1995, Tromsø 5: 2001, Tromsø 6: 2007 to 2008, Tromsø 7: 2015 to 2016) (details described elsewhere³²).

Participants

All inhabitants in Tromsø municipality aged 40 years and older were invited to Tromsø 7. A total of 21 083 (65% of 32 591 invited participants) participants attended a first visit including questionnaires, biological sampling and clinical examinations. A random selection of 8346 participants attended a second visit at a later time point (>7 days), where 6778 participants were invited to wear an accelerometer, of which 6332 (93%) participants accepted. Of those who provided valid accelerometry data, 4040 participants completed both the leisure time and occupational time SGPALS; 5902 participants

completed the PAFID questionnaire, and 5186 and 5088 participants completed the sitting question from the IPAQ short-form for week and weekend, respectively, where 4896 completed both.

All participants gave written informed consent.

Patient and public involvement

The Tromsø study advisory board includes patient (University hospital of Northern Norway) and public (eg, Norwegian Health Association, Tromsø municipality) representatives. Some participants are invited as ambassadors when data collection is ongoing, where they actively contribute to recruitment of participants. We have together with the Norwegian Health Association provided individual feedback on levels of physical activity to participants in Tromsø 7. There was no public involvement when designing this study.

Data collection

Height and weight were measured in light clothing without shoes. Body mass index (BMI) was calculated (kg/m^2) and defined as normal and underweight ($<25 \text{ kg}/\text{m}^2$), overweight (25 to $29.9 \text{ kg}/\text{m}^2$) and obese ($\geq 30 \text{ kg}/\text{m}^2$). Educational level was collected from questionnaires and categorised in; (1) primary school, (2) high school diploma, (3) university education <4 years and (4) university education ≥ 4 years.

The physical activity and sitting questionnaires

The Saltin-Grimby physical activity level scale

The SGPALS asks participants to rank their leisure time and occupational time physical activity level separately, choosing one of four options. Based on the idea of the original questionnaire by Saltin and Grimby¹⁵, the SGPALS used in Tromsø 7 is a slight modification of Saltin and Grimby¹⁵ according to Rödger *et al.*¹⁷ The SGPALS is presented in online supplementary table 1.

We computed the SGPALS as combined leisure time and occupational time where individuals were categorised as (1) *inactive*, (2) *moderately inactive*, (3) *moderately active* and (4) *active* according to Wareham *et al.*³³ with some modifications. In order to calculate physical activity volume, we assigned a physical activity level (PAL) value from the combined leisure time and occupational time SGPALS, which we derived from a previous validation study that calculated PAL as energy expenditure obtained from doubly labelled water divided by the estimated basal metabolic rate.²⁵ The classifications and the assigned PAL value are presented in table 1.

The physical activity frequency, intensity and duration questionnaire

The PAFID questionnaire (table 2) includes three questions referring to frequency, intensity and duration of physical activity. We generated an index to reflect METs by multiplying intensity (METs) by duration (minutes) by frequency (times per week), and the outcome was expressed as MET-hours per week.^{34 35} We also grouped

Table 1 Physical activity classification by the combined leisure time and occupational time SGPALS (n=4040)

	Light LPA (n=532)	Moderate LPA (n=2429)	Hard LPA (n=969)	Very hard LPA (n=109)
Light OPA (n=2263)	Inactive (n=349, 8.6%) PAL: 1.4	Moderately inactive (n=1346, 33.3%) PAL: 1.5	Active (n=507, 12.6%) PAL: 1.7	Active (n=61, 1.5%) PAL: 1.9
Moderate OPA (n=1018)	Moderately inactive (n=105, 2.6%) PAL: 1.5	Moderately active (n=648, 16.0%) PAL: 1.6	Active (n=234, 5.8%) PAL: 1.8	Active (n=31, 0.8%) PAL: 2.0
Heavy OPA (n=651)	Moderately active (n=61, 1.5%) PAL: 1.6	Active (n=386, 9.6%) PAL: 1.7	Active (n=190, 4.7%) PAL: 1.9	Active (n=14, 0.3%) PAL: 2.2
Very Heavy OPA (n=108)	Active (n=17, 0.4%) PAL: 1.7	Active (n=50, 1.2%) PAL: 1.8	Active (n=38, 0.9%) PAL: 2.1	Active (n=3, 0.1%) PAL: 2.3

Data are shown as n and %. The number of participants and percentage distribution derives from our study sample. The assigned PAL value derives from Johansson and Westertep,²⁵ who divided energy expenditure obtained from doubly labelled water by the estimated basal metabolic rate of their participants. LPA, leisure time physical activity; OPA, occupational time physical activity; PAL, physical activity level; SGPALS, Saltin-Grimby Physical Activity Level Scale.

MET-hours per week in quartiles in order to assess the validity of ranking physical activity in this PAQ.

The International physical activity questionnaire, sitting question

In this study, the IPAQ short-form sitting question²¹ was employed, asking participants to estimate their average amount of sitting hours on a typical week and weekend day during the last week. In addition to the reported volume, we also grouped sitting hours in quartiles to assess the validity of ranking sitting hours.

Accelerometry data processing

Accelerometry measured physical activity was measured with the triaxial (three planes; axial, coronal and sagittal) ActiGraph wGT3X-BT accelerometer (ActiGraph, LLC, Pensacola, USA), firmware 1.2.0 to 1.8.0. Trained technicians attached the accelerometer to the participants' right hip and instructed them to wear the accelerometer for 24 hours a day on eight consecutive days (the rest of the day following the visit in the clinic and seven more days) and only to remove the accelerometer during water-based activities (eg, showering or swimming) and contact sports. The accelerometer was returned by mail in a prepaid envelope. The ActiLife software (ActiGraph, LLC, Pensacola, USA) was used for initialisation and downloading the data. The accelerometer was initialised for raw data mode with a sampling frequency of 100 Hertz and recordings started at 00:00 the day following the visit in the clinic.

The raw acceleration files were filtered to 10s epochs using the normal (default) proprietary filter in the ActiLife software. The acceleration units are expressed in triaxial vector magnitude (VM) (the square root of the sum of squared activity counts) counts per minute (CPM). We also extracted the number of steps in the accelerometer, which derives from the axial plane in a proprietary algorithm by the manufacturer. The .agd-files (epoch files) were further converted to .csv-files and further analysed in the Quality Control & Analysis Tool software (a custom-made software developed in Matlab: The MathWorks, Inc, Natick, Massachusetts, USA).

The 10s epochs were further aggregated to 60s and an epoch was classified as wear time if two of the following three criteria were fulfilled: (1) an epoch >5VM CPM, (2) if at least two epochs >5VM CPM in the preceding 20 min or (3) at least two epochs >5VM CPM in the following 20 min. Otherwise the acceleration was considered to be noise and classified as non-wear time.³⁶

The triaxial VM CPM cut-points for different intensities are <150 VM CPM for sedentary behaviour³⁷ and ≥2690 VM CPM for moderate and vigorous physical activity

Table 2 Physical activity frequency, intensity and duration (PAFID) questionnaire. Number, MET-values and minutes in parentheses in answering alternatives represents the values for the calculation of MET-hours per week

Frequency (days)	Intensity (METs)	Duration (minutes)
<i>How frequently do you exercise? With exercise, we mean walking, cross-country skiing, swimming or other exercise/sports.</i>	<i>On average, how hard is the exercise?</i>	<i>On average, how long do you exercise?</i>
Never (0)	I take it easy without breaking into a sweat or losing my breath (3 METs)	<15 min (10 min)
Less than once a week (0.5)	I push myself so hard that I break into a sweat and lose my breath (6 METs)	15–29 min (22.5 min)
Once a week (1)	I push myself to near-exhaustion (9 METs)	30–60 min (45 min)
Two to three times per week (2.5)	N/A	>60 min (60 min)
Almost every day (5)	N/A	N/A

METs, metabolic equivalents of tasks.

(MVPA),³⁸ where light physical activity is between 150 to 2689 VM CPM.

Extracted accelerometry measures were volume measures (steps per day and mean VM CPM per day) in addition to intensity measures (minutes per day in sedentary behaviour, light physical activity, MVPA and ≥ 10 min bouts MVPA).

Statistical analyses

We calculated Pearson correlation coefficients to assess the correlation between the PAQs volume outcomes (SGPALS: PAL score, PAFID: MET-hours-week⁻¹, IPAQ sitting: hours spent sitting) and accelerometry outcomes (VM CPM, steps per day, minutes in sedentary behaviour, light physical activity, non-bouted and bouted MVPA) where a coefficient of 0.00 to 0.10, 0.10 to 0.39, 0.40 to 0.69 and ≥ 0.70 was considered a negligible, weak, moderate and strong correlation, respectively.³⁹ Univariate analyses of variance (ANOVA) were performed to assess associations of accelerometry measures (VM CPM, steps, minutes in sedentary behaviour, light physical activity, non-bouted and bouted MVPA) with the SGPALS physical activity ranking, quartiles of MET-hours per week from the PAFID questionnaire and quartiles of reported sitting from the IPAQ. For the IPAQ sitting question, a Bland-Altman plot was created (online supplementary figure 1). The Alpha level was set to 0.05 and data are presented as mean \pm SEM unless otherwise is stated. All data were confirmed to follow normal distribution by visual inspection of residuals when performing the above-mentioned analyses. The analyses were performed overall and in strata of sex, age (10 year groups), BMI (<25, 25 to 29, ≥ 30 kg·m⁻²) and education (primary, high school, <4 years university, ≥ 4 years university). The Statistical Package for Social Sciences (V.25, International Business Machines Corporation, Armonk, New York, USA) was used to perform all statistical analyses.

Patient and public involvement

Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

RESULTS

The descriptive characteristics of the participants wearing the accelerometers and completing the PAQs are presented in table 3.

PAL scores calculated from the SGPALS correlated weakly with VM CPM ($r=0.32$), steps per day ($r=0.27$), sedentary behaviour ($r=-0.20$), light physical activity ($r=0.22$), non-bouted MVPA ($r=0.25$) and bouted MVPA ($r=0.16$) (all $p<0.05$), which was consistent across sex, age, BMI and educational level (all $p<0.05$) (online supplementary table 2). All accelerometry measures increased by increasing rank of self-reported physical activity ($P_{\text{trend}} < 0.001$) (table 4).

Calculated MET-hours per week from the PAFID questionnaire showed negligible correlation with

Table 3 Participant characteristics

SGPALS	Women (n=1983)	Men (n=2057)	Total (n=4040)
Age (yrs)	58.9 \pm 9.5	61.0 \pm 9.9	60.0 \pm 9.7
Height (cm)	164.5 \pm 6.3	177.4 \pm 6.7	171.1 \pm 9.2
Weight (kg)	71.9 \pm 12.8	87.8 \pm 13.8	80.0 \pm 15.5
BMI (kg/m ²)	26.6 \pm 4.7	27.9 \pm 4.0	27.2 \pm 4.4
PAFID	Women (n=3174)	Men (n=2728)	Total (n=5902)
Age (yrs)	63.3 \pm 10.3	63.7 \pm 10.2	63.5 \pm 10.2
Height (cm)	163.6 \pm 6.3	176.9 \pm 6.7	169.8 \pm 9.3
Weight (kg)	71.7 \pm 12.8	87.0 \pm 13.8	78.8 \pm 15.3
BMI (kg/m ²)	26.8 \pm 4.7	27.8 \pm 3.9	27.3 \pm 4.4
IPAQ combined	Women (n=2495)	Men (n=2401)	Total (n=4896)
Age (yrs)	61.4 \pm 10.1	62.6 \pm 10.0	62.0 \pm 10.1
Height (cm)	164.2 \pm 6.2	177.2 \pm 6.6	170.6 \pm 9.2
Weight (kg)	71.5 \pm 12.8	87.1 \pm 13.7	79.1 \pm 15.4
BMI (kg/m ²)	26.5 \pm 4.7	27.7 \pm 3.9	27.1 \pm 4.3

Data are shown as mean \pm SD.

BMI, body mass index; IPAQ combined, International Physical Activity Questionnaire combined: mean of week and weekend; PAFID, Physical Activity Frequency, Intensity and Duration; SGPALS, Saltin-Grimby Physical Activity Level Scale.

accelerometry measured light physical activity ($r=0.06$), weak correlation with VM CPM ($r=0.34$), moderate correlation with steps per day ($r=0.43$) and weak and moderate correlation with non-bouted MVPA ($r=0.39$) and bouted MVPA ($r=0.44$), respectively ($p<0.001$). This was consistent across sex, age, BMI and educational level ($p<0.05$) except for light physical activity, which did not correlate with MET-hours per week in some age groups (40 to 49 years; $p=0.19$, 50 to 59 years; $p=0.13$, 60 to 69 years; $p=0.79$), BMI classifications (<25 kg/m²; $p=0.54$ and 25 to 29 kg/m²; $p=0.31$) and educational levels (high school; $p=0.07$ and university ≥ 4 years; $p=0.051$) (online supplementary table 3).

Quartiles of MET-hours per week from the PAFID questionnaire showed positive association with all accelerometry measures ($P_{\text{trend}} < 0.001$) (table 5).

Accelerometry measured sedentary hours per day correlated weakly with reported sitting hours from the IPAQ sitting question (week day; $r=0.22$, weekend day; $r=0.15$), combined (mean of week and weekend; $r=0.22$, all $p<0.01$), which was consistent across sex, age, BMI and educational level ($p<0.01$) (online supplementary table 4). There was a positive association between quartiles of reported sitting in the IPAQ and accelerometry measured sedentary time ($P_{\text{trend}} < 0.001$) (table 6).

DISCUSSION

We assessed the criterion validity of two PAQs inheriting different physical activity measurement properties (physical activity ranking, volume calculation) and one sedentary

Table 4 The combined leisure time and occupational time SGPALS, and the associations with the accelerometry estimates

n=4040	Inactive (n=349)	Moderately inactive (n=1451)	Moderately active (n=709)	Active (n=1531)
VM CPM*†	410.2±7.8	527.9±4.1	571.3±6.2	618.4±4.6
Steps per day*†	4900.5±107.8	7177.0±71.1	7487.2±103.7	8291.9±73.6
Light physical activity (min·day ⁻¹)*†	360.9±4.8	391.6±2.2	432.0±3.4	425.9±2.2
MVPA (min·day ⁻¹)*†	23.8±1.1	42.3±0.7	43.0±1.1	53.8±0.8
Bouted MVPA (min·day ⁻¹)*†	4.3±0.4	14.1±0.4	12.5±0.6	17.6±0.5

Data are shown as mean±SEM.

*Significant difference between ranks: $p<0.001$.

†Significant linear trend by increasing rank: $p<0.001$.

MVPA, moderate and vigorous physical activity; SGPALS, Saltin-Grimby Physical Activity Level Scale; VM CPM, vector magnitude counts per minute.

time questionnaire, processed as both ranking and volume calculations, against accelerometry as our criterion measure. We found positive associations between ranking of physical activity in both the SGPALS and the PAFID questionnaire, and accelerometry measured physical activity. When processed as calculated volume, we found at best moderate correlations between self-reported and accelerometry measured physical activity. The IPAQ sitting question showed weak correlations and a narrow range in mean accelerometry measured sedentary time between quartile 1 and 4 in the IPAQ (within 1 hour per day).

The validity of the questionnaires

We found positive associations between accelerometry measured physical activity and ranking in the SGPALS. For example, those who categorised themselves in the lowest rank in the combined SGPALS accumulated on average ~4900 steps and 23 min of MVPA per day, respectively, which is about half of the accumulated steps and MVPA per day in the highest rank (~8290 steps and 53 min MVPA). This illustrates the ability of the SGPALS to rank physical activity levels in a large cohort of adults and elderly. The findings of positive associations between SGPALS rankings and accelerometry measured physical activity are consistent with previous validation studies of the SGPALS.^{23 26}

In contrast, when estimating PAL volume scores from the SGPALS, the correlations between PAL scores and accelerometry measured physical activity were weak, which accentuates the biases associated with self-reported physical activity.^{2-4 6} These findings may suggest that the biases associated with self-reported physical activity are more pronounced when physical activity is processed as total volume (eg, PAL, MET-hours per week) compared with ranking individuals according to their self-reported physical activity.

We found positive associations between quartiles of MET-hours per week from the PAFID questionnaire and accelerometry estimates. However, correlations between MET-hours per week from the PAFID questionnaire and accelerometry estimates were weak and only moderate for bouted MVPA. Such correlations are consistent with a previous validation study of the PAFID questionnaire.¹⁸ As with the SGPALS, ranking by quartiles may be the preferred way of expressing self-reported physical activity.

Although we found a positive association between quartiles of reported sitting hours from the IPAQ and accelerometry measured sedentary time, the narrow 1 hour range between quartile 1 and 4 in the IPAQ suggests small differences in real sedentary time between quartiles in the IPAQ.

Table 5 Quartiles of MET-hours per week from the PAFID (n=5902)

Quartiles	1 (n=1355)	2 (n=1498)	3 (n=1473)	4 (n=1576)
Range MET-hours·week ⁻¹	0.00–2.50	2.81–9.00	11.25–11.25	15.00–45.00
MET-hours·week ⁻¹	1.03±0.02	5.13±0.03	11.25±0.00	21.44±0.16
VM CPM*†	448.3±4.4	508.9.5±4.3	557.4±4.3	610.5±4.6
Steps per day*†	5207.8±62.1	6342.9±63.4	7441.3±69.9	8559.4±78.2
Light physical activity (min·day ⁻¹)*†	386.4±2.6	406.2±2.3	407.9±2.3	404.9±2.0
MVPA (min·day ⁻¹)*†	25.9±0.7	35.2±0.7	44.6±0.7	55.8±0.8
Bouted MVPA (min·day ⁻¹)*†	3.9±0.2	9.4±0.3	15.2±0.4	23.7±0.5

Data are shown as mean±SEM.

*Significant difference between quartiles: $p<0.001$.

†Significant trend by increasing quartile: $p<0.001$.

Bouted MVPA, moderate and vigorous physical activity in ≥10 min bouts; MET, metabolic equivalent of tasks; MVPA, moderate and vigorous physical activity; PAFID, Physical Activity Frequency, Intensity and Duration; VM CPM, vector magnitude counts per minute.

**Table 6** Quartiles of reported hours sitting from the IPAQ sitting question, for a typical week and weekend day combined, and the association with accelerometry measured sedentary time

Quartiles	1 (n=783)	2 (n=1432)	3 (n=1277)	4 (n=1359)
Range IPAQ (hours·day ⁻¹)	0.0–4.0	4.0–5.0	6.0–7.0	8.0–24.0
IPAQ (hours·day ⁻¹)	2.8±0.03	4.7±0.03	6.6±0.5	9.7±1.9
Accelerometry sedentary time (hours·day ⁻¹)*†	9.3±0.06	9.6±0.04	9.9±0.04	10.3±0.04

Data are shown mean±SEM.

*Significant difference between quartiles: $p < 0.001$.

†Significant trend by increasing quartile: $p < 0.001$.

IPAQ, International Physical Activity Questionnaire.

Strengths

This study included one of the largest sample sizes in validation studies of PAQs, allowing us to assess the validity in a large heterogeneous sample with high participation rate, which may represent the heterogeneous population to a larger extent than smaller sample sizes. Consequently, the generalisability of the findings from this study is likely high, at least for adults >40 years in western high-income countries.

Limitations

Validation of PAQs is challenging. First of all, in contrast to doubly labelled water, which is the gold standard for measuring free-living energy expenditure,^{40–41} there is no gold standard to measure all aspects (domain, context, intensity, duration, frequency and volume) of physical activity accurately.^{16 42 43}

Second, we used specific cut-points to split intensity in the accelerometry data, which may not reflect the intended intensity by the participants when answering the PAQs. However, in general, accelerometry measured physical activity shows greater validity than self-reported methods when compared with energy expenditure estimated from doubly labelled water,^{44–46} thus, a criterion validation from accelerometry can be considered applicable.

Third, the time periods for self-reported physical activity and sedentary time were not aligned with the accelerometry assessment. However, most physical activity instruments are intended to assess habitual physical activity.⁴⁷ Moreover, as all included questionnaires (SGPALS: Kappa: 0.69,¹⁶ PAFID: Spearman's rho (ρ): 0.76 to 87),¹⁸ IPAQ: ρ : 0.50 to 0.94²¹) and a 7 day accelerometry recording with four valid days (intraclass correlation: 0.8)⁴⁷ are found to provide acceptable reliability, we believe that the included instruments provide reasonable estimates of habitual physical activity and our comparison is justified.

Finally, the waist placement of accelerometers in our study does not assess sitting per se. Other placements, such as thigh-worn accelerometers, may be more suitable for validating self-reported sitting. Nevertheless, our results are consistent with a previous study that employed thigh-worn accelerometers,⁴⁸ suggesting that hip-worn accelerometers are able to measure sedentary time more accurately than self-reported methods.

CONCLUSION

Ranking of physical activity seems to be the preferred method to process PAQs, exhibiting higher validity against accelerometry measures than volume calculations of self-reported physical activity. Self-reported sedentary time poorly reflects accelerometry measured sedentary time. The two PAQs can be used for ranking individuals into different physical activity categories supporting previous studies using these instruments when assessing associations with health outcomes.

Author affiliations

¹Faculty of Health Sciences, School of Sport Sciences, UiT Arctic University of Norway, Tromsø, Troms, Norway

²Faculty of Health Sciences, Department of Community Medicine, UiT Arctic University of Norway, Tromsø, Troms, Norway

³Faculty of Health and Sport Sciences, Department of Sport Science and Physical Education, University of Agder, Kristiansand, Vest-Agder, Norway

⁴MRC Epidemiology Unit, University of Cambridge, Cambridge, UK

⁵Department of Sports Science and Clinical Biomechanics, Faculty of Health Sciences, University of Southern Denmark, Odense, Syddanmark, Denmark

⁶Faculty of Science and Technology, Department of Computer Science, UiT Arctic University of Norway, Tromsø, Troms, Norway

⁷Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Oslo, Norway

⁸Department of Chronic Diseases and Ageing, Norwegian Institute of Public Health, Oslo, Oslo, Norway

Twitter Edvard H Sagelv @edvardhsagelv

Acknowledgements The authors would like to acknowledge PhD Ola Løvsletten for his advice on statistical analyses.

Contributors Significant manuscript writer: EHS. Significant manuscript reviewer/reviser: BM, UE, LAH, SB, BHH, JJ and AH. Concept and design: EHS, BM, UE and LAH. Data acquisition, analysis and interpretation: EHS, BM, LAH, UE, SB, BHH, JJ and AH. Statistical expertise: Ola Løvsletten mentioned in acknowledgements.

Funding The article processing charges are funded by the publication fund at the University Library at UiT the Arctic University of Norway. The work of Edvard H Sagelv and Jonas Johansson are funded by the Population Studies in the High North (Befolkningsundersøkelser i Nord: BiN), an internally funded project by UiT The Arctic University of Norway. The work of Søren Brage is funded by the UK Medical Research Council (MC_UU_12015/3) and the NIHR Biomedical Research Centre in Cambridge (IS-BRC-1215-20014). The remaining authors were funded internally by their institutional tenures/positions.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Tromsø 7 and this present study were approved by the Regional Ethics Committee for Medical Research (REC North ref. 2014/940 and 2016/758410, respectively) and the Norwegian Data Protection Authority.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. The data that support the findings of this study are available from the Tromsø Study but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. The data can be made available from the Tromsø Study upon application to the Data and Publication Committee for the Tromsø Study.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Edvard H Sagelv <http://orcid.org/0000-0003-2213-5806>

REFERENCES

- Knuth AG, Hallal PC. Temporal trends in physical activity: a systematic review. *J Phys Act Health* 2009;6:548–59.
- Sallis JF, Saelens BE. Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport* 2000;71 Suppl 2:1–14.
- Prince SA, Adamo KB, Hamel ME, et al. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56.
- Silbury Z, Goldsmith R, Rushton A. Systematic review of the measurement properties of self-report physical activity questionnaires in healthy adult populations. *BMJ Open* 2015;5:e008430.
- Lee PH, Macfarlane DJ, Lam TH, et al. Validity of the International physical activity questionnaire short form (IPAQ-SF): a systematic review. *Int J Behav Nutr Phys Act* 2011;8:115.
- Helmerhorst HJF, Brage S, Warren J, et al. A systematic review of reliability and objective criterion-related validity of physical activity questionnaires. *Int J Behav Nutr Phys Act* 2012;9:103.
- Lee I-M, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* 2012;380:219–29.
- Wen CP, Wai JPM, Tsai MK, et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011;378:1244–53.
- Lear SA, Hu W, Rangarajan S, et al. The effect of physical activity on mortality and cardiovascular disease in 130 000 people from 17 high-income, middle-income, and low-income countries: the PURE study. *Lancet* 2017;390:2643–54.
- Blair SN. Physical inactivity: the biggest public health problem of the 21st century. *Br J Sports Med* 2009;43:1–2.
- Morris JN, Heady JA, Raffle PA, et al. Coronary heart-disease and physical activity of work. *Lancet* 1953;262:1111–20.
- Carnethon MR, Activity P. Physical activity and cardiovascular disease: how much is enough? *Am J Lifestyle Med* 2009;3:44S–9.
- Haennel RG, Lemire F. Physical activity to prevent cardiovascular disease. how much is enough? *Can Fam Physician* 2002;48:65–71.
- Messerli FH, Ketelhut RG, Ketelhut K. Cardiovascular risk factors and physical activity: how much is enough? *Arch Intern Med* 1999;159:882–3.
- Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes. Comparison with still active athletes of the same ages. *Circulation* 1968;38:1104–15.
- Grimby G, Börjesson M, Jonsdottir IH, et al. The "Saltin-Grimby Physical Activity Level Scale" and its application to health research. *Scand J Med Sci Sports* 2015;25 Suppl 4:119–25.
- Rödger L, Jonsdottir IH, Rosengren A, et al. Self-Reported leisure time physical activity: a useful assessment tool in everyday health care. *BMC Public Health* 2012;12:693.
- Kurtze N, Rangul V, Hustvedt B-E, et al. Reliability and validity of self-reported physical activity in the Nord-Trøndelag health study: Hunt 1. *Scand J Public Health* 2008;36:52–61.
- Ekelund U, Steene-Johannessen J, Brown WJ, et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* 2016;388:1302–10.
- de Rezende LFM, Rodrigues Lopes M, Rey-López JP, et al. Sedentary behavior and health outcomes: an overview of systematic reviews. *PLoS One* 2014;9:e105620.
- Craig CL, Marshall AL, Sjöström M, Sjögren M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381–95.
- Wilhelmsen L, Tibblin G, Aurell M, et al. Physical activity, physical fitness and risk of myocardial infarction. *Adv Cardiol* 1976;18:217–30.
- Emaus A, Degerstrom J, Wilsgaard T, et al. Does a variation in self-reported physical activity reflect variation in objectively measured physical activity, resting heart rate, and physical fitness? results from the Tromsø study. *Scand J Public Health* 2010;38:105–18.
- Graff-Iversen S, Anderssen SA, Holme IM, et al. Two short questionnaires on leisure-time physical activity compared with serum lipids, anthropometric measurements and aerobic power in a suburban population from Oslo, Norway. *Eur J Epidemiol* 2008;23:167–74.
- Johansson G, Westerterp KR. Assessment of the physical activity level with two questions: validation with doubly labeled water. *Int J Obes* 2008;32:1031–3.
- Matthiessen J, Biltoft-Jensen A, Rasmussen LB, et al. Comparison of the Danish physical activity questionnaire with a validated position and motion instrument. *Eur J Epidemiol* 2008;23:311–22.
- Loe H, Rognmo Øivind, Saltin B, et al. Aerobic capacity reference data in 3816 healthy men and women 20–90 years. *PLoS One* 2013;8:e64319.
- Lindén V, Lindén V. Vitamin D and serum cholesterol. *Scand J Soc Med* 1975;3:83–5.
- Thelle DS, Førde OH, Try K, et al. The Tromsø heart study. *Acta Med Scand* 1976;200:107–18.
- Miller NE, Thelle DS, Førde OH, et al. The Tromsø heart-study. high-density lipoprotein and coronary heart-disease: a prospective case-control study. *Lancet* 1977;1:965–8.
- Nordstoga AL, Zotcheva E, Svedahl ER, et al. Long-Term changes in body weight and physical activity in relation to all-cause and cardiovascular mortality: the HUNT study. *Int J Behav Nutr Phys Act* 2019;16:45.
- Jacobsen BK, Eggen AE, Mathiesen EB, et al. Cohort profile: the Tromsø study. *Int J Epidemiol* 2012;41:961–7.
- Wareham NJ, Jakes RW, Rennie KL, et al. Validity and repeatability of a simple index derived from the short physical activity questionnaire used in the European prospective investigation into cancer and nutrition (EPIC) study. *Public Health Nutr* 2003;6:407–13.
- Ainsworth BE, Haskell WL, Leon AS, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993;25:71–80.
- Ainsworth BE, Haskell WL, Herrmann SD, et al. 2011 compendium of physical activities: a second update of codes and Met values. *Med Sci Sports Exerc* 2011;43:1575–81.
- Hecht A, Ma S, Porszasz J, et al. Methodology for using long-term accelerometry monitoring to describe daily activity patterns in COPD. *COPD* 2009;6:121–9.
- Peterson NE, Sirard JR, Kulbok PA, et al. Validation of accelerometer thresholds and inclinometry for measurement of sedentary behavior in young adult university students. *Res Nurs Health* 2015;38:492–9.
- Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors. *J Sci Med Sport* 2011;14:411–6.
- Schober P, Boer C, Schwarte LA. Correlation coefficients: appropriate use and interpretation. *Anesth Analg* 2018;126:1763–8.
- Westerterp KR. Assessment of physical activity: a critical appraisal. *Eur J Appl Physiol* 2009;105:823–8.
- Lifson N, Gordon GB, McClINTOCK R, Mc CR. Measurement of total carbon dioxide production by means of D2O18. *J Appl Physiol* 1955;7:704–10.
- Hills AP, Mokhtar N, Byrne NM. Assessment of physical activity and energy expenditure: an overview of objective measures. *Front Nutr* 2014;1:5.
- Dencker M, Andersen LB. Accelerometer-measured daily physical activity related to aerobic fitness in children and adolescents. *J Sports Sci* 2011;29:887–95.
- Plasqui G, Westerterp KR. Physical activity assessment with accelerometers: an evaluation against doubly labeled water. *Obesity* 2007;15:2371–9.
- Chomistek AK, Yuan C, Matthews CE, et al. Physical activity assessment with the ActiGraph GT3X and doubly labeled water. *Med Sci Sports Exerc* 2017;49:1935–44.
- Bonn SE, Rimm EB, Matthews CE, et al. Associations of sedentary time with energy expenditure and anthropometric measures. *Med Sci Sports Exerc* 2018;50:2575–83.
- Baranowski T, Mâsse LC, Ragan B, et al. How many days was that? we're still not sure. but we're asking the question better! *Med Sci Sports Exerc* 2008;40:544–9.
- Lagersted-Olsen J, Korshøj M, Skotte J, et al. Comparison of objectively measured and self-reported time spent sitting. *Int J Sports Med* 2014;35:534–40.